

1917 2021
104

RHODE ISLAND
MEDICAL JOURNAL



SPECIAL SECTION
CLIMATE CHANGE *and* HEALTH

THEME EDITOR: WILLIAM BINDER, MD

NOVEMBER 2021 VOLUME 104 • NUMBER 9 ISSN 2327-2228



COVERYS[®]

IT'S TIME TO EXPECT MORE FROM YOUR MEDICAL LIABILITY INSURANCE COMPANY.

**More means combining insurance protection
with unique claims analytics and risk management.**

So providers can reduce distractions and focus on improving outcomes.

Coverys is rated A (Excellent)* and has over 45+ years' experience protecting healthcare.
Visit coverys.com.



William Binder, MD

**9 Climate Change and Health:
A Special Edition of the *Rhode Island Medical Journal***

*Examining the impacts of climate change on health,
health care institutions, and mitigation strategies*

WILLIAM BINDER, MD
THEME EDITOR



On the Cover:

The Fox Point Hurricane Barrier's three 40-ft gates can be closed in 30 minutes, preventing storm surges from inundating the low-lying areas of downtown Providence. The barrier was built 1960–1966, and in addition to the Providence River gates, consists of shore dikes with vehicular gates where roads pass through them, canal gates near the power station, and a pumping station.

[PHOTO BY ANDREW E. BINDER]

11 Climate Change and Human Health

NITIN S. DAMLE, MD, MS, MACP

**13 Medical Society Consortium on Climate
and Health Consensus Statement**

**14 Who's at Risk in a Changing Climate?
Mapping Electricity-Dependent
Patient Populations in a Coastal City**

EMMA WEBB, BA
LAKSHMAN BALAJI, BDS, MPH
LARRY A. NATHANSON, MD
SATCHIT BALSARI, MD, MPH
CALEB DRESSER, MD, MPH

**20 Asthma Exacerbations Attributable to
Ozone Air Pollution in New England**


NICHOLAS J. NASSIKAS, MD
KEITH SPANGLER, PhD
GREGORY A. WELLENIUS, ScD

**24 Increased Temperatures Are Associated with
Increased Utilization of Emergency Medical Services
in Rhode Island**

KATELYN MORETTI, MD, MS
BENJAMIN GALLO MARIN, AB, MD'23
LUKE B. SOLIMAN, MTS, MD'23
NICHOLAS ASSELIN, DO
ADAM R. ALUISIO, MD, MSc, DTM&H

**29 Potential Effects of Climate Change on
Tick-borne Diseases in Rhode Island**

HOWARD S. GINSBERG, PhD
JANNELLE COURET, PhD, MEM
JASON GARRETT, BSN, MPH
THOMAS N. MATHER, PhD
ROGER A. LEBRUN, PhD



34 Wasting No Time:
Implementation and the Climate Impact of a Solid Waste Stream Process Intervention in a Large Academic Emergency Department

KATELYN MORETTI, MD, MS
REBECCA KARB, MD, PhD
ROGER DURAND
LEO KOBAYASHI, MD
ALISON S. HAYWARD, MD, MPH

38 Trends in Waste Production at a Community Hospital during the COVID-19 Pandemic

KYLE DENISON MARTIN, DO, MA, MPH, DTM&H
JANE J. CHEN, BS, MD'24
JAMIE THORNDIKE, BS, MD'24
WINSTON MCCORMICK, BS, MD'23
JOHN ROTA, BA
BRIAN BERG, DO
ANNIE DULSKI, DO

43 Trash Talk in the ED:
Takeaways from Waste Audits at New England Hospitals

KYLE DENISON MARTIN, DO, MA, MPH, DTM&H
WINSTON MCCORMICK, BS, MD'23
JULIA CAPACCI, DO, MS
KATELYN MORETTI, MD, MS

45 The Rhode Island Climate Change and Health Program:
Building Knowledge and Community Resilience

RACHEL CALABRO, MS
CAROLINE HOFFMAN, MPH

49 Climate Change and Health in New England:
A Review of Training and Policy Initiatives at Health Education Institutions and Professional Societies


CALEB DRESSER, MD, MPH
EMILY GENTILE, BS
RACHAEL LYONS, BS
KALI SULLIVAN, BS
SATCIT BALSARI, MD, MPH

55 Beyond the Hazard Vulnerability Analysis:
Preparing Health Systems for Climate Change

JOSHUA BAUGH, MD, MPP, MHCM
KATIE KEMEN, MBA
JOHN MESSERVY, AIA
PAUL BIDDINGER, MD

60 Providence's Vulnerability to Floods:
Impacts of Sea Level Rise, Stronger Storms, and Heavier Rainfall

ANDREW E. BINDER, BA
SELIM SUNER, MD, MS
H. CURTIS SPALDING, MPA
ERICH OSTERBERG, PhD



PUBLISHER

RHODE ISLAND MEDICAL SOCIETY

PRESIDENT

ELIZABETH B. LANGE, MD

PRESIDENT-ELECT

THOMAS A. BLEDSOE, MD

VICE PRESIDENT

HEATHER SMITH, MD

SECRETARY

KARA STAVROS, MD

TREASURER

KWAME DAPAAH-AFRIYIE, MD, MBA

EXECUTIVE DIRECTOR

NEWELL E. WARDE, PhD

EDITORS-IN-CHIEF

WILLIAM BINDER, MD

EDWARD FELLER, MD

ASSOCIATE EDITOR

KENNETH S. KORR, MD

EDITOR-IN-CHIEF EMERITUS

JOSEPH H. FRIEDMAN, MD

CO-EDITOR-IN-CHIEF EMERITUS

EDWARD FELLER, MD

PUBLICATION STAFF

MANAGING EDITOR

MARY KORR

mkorr@rimed.org

GRAPHIC DESIGNER

MARIANNE MIGLIORI

ADVERTISING ADMINISTRATOR

DULCINEIA COSME

dcosme@rimed.org

RHODE ISLAND MEDICAL JOURNAL (USPS 464-820), a monthly publication, is owned and published by the Rhode Island Medical Society, 405 Promenade Street, Suite A, Providence RI 02908, 401-331-3207. All rights reserved. ISSN 2327-2228. Published articles represent opinions of the authors and do not necessarily reflect the official policy of the Rhode Island Medical Society, unless clearly specified. Advertisements do not imply sponsorship or endorsement by the Rhode Island Medical Society.

© COPYRIGHT 2013–2021, RHODE ISLAND MEDICAL SOCIETY, ALL RIGHTS RESERVED.



RHODE ISLAND MEDICAL JOURNAL

EMERGENCY MEDICINE RESIDENCY CPC

63 A Case of Heat Stroke in the Era of Climate Change

FRED VARONE, MD

WILLIAM BINDER, MD

CASE REPORTS

70 Complete Resolution of Traumatic Indirect Carotid-Cavernous Fistula Two Years After Inciting Event

JAMES O. ROBBINS, BA, MD-ScM'22

NISARG CHHAYA, MD

JAMIE L. SCHAEFER, MD

73 An Unusual Presentation of Kikuchi-Fujimoto Disease with Recurrent Fever and Syncope

QUYNH-LAN DAO, MD

BRIDGET HOTTENSTEIN, MD

VIJAI RAM SELVARAJ, MD, MPH

IMAGES IN MEDICINE

76 An Incidental Appendix Cystadenoma in a 47-Year-Old Male

WINSTON McCORMICK, BS, MD'23

YIGIT BAYKARA, MD

JESSICA YOON, MD

STEVEN SCHECHTER, MD

PUBLIC HEALTH

79 Adverse Childhood Experiences and Long-Term Health among Adults in Rhode Island

TRACY L. JACKSON, PhD, MPH

KARINE MONTEIRO, MPH

83 Vital Statistics

ROSEANN GIORGIANNI, DEPUTY STATE REGISTRAR

RHODE  ISLAND
MEDICAL JOURNAL




67 SPOTLIGHT

Frozen Obsession documentary follows URI team studying climate change in the Arctic
MARY KORR


85 COMMENTARY

My Newest Favorite Hallucination
JOSEPH H. FRIEDMAN, MD



87 BOOKS

Q&A with Providence College Authors of 'Today's Health Care Issues'
MARY KORR



91 RIMS NEWS

Working for You

RHODE ISLAND MEDICAL JOURNAL



IN THE NEWS

- HOPEHEALTH HULITAR HOSPICE CENTER 94** completes expansion
- FEDERAL DELEGATION 95** announces \$6.1M for capital improvements at RI Community Health Centers
- 95 MIRIAM** receives \$11.1M grant to fund COBRE center on stress, trauma in early life
- 96 ORTHO RHODE ISLAND** now offering robotic technology in total and partial joint replacement

PEOPLE/PLACES



MUKESH K. JAIN, MD 97
appointed Dean of
Medicine and Biological
Sciences at Brown



MEGAN RANNEY, MD 98
appointed Academic Dean at
Brown School of Public Health



MARK DEITCH, MD 98
named VP, CMO
at Miriam

99 KYLE S. NULAND, MD
joins University Orthopedics
Adult Reconstruction Division

99 ROSS BUDACKI, MD
joins University Orthopedics
Sports Medicine division



100 OBITUARIES
Louis A. Corvese, MD
Zhou Weng, MD



Are insurance costs affecting your bottom line?

As record setting malpractice verdicts occur with increasing frequency, insurance premiums for medical practices will continue to rise. It's critical to partner with an experienced insurance broker who can advocate on your behalf to secure the best coverage at the best price.

HUB can help. Our insurance and risk management specialists work with you to develop a strategy to manage your insurance costs — so you can continue to focus on patient care.

hubinternational.com/rimed

Put our resources to work for you.

Daniel Nissi, LIA [508-259-9480](tel:508-259-9480) daniel.nissi@hubinternational.com



Climate Change and Health: A Special Edition of the *Rhode Island Medical Journal*

Examining the impacts of climate change on health, health care institutions, and mitigation strategies

WILLIAM BINDER, MD
THEME EDITOR

The impact of humans on climate has been recognized for centuries. Eighteenth-century Scottish Enlightenment philosopher David Hume postulated that the moderation of Europe's climate, in comparison to Roman times, was due to the gradual advance of cultivation in Europe. He further remarked, "Our northern colonies in America become more temperate, in proportion as the woods are felled..."¹

While Hume could not cite rigorous data as a source for his opinion, scientists and philosophers continued to ponder the impact of humans on their environment. A century after Hume, Irish physicist John Tyndall ushered in the modern era of climate science in 1859 when he discovered the connection between atmospheric CO₂ and the greenhouse effect. By mid-20th century, 200 years after Hume, one scientist remarked, "Human beings are now carrying out a large-scale geophysical experiment of a kind which could not have happened in the past, nor be reproduced in the future. Within a few hundred years we are returning to the air and oceans the concentrated organic carbon stored over hundreds of millions of years."²

Data supporting the impact of humans on atmospheric warming continues to accumulate and this year's Nobel Prize in physics acknowledges the complex interaction between humans, climate, and weather. Rising seas, extreme weather events, and species extinction are no longer versions of apocalyptic science fiction but instead a harsh, modern reality with existential ramifications requiring mitigation and adaptation.

While some skeptics continue to doubt the incontrovertible data, the scientific, medical, and business communities, as well as the US military, have moved past reactionary arguments. In this special issue of the *Rhode Island Medical Journal* (RIMJ), we present an array of articles devoted to the impact of climate change on health in southern New England. Authors from throughout New England and representing multiple health systems, medical schools, universities, primary care practices, the United States Geological Survey, and the Rhode Island Department of Health have all contributed to this important edition of RIMJ.

NITIN S. DAMLE frames the issue by noting that extreme weather events and air pollution have precipitated increasing morbidity and mortality through heat waves, and increases in infectious and respiratory diseases. His commentary also alludes to the societal impact of climate change, as it

can lead to food insecurity, mental health crises, and mass migration. The **RHODE ISLAND MEDICAL SOCIETY'S** consensus statement, signed in March 2020, endorses the perspective of multiple medical societies, and recognizes the impact of climate change on the health of every American.

EMILY WEBB et al., **NICHOLAS J. NASSIKAS** et al., and **KATELYN MORETTI** et al. offer supporting data on the impact of climate change on health in New England. **Webb** details the impact of extreme weather events such as hurricanes and floods – think Hurricanes Sandy, Katrina, and Ida – on electricity-dependent asthmatic patients in Massachusetts. Without electricity, asthmatic patients, as well as others requiring electrically dependent durable medical equipment, are at grave risk. **Nassikas** examines the consequences of pollution on asthma and reports on thousands of excess Emergency Department (ED) visits for summertime ozone-attributable asthma exacerbations across New England. **Moretti** reports on the straightforward association between higher temperatures and increasing Emergency Medical Services' (EMS), and de facto ED, utilization, suggesting that further research may help with planning and resource allocation during summer months. We are also fortunate to feature an article by **HOWARD S. GINSBERG** et al. from the US Geological Survey and the University of Rhode Island (URI) on the effects of climate change on tick-borne disease in Rhode Island and southern New England. The authors document the increasing incidence of Lyme, babesiosis and anaplasmosis in Rhode Island, and note that Lone star ticks, the Gulf Coast tick, and the introduced Asian longhorned tick have been spreading northward into our state. The **USGS/URI group** presents a balanced view, noting that climate change, as well as numerous other environmental and socioeconomic factors, may also contribute to the expanding range of ticks and tick-borne diseases.

The current issue of RIMJ also presents data on mitigation strategies designed to decrease hospital-based waste and reduce emissions. Previous reports suggest that the healthcare industry is responsible for 10% of greenhouse gases and two million tons of waste in the US annually.³ **KATELYN MORETTI** et al. describe a waste mitigation project at Rhode Island Hospital that led to a 23% reduction in solid waste and significant reductions in greenhouse gas emissions. **KYLE DENISON MARTIN** et al. similarly performed a waste audit at Kent Hospital, noting the impact

of the Covid-19 pandemic on waste and energy use in the ED. **Martin** et al. also provide an accompanying commentary regarding important takeaways of waste audits, offering ideas for an environmentally responsible pathway for large health systems.

Paraphrasing the axiom that all politics is local, it can be said that climate change education begins at a local and regional level. **RACHEL CALABRO** and **CAROLINE HOFFMAN** offer an introduction to the Rhode Island Department of Health's Climate Change and Health Program. As part of a US Centers for Disease Control and Prevention initiative, the program has partnered with community groups and state and local agencies to provide educational resources, fund projects fostering community resilience, and offer technical assistance. **CALEB DRESSER** et al. assess the extent of educational activity on a regional basis. In a first-of-its-kind study, they report on the scale of climate and health activities within educational institutions and professional societies in the New England region.

Rounding out this issue is a report by **JOSHUA BAUGH** et al. examining a large health system's sophisticated approach toward a vulnerability analysis, recognizing that previous analyses using historical data is inadequate in an era of rapid climate change and extreme weather events. This Harvard-wide review is a broadly applicable approach for other health systems and should be a model for other institutions moving forward. Recognizing the importance of Baugh's discussion, **ANDREW E. BINDER** et al. provide data and commentary on creating an accurate hazard analysis in order to mitigate the impact of future flooding on Providence's critical health care facilities.

We hope readers appreciate this edition of the RIMJ and that it piques an interest in what is arguably the most salient and pressing issue of our time.

References

1. James Rodger Fleming. Historical Perspectives on Climate Change. 1998; Oxford University Press (NY).
2. Butler CD. Climate Change, Health and Existential Risks to Civilization: A Comprehensive Review (1989-2013). *Int J Environ Res Public Health*. 2018;15(10):2266. Published 2018 Oct 16. doi:10.3390/ijerph15102266.
3. Hsu S, Thiel CL, Mello MJ, Slutzman JE. Dumpster Diving in the Emergency Department: Quantity and Characteristics of Waste at a Level 1 Trauma Center. *West J Emerg Med*. 2020;21:1211-1217.

Theme editor

William Binder, MD, Associate Professor of Emergency Medicine, Warren Alpert Medical School of Brown University; Attending Physician: Rhode Island and Miriam Hospitals, Providence, RI; Editor-in-Chief, *Rhode Island Medical Journal*.

Correspondence

william_binder@brown.edu

Climate Change and Human Health

NITIN S. DAMLE, MD, MS, MACP

In 2015, 186 countries signed the Paris climate agreement to limit global warming to “well below 2 degrees Celsius” and avoid the harmful impacts on human health. Integral to this goal was to reduce carbon emissions to near zero over the next several decades.

Unfortunately, carbon emissions have continued to rise, and the global average temperature has risen by 1.2 degrees Celsius, resulting in the five hottest years on record, all since 2015. The environmental impact is well documented and includes sea level rise, ocean acidification, ice melt, and loss of glacier mass. These changes have resulted in a 46% increase (since 1980) in extreme heat waves, flooding, droughts, forest fires, and intense weather events (hurricanes and cyclones).

The above weather events and air pollution lead to downstream consequences on human health. This article will focus on six health effects.

HEAT WAVES

The number and intensity of heat waves has increased significantly during the past 20 years. This burden is borne mostly by the elderly, children, those with cardiovascular and respiratory diseases, outside laborers, the homeless, and the poor. There has been a 53.7% increase in heat-related mortality for people over the age of 65. Heat waves and temperature rise increase the risk of myocardial infarction, stroke, and acute and chronic renal injury. Studies indicate for each day that the heat index is 95 degrees Fahrenheit (compared to 75 degrees F), emergency department visits increase by 6.6% over the following seven days, heat-related emergency department visits increase by 89% for seven days and death rates increase by 5.8% for each day.

Mitigation is essential and includes early warning systems and response plans, increased access to air conditioning, communication by clinical teams to reach vulnerable populations, and education about the risk of heat illness.

PARTICULATE MATTER AND RESPIRATORY DISEASES

The primary drivers of respiratory diseases are automobile, power plant emissions and forest fires. Measured

particulate matter includes sulfates, nitrates, black carbon, and nitrous oxide.

Forty-three million people in the United States and 92% of the world live in areas in excess of the World Health Organization (WHO) particulate limit, and in 2019 there were eight million deaths worldwide attributable to air pollution.

Increased carbon dioxide production leads to increased growth of allergen-producing weeds, grasses, and trees with rain and floods leading to increased mold and fungal growth. The aeroallergen season has increased from 177 days in the 1970s to 190 days today and is projected to reach 214 days by mid-century. Allergy exacerbations lead to 11 million office visits per year at a cost of \$11.2 billion per year and asthma and COPD exacerbations result in two million emergency department visits, 500,000 hospitalizations, and 3,600 deaths at a cost of \$56 billion per year. Clinicians will need to educate patients about weather trends, knowledge about air pollution, early symptoms, and prevention.

VECTOR-BORNE AND WATER-BORNE DISEASES

The primary vectors in the United States are mosquitoes, sand flies, and ticks. There has been an increase in cases of dengue, Chikungunya, West Nile fever, Zika, and even malaria. The number of cases of Lyme disease has risen from 50,000 in the 1990s to over 400,000 in 2018 and is reported in most regions of the country.

Heavy rains and flooding lead to contamination of water systems, reservoirs, and surface water that increase risk for *E. coli*, *Campylobacter*, *Leptospira*, and *Salmonella* infection. *Vibrio parahaemolyticus* infection is a significant seafood-related public health concern. In addition, rates of parasitic infection have increased, in particular *Cryptosporidium*, *Cyclospora*, and *Giardia Lamblia* in contaminated drinking water and fresh produce. Globally, there are over one billion people that lack access to safe drinking water and 2.5 billion lack access to adequate sanitation, with an estimated five million deaths annually in children.

FOOD SECURITY

The food system involves a network of interactions with our physical and biological environments as food moves from

production to consumption, or from “farm to table.” There has been an increase in crop losses from fungi, bacteria, and viruses. There has been a decrease in food production by 2% per year in the face of a 14% per year increase in global demand, with a projected further decline of 6% in global wheat and 10% in global rice production for each one degree rise in temperature.

The nutrient content (protein, iron, and zinc) has decreased from 20% to 17% with an increase in carbon dioxide concentration from 385 parts per million (ppm) to 450 ppm between 2008 and 2020. Carbon dioxide production is 10 times greater for the consumption of beef, lamb, and pork versus more plant, fish, turkey, and chicken-based diets.

MENTAL HEALTH

The impact on mental health is well established. Forty-nine percent of Hurricane Katrina victims reported increased stress, anxiety, and depression and one in six have been diagnosed with post traumatic stress disorder. Studies show an increased rate of strained interpersonal relationships, substance use disorder, interpersonal aggression, violence, crime, and decreased community cohesion.

There also appears to be a relationship between increased temperature and the number of suicides, and people who meet criteria for mental illness are more vulnerable to the effects of climate change on physical as well as mental health. Much more research is necessary on the impact of climate change and prioritizing mental health benefits with mitigation efforts.

MASS MIGRATION

Extreme weather, sea-level rise, floods, food insecurity, and drought have forced two hundred million people in the world to migrate from their homes between 2008 and 2018. Modeling predicts that one billion people will need to migrate by the end of the century for these same reasons. Without climate change mitigation, this rate will significantly outpace other causes of migration, such as interstate conflict, failure of national governance, and unemployment or underemployment.

CONCLUSION

The effects of climate change have been predicted since the 1970s. There were non-binding agreements in 1979, 1989, and 1997 that did not succeed in slowing carbon dioxide emissions. It is time for the United States to rejoin and strengthen the Paris Agreement of 2015 and actively engage in setting standards at the upcoming UN Climate Change Conference (COP26) this month.

If the United States health sector were a country, it would be the fifth largest emitter on the planet, with \$9 billion

annually in energy costs. A 30% cut in healthcare greenhouse gas emissions by 2030 would prevent an estimated 4,130 premature deaths, 85,000 asthma attacks, four million respiratory symptom events, and 3,750 hospital visit incidents and save about \$1.2 billion in medical costs.

As a physician you have the gravitas to make a significant impact on your community by raising awareness through talks in town halls and community gatherings, in addition to educating your colleagues and patients. You can lobby your local, state, and national representatives to make climate change and health a high priority issue and you can engage in climate-healthy habits, including plant-based diets, electric cars, energy-efficient homes and work environments.

I have painted a sobering outlook with one last opportunity to mitigate the effects of climate change. A recent report by the United Nations Intergovernmental Panel on Climate Change (IPCC) stated that it is possible to limit global warming to 1.5 degrees Celsius, but only with large-scale reductions of all greenhouse gases immediately. If countries move slowly, the Earth could warm by four degrees Celsius (7.2 degrees F) by the end of the century. Experts predict that such a rise would make the Earth uninhabitable. Therefore climate change is the existential threat of our time and now is the moment for all of us to educate and amplify our voices to protect future generations and the planet.

Author

Nitin S. Damle, MD, MS, MACP
Past President, American College of Physicians;
Past President, Rhode Island Medical Society;
Clinical Assistant Professor of Medicine,
Alpert Medical School of Brown University;
Senior Partner, South County Internal Medicine

Correspondence

nsdamle@scim.necoxmail.com

Medical Society Consortium on Climate and Health Consensus Statement

Editor's note: *The following was adopted by the Rhode Island Medical Society in March 2020, along with Affiliates of the Medical Society Consortium on Climate and Health (medsocietiesforclimatehealth.org) regarding the health threats of climate change and the health benefits of solutions (decreasing fossil fuel use).*

We – the undersigned medical societies – support the international scientific consensus, as established in multiple national and international assessments, that the Earth is rapidly warming, and that human actions (especially burning of fossil fuels) are the primary causes. As established in the 2016 U.S. Climate and Health Assessment – The Impacts of Climate Change on Human Health in the United States: A Scientific Assessment – the resulting changes in our climate are creating conditions that harm human health through extreme weather events, reduced air and water quality, increases in infectious and vector-borne diseases, and other mechanisms. While climate change threatens the health of every American, some people are more vulnerable and are most likely to be harmed, including: infants and children; pregnant women; older adults; people with disabilities; people with pre-existing or chronic medical conditions, including mental illnesses; people with low-income; and indigenous peoples, some other communities of color, and immigrants with limited English proficiency.

As medical professionals, many of our members know

firsthand the harmful health effects of climate change on patients. We know that addressing climate change through reduction in fossil fuel use will lead to cleaner air and water, to immediate health benefits for Americans, and will help to limit global climate change.

We support educating the public and policymakers in government and industry about the harmful human health effects of global climate change, and about the immediate and long-term health benefits associated with reducing greenhouse gas emissions (i.e., heat-trapping pollution) and taking other preventive and protective measures that contribute to sustainability. We support actions by physicians and hospitals within their workplaces to adopt sustainable practices and reduce the carbon footprint of the health delivery system.

We recognize the importance of health professionals' involvement in policymaking at the local, state, national, and global level, and support efforts to implement comprehensive and economically sensitive approaches to limiting climate change to the fullest extent possible.

Our organizations are committed to working with officials at all levels to reduce emissions of heat-trapping pollution, and to work with health agencies to promote research on effective interventions and to strengthen the public health infrastructure with the aim of protecting human health from climate change.

Who's at Risk in a Changing Climate?

Mapping Electricity-Dependent Patient Populations in a Coastal City

EMMA WEBB, BA; LAKSHMAN BALAJI, BDS, MPH; LARRY A. NATHANSON, MD;
SATCHIT BALSARI, MD, MPH; CALEB DRESSER, MD, MPH

ABSTRACT

BACKGROUND: Climate change is causing increasingly frequent extreme weather events. This pilot study demonstrates a GIS-based approach for assessing risk to electricity-dependent patients of a coastal academic medical center during future hurricanes.

METHODS: A single-center retrospective chart review was conducted and the spatial distribution of patients with prescriptions for nebulized medications was mapped. Census blocks at risk of flooding in future hurricanes were identified; summary statistics describing proportion of patients at risk are reported.

RESULTS: Out of a local population of 2,101 patients with prescriptions for nebulized medications in the preceding year, 521 (24.8%) were found to live in a hurricane flood zone.

CONCLUSIONS: Healthcare systems can assess risk to climate-vulnerable patient populations using publicly available data in combination with hospital medical records. The approach described here could be applied to a variety of environmental hazards and can inform institutional and individual disaster preparedness efforts.

KEYWORDS: electricity-dependent, mapping, nebulizer, hurricane, climate change, power outage

INTRODUCTION

Climate change, an aging and medically complex population, increasing technological dependence, and fragile public infrastructure present mounting challenges to the health of patients and the operation of healthcare systems.¹⁻³ Health risks from heat waves, hurricanes, flooding, fires, and other hazards are already increasing as result of anthropogenic climate change.⁴ This is of particular concern because an increasing number of patients are reliant on medical devices and a stable supply of electricity, which can be compromised by climate-related hazards and infrastructure failures.⁵⁻⁷

The use of Durable Medical Equipment (DME) has been rising in the United States with the growing prevalence of non-communicable diseases and an aging population.⁸⁻¹⁰ As of 2016, over 2.5 million Medicare patients were dependent

on electrically-powered DME (EDME) including ventilators, oxygen concentrators, and home hemodialysis machines.¹¹ During power outages, EDME-dependent patients (EDPs) may experience device failure or problems continuing their care in the community and may seek assistance at hospitals.^{2,12-14} Many of the patients facing this situation are elderly, have multiple chronic comorbidities, are on multiple medications, or are physically disabled,^{15,16} putting them at elevated risk of immediate or long-term health harm from EDME dysfunction.

Sea level rise, hurricanes, severe storms, wildfires, and heat waves are expected to intensify as a result of climate change and are a substantial threat to power grid stability and access to electricity.^{17,18,19} The frequency of power outages in the US has risen in recent decades because of an increase in the incidence of weather-related outages; severe weather was responsible for 80% of major outages between 2003 and 2012.³ Extreme weather can cause power grid disruptions in a variety of many ways, ranging from the flooding of low-lying electrical infrastructure that occurred during Hurricane Sandy⁷ to increases in demand for electricity in combination with reduced power plant efficiency and production capacity, as seen during heat waves in Southern California.^{5,6,20} Power companies also sometimes initiate Public Safety Power Shutoffs, preemptively turning off power in an effort to reduce risk of wildfire ignition during hot, dry, and windy conditions.²¹

As increasing numbers of patients become dependent on electricity for their healthcare, the value of proactive patient engagement and institutional preparedness for disaster-related threats to the electricity supply will increase, particularly in areas at high risk of outages and with high numbers of EDPs. Information on this population and the hazards it faces are crucial to these initiatives, and efforts to map EDPs at national, state, and municipal levels using insurance claims data, medical record searches, and geographic information systems (GIS) are ongoing.^{1,11,22-24} The most comprehensive existing tool is the Department of Health and Human Services emPOWER system, which maps EDME users at the zip code level based on Medicare claims and can provide address-level information to EMS agencies seeking to reach specific patients during a disaster.¹¹ However, the tool does not provide information about where patients obtain healthcare, excludes patients covered under other

insurance providers or who have no insurance, and does not allow for disaggregated analysis at the municipal or institutional level.^{11,24}

In this article we present a complementary, locally actionable approach that utilizes retrospective medical records review in combination with publicly available hazard data to assess the vulnerability of institution-specific EDP populations to climate-related natural hazards. For purposes of this pilot study, we focused on home nebulizers as a representative EDME and hurricane-related coastal flooding as a representative climate-related hazard.

METHODS

This study consisted of a retrospective chart review followed by spatial aggregation, mapping, and analysis of presumed nebulizer users in relation to publicly available data describing hurricane inundation zones. Nebulizers were selected as the representative EDME because they are commonly used, use can be inferred from prescription information, and device failure can contribute to adverse clinical sequelae. Inundation zones were chosen because of the strong relationship between flooding and electrical outages which has been demonstrated in recent hurricane events.^{3,6,7}


All medical records in the hospital online medical record (OMR) were queried for the presence of prescriptions for nebulized medications within the preceding year (10/25/2019 to 10/25/2020). Duplicate records from patients with multiple prescriptions within the time period were removed. Home addresses were converted to spatial coordinates via an application programming interface (API), and records were then aggregated to the level of census blocks to create the de-identified dataset used for this analysis. Hurricane Surge Inundation Zone data from the U.S. Army Corps of Engineers was obtained from MassGIS and overlaid with the frequency maps of nebulizer users.²⁵ A clip function was used to restrict the block-level nebulizer frequency maps to areas of overlap with hurricane inundation zones and quantify the population of patients with recent prescriptions for nebulized medication who would be exposed to storm surge inundation during a hurricane of a given Saffir-Simpson category. Attribute tables were exported and the sum and proportion of the total nebulizer population at risk and the increase in number and percent of patients expected with each increase in hurricane category were computed. Maps of nebulizer prescription frequency were created using 2010 US Census geographic data and are shown at the census tract level to protect patient anonymity.

Conversion of addresses to geo-coordinates was performed via the Google Maps API implemented in the gmap package in R v3.6.1.^{26,27} Spatial analysis and mapping was performed in ArcMap²⁸; summary statistics were computed in MS Excel.²⁹ This study was reviewed and determined to be exempt by the BIDMC Institutional Review Board.

RESULTS

A total of 1,384,134 prescription entries for 185,906 patients were screened, from which a list of 3,649 nebulizer prescriptions was compiled. Of these, 1,417 duplicates (e.g. multiple prescriptions for the same patient during the study time-frame) were excluded, leaving 2,232 unique patient records. Seven records could not be successfully geocoded and were removed from the dataset, leaving 2,225 records representing presumed nebulizer users, from which the 2,101 patients residing in Massachusetts were included in the analysis of local vulnerability to hurricane inundation, as summarized in **Figure 1**.

Figure 1. Medical records screening and inclusion process. All records are from a single medical center during the period 10/25/2019 to 10/25/2020.




A total of 30 different medications were prescribed, including albuterol sulfate (1,237 patients), ipratropium-albuterol (372), budesonide (332), and other medications or combinations of medications (284). Most patients resided in Massachusetts (2,101), followed by New Hampshire (45), Maine (16), Rhode Island (14), and other states (49). Within Massachusetts, 730 lived in Suffolk County, 512 in Middlesex County, and 859 in other counties. See **Table 1**.

Within Massachusetts, 1,886 census blocks in 794 census tracts contained at least 1 presumed nebulizer user. All census tracts containing more than 10 patients were located in

Table 1. Study population characteristics, including locations of residence and nebulized prescriptions information.

	Number	Proportion (%)
Patients	2,225	100%
Prescription Type		
Albuterol	1,237	55.6 %
Ipratropium-Albuterol	372	16.7 %
Budesonide	332	14.5 %
Other / Combination	284	12.8 %
State of Residence		
Massachusetts	2,101	94.4 %
New Hampshire	45	2.0 %
Maine	16	0.7 %
Rhode Island	14	0.6 %
Other	49	2.2 %

Figure 2. Distribution of presumed nebulizer users in and around Boston, MA. Darker coloration corresponds to a larger number of presumed nebulizer users residing within the census tract.




Dorchester, Saugus, and Revere. The spatial distribution of nebulizer users at the census tract level is shown in **Figure 2**.

At the block level, 521 presumed nebulizer users resided in census blocks that overlapped with hurricane flood zones. This population represents 24.8% of patients who received care at the study site during the study timeframe, were nebulizer users, and lived in Massachusetts. Subsets of this population living in each hurricane inundation zone were computed; stratified results are presented in **Table 2**. A total of 186 presumed nebulizer users (8.9%) reside in a zone that would flood in a Category 1 hurricane, and 372 (17.7%) reside in a zone that would flood in a Category 2 hurricane; the population of interest at risk of inundation in a Category 2 storm is approximately twice that at risk in a Category 1 storm. The spatial distribution of nebulizer users living in locations at risk of hurricane-related flooding is shown in **Figure 3**; data is presented at the census tract level to preserve anonymity.

Table 2. Population of presumed nebulizer users who received care at the study site during the study period and are at risk of inundation of their dwelling location in Category 1 through Category 4 hurricanes.

	Category 1 zone	Category 2 zone	Category 3 zone	Category 4 zone
Presumed nebulizer users at risk of flooding in this zone (n)	186	372	456	521
Proportion of Massachusetts nebulizer users (n=2,101) at risk of flooding in a storm of this category (%)	8.9	17.7	21.7	24.8
Additional patients at risk relative to next lowest category hurricane (n)	-	186	84	65
Proportional increase in number of patients at risk relative to next lowest category hurricane (%)	-	100%	22.6%	14.3%

Figure 3. Distribution of presumed nebulizer users residing in hurricane inundation zones. Darker red indicates a larger number of presumed nebulizers who may be at risk of inundation during a major hurricane.



DISCUSSION

This pilot study demonstrates a simple, reproducible methodology for mapping climate-related risk to electricity-dependent patients. Our approach, which uses publicly available hazard data in combination with internal medical records data, is not technologically complex and, thanks to the widespread use of electronic medical records, could be implemented by many healthcare institutions.

Our results show that nearly one-quarter of Massachusetts nebulizer users who received care at the study site are living in census blocks at risk of flooding during hurricanes, and that the number at risk approximately doubles in a Category 2 hurricane in comparison to a Category 1 hurricane. At the institutional level, this suggests there could be an increase in respiratory presentations in the hours or days following

a hurricane, such information can help guide preparedness efforts. In addition, awareness of this risk may help primary care providers, pulmonologists, and patients make preparations so that alternative breathing treatments, sources of electricity, or temporary accommodations are available in the event of a major storm.

It was also noteworthy that the highest densities of presumed nebulizer users at risk of inundation in this study were in Dorchester, Saugus, and Revere – communities with substantial minority, historically marginalized, or financially constrained populations that may face substantial barriers to climate change adaptation. Such communities may benefit from hospital outreach efforts including collaboration with local governments or civil society organizations on community-based interventions to address these and other health risks.

This pilot study demonstrates that individual healthcare organizations already possess or can access the data needed to perform basic evaluations of the risk to their high-vulnerability patient populations posed by climate-related natural hazards. Practical information on the hazard exposure of vulnerable populations can be obtained using freely available datasets and analytical tools in combination with medical records data. Improving electronic medical records utilization, open-source statistical and geographic information systems, and availability of free training on their use mean that simple analyses should be within the reach of many institutions. While true real-time translational data readiness for disaster response remains an urgent and complex challenge,³⁰ static pre-disaster risk assessments such as the pilot results presented here can be performed at the level of individual institutions. Availability of such information may enhance engagement with community resilience and climate change adaptation efforts and could be expanded to a wide range of patient populations, hazards, and infrastructure issues.

At the organizational level, this information could contribute to forecasting patient surges and anticipating acute care needs. For example, Emergency Department (ED) demand forecasting is often done hour-to-hour³¹⁻³⁴, the information and methods outlined here could be extended to allow organizations to predict which locations could receive a higher volume of EDME-related ED arrivals based on proximity to high-risk areas and vulnerable populations during disaster events. Community resilience efforts may also benefit from information on the hazard exposure of specific high-risk patient populations; such work may help avoid ED crowding related to DME problems and exacerbations of chronic conditions during disasters, which can lead to operational challenges, delays in care, and increased ED length of stay.³⁵⁻³⁸

Information on exposure to climate-related hazards can also inform care in the outpatient setting. Outpatient providers may find such information useful when counseling patients on issues such as respiratory health, heat wave

safety, and disaster preparedness. Home Health Care (HHC) presents another intervention opportunity³⁹; HHC organizations are uniquely positioned to assist EDME users with disaster preparedness at home – for example, using the National Association for Homecare and Hospice Emergency Preparedness Checklist.⁴⁰

Analyses of climate change hazards in relation to vulnerable populations will be of growing importance in coming decades. There is a particularly urgent need to develop methods to proactively identify populations at high risk during disaster-related electrical outages in order to reduce suffering, avoid long-term health impacts associated with device failure, anticipate and reduce ED surges, and ease strain on hospital systems. Healthcare institutions have a responsibility to prepare for anticipated hazards that can affect their ability to provide care, including the risk of patient surges during extreme events. Healthcare providers are uniquely positioned to counsel patients on preparedness for natural hazards as a means of protecting their health and advocate for measures to adapt to climate change and protect health in the community. Information on local climate-related risk profiles can help inform such conversations.

LIMITATIONS

The population in this analysis is limited to nebulizer users and does not include persons dependent on other forms of EDME, which may have different spatial distributions. This medical records review was limited to a single medical center, and provides information specific to the hospital under assessment, rather than a portrait of the region as a whole. Our analysis infers that inundation from a hurricane is a predictor of electrical outage based on a variety of case studies and other evidence,^{3,6,7} but lack of recent hurricanes means the degree of this correlation in the study region is not definitively known. In addition, quantifying the population at risk on the basis of residence within an inundation zone may underestimate the true number at risk, as damage to electrical infrastructure within flood zones may result in power outages affecting non-inundated areas, as occurred in large parts of Manhattan following a transformer failure during Hurricane Sandy.^{6,7} Finally, a variety of factors may affect whether inundation actually harms electricity-dependent patients, including locations and fortifications of power stations and transmission lines, availability of public sites at which power is available, such as libraries or community centers, and whether or not a household is on a priority power restoration list or executes a pre-disaster evacuation plan.

CONCLUSION

Extreme weather related to climate change and the use of electricity-dependent medical devices are intersecting issues that combine to increase the risk of adverse health outcomes

resulting from weather-related electrical outages. This pilot study uses publicly available hazard data and internal medical records data to show that a quarter of study-site patients who use nebulizers reside in locations that are expected to be inundated during a major hurricane and may experience electrical outages. Our findings suggest preparedness opportunities at both the population and institutional levels. The methodology necessary to conduct this type of assessment is not complex and could be reproduced in other settings to assess a variety of hazards and patient populations.

References

- Molinari NA, Chen B, Krishna N, Morris T. Who's at Risk When the Power Goes Out? The At-home Electricity-Dependent Population in the United States, 2012. *J Public Health Manag Pract.* 2017;23(2):152-9.
- Mokdad AH, Mensah GA, Posner SF, Reed E, Simoes EJ, Engelgau MM. When chronic conditions become acute: prevention and control of chronic diseases and adverse health outcomes during natural disasters. *Prev Chronic Dis.* 2005;2 Spec no(Spec No):A04.
- Kenward A, Raja U. Blackout: Extreme Weather, Climate Change, and Power Outages. *Climate Central*; 2014.
- USGCRP, 2018: Impacts, Risks, and Adaptation in the United States: Fourth National Climate Assessment, Volume II [Reidmiller, D.R., C.W. Avery, D.R. Easterling, K.E. Kunkel, K.L.M. Lewis, T.K. Maycock, and B.C. Stewart (eds.)]. U.S. Global Change Research Program, Washington, DC, USA, 1515 pp. doi: 10.7930/NCA4.2018.
- Miller NL, Hayhoe K, Jin J, Auffhammer M. Climate, Extreme Heat, and Electricity Demand in California. *Journal of Applied Meteorology and Climatology.* 2008;47(6):1834-44.
- Davis M, Clemmer S. Power Failure: How Climate Change Puts Our Electricity at Risk— and What We Can Do. [Internet] 2014. [Available from: <https://www.ucsusa.org/resources/power-failure>]
- Kleinbaum J. Hurricane Sandy Plunges Lower Manhattan into Darkness [Internet]: NBC New York, 2012 [Available from: <https://www.nbcnewyork.com/news/local/lower-manhattan-plunged-into-darkness-during-historic-storm/1966520/#:~:text=In%20all%2C%20Sandy%20left%20about,or%20flying%20debris%2C%20Miksd%20said>].
- Raghupathi W, Raghupathi V. An Empirical Study of Chronic Diseases in the United States: A Visual Analytics Approach. *Int J Environ Res Public Health.* 2018;15(3):431.
- LaPlante MP, Kaye HS. Demographics and trends in wheeled mobility equipment use and accessibility in the community. *Assist Technol.* 2010;22(1):3-17.
- Seeman TE, Merkin SS, Crimmins EM, Karlamangla AS. Disability trends among older Americans: National Health And Nutrition Examination Surveys, 1988-1994 and 1999-2004. *Am J Public Health.* 2010;100(1):100-7.
- U.S. Department of Health and Human Services. HHS emPOWER Map 3.0 [Internet] 2020. Available from: <https://empowermap.hhs.gov/#:~:text=Over%202.5%20million%20Medicare%20beneficiaries,life%2Dthreatening%20for%20these%20individuals>.
- Howard D, Zhang R, Huang Y, Kutner N. Hospitalization rates among dialysis patients during Hurricane Katrina. *Prehosp Disaster Med.* 2012;27(4):325-9.
- Greenwald PW, Rutherford AF, Green RA, Giglio J. Emergency department visits for home medical device failure during the 2003 North America blackout. *Acad Emerg Med.* 2004;11(7):786-9.
- Arrieta MI, Foreman RD, Crook ED, Icenogle ML. Providing continuity of care for chronic diseases in the aftermath of Katrina: from field experience to policy recommendations. *Disaster Med Public Health Prep* [Internet]. 2009;3(3):174-82.
- AARP Public Policy Institute. Chronic Conditions Among Older Americans. https://www.aarp.org/health/medicare-insurance/info-03-2009/beyond_50_hcr.html; 2009.
- Hayes BD, Klein-Schwartz W, Barrueto F, Jr. Polypharmacy and the geriatric patient. *Clin Geriatr Med.* 2007;23(2):371-90, vii.
- Oppenheimer M, Glavovic BC, Hinkel J, Wal Rvd, Magnan AK, Abd-Elgawad A, et al. Sea Level Rise and Implications for Low-Lying Islands, Coasts and Communities. IPCC Special Report on the Ocean and Cryosphere in a Changing Climate: Intergovernmental Panel on Climate Change; 2019.
- Collins M, Sutherland M, Bouwer L, Cheong S-M, Frölicher T, Combes HJD, et al. Extremes, Abrupt Changes and Managing Risk. IPCC Special Report on the Ocean and Cryosphere in a Changing Climate: Intergovernmental Panel on Climate Change; 2019.
- Hoegh-Guldberg O, Jacob D, Taylor M, Bindi M, Brown S, Camilloni I, et al. Impacts of 1.5°C Global Warming on Natural and Human Systems. Global Warming of 1.5°C. An IPCC Special Report on the impacts of global warming of 1.5°C above pre-industrial levels and related global greenhouse gas emission pathways, in the context of strengthening the global response to the threat of climate change, sustainable development, and efforts to eradicate poverty; 2019.
- Wilson W. Constructed Climates: A Primer on Urban Environments. 2011. Figure 6.6, electricity supplied by southern California in 2004 vs. maximum daily temperature. In: *Constructed Climates: A Primer on Urban Environments* [Internet]. The University of Chicago Press. Available from: <http://people.duke.edu/~wgv/ConClim/chap-6-human-health-urban-inequities/6-6-air-conditioning.html>.
- Pacific Gas & Electric. Learn about Public Safety Power Shutoff (PSPS) events. Date Unknown. Available from: https://www.pge.com/en_US/residential/outages/public-safety-power-shutoff/learn-about-psps.page.
- Xi N, Wallace R, Agarwal G, Chan D, Gershon A, Gupta S. Identifying patients with asthma in primary care electronic medical record systems Chart analysis-based electronic algorithm validation study. *Can Fam Physician.* 2015;61(10):e474-83.
- Wilson JL, Neuffer E. Estimating Medically Fragile Population Exposures to Tropical Storm Surges. *International Journal of Geospatial and Environmental Research.* 2015;2(1).
- DeSalvo K, Lurie N, Finne K, Worrall C, Bogdanov A, Dinkler A, et al. Using Medicare data to identify individuals who are electricity dependent to improve disaster preparedness and response. *Am J Public Health.* 2014;104(7):1160-4.
- Massachusetts Bureau of Geographic Information. MassGIS Data: Hurricane Surge Inundation Zones 2013 [Available from: <https://www.mass.gov/info-details/massgis-data-hurricane-surge-inundation-zones#downloads>].
- Kahle D, Wickham H. ggmap: Spatial Visualization with ggplot2. *The R Journal,* 5(1), 144-161. URL <http://journal.r-project.org/archive/2013-1/kahle-wickham.pdf>
- R Core Team (2021). R: A language and environment for statistical computing. R Foundation for Statistical Computing, Vienna, Austria. URL <https://www.R-project.org/>.
- ESRI. Resources for ArcMap [date unknown] [Available from: <https://www.esri.com/en-us/arcgis/products/arcgis-desktop/resources>].
- Microsoft. Microsoft 365 Excel [date unknown] [Available from: <https://www.microsoft.com/en-us/microsoft-365/excel>].
- Balsari S, Kiang M, Buckee C. Data in Crisis – Rethinking Disaster Preparedness in the United States. *NEJM.* September 1, 2021 DOI: 10.1056/NEJMms2104654
- McCarthy ML, Zeger SL, Ding R, Aronsky D, Hoot NR, Kelen GD. The challenge of predicting demand for emergency department services. *Acad Emerg Med.* 2008;15(4):337-46.

32. Hoot NR, Leblanc LJ, Jones I, Levin SR, Zhou C, Gadd CS, et al. Forecasting emergency department crowding: a prospective, real-time evaluation. *J Am Med Inform Assoc.* 2009;16(3):338-45.
33. Chase VJ, Cohn AE, Peterson TA, Lavieri MS. Predicting emergency department volume using forecasting methods to create a "surge response" for noncrisis events. *Acad Emerg Med.* 2012;19(5):569-76.
34. Asheim A, Bache-Wiig Bjornsen LP, Naess-Pleym LE, Uleberg O, Dale J, Nilsen SM. Real-time forecasting of emergency department arrivals using prehospital data. *BMC Emerg Med.* 2019;19(1):42.
35. Platz E, Cooper HP, Silvestri S, Siebert CF. The impact of a series of hurricanes on the visits to two central Florida Emergency Departments. *J Emerg Med.* 2007;33(1):39-46.
36. Lane K, Charles-Guzman K, Wheeler K, Abid Z, Graber N, Matte T. Health effects of coastal storms and flooding in urban areas: a review and vulnerability assessment. *J Environ Public Health.* 2013;2013:913064.
37. Saulnier DD, Brolin Ribacke K, von Schreeb J. No Calm After the Storm: A Systematic Review of Human Health Following Flood and Storm Disasters. *Prehosp Disaster Med.* 2017;32(5):568-79.
38. Morley C, Unwin M, Peterson GM, Stankovich J, Kinsman L. Emergency department crowding: A systematic review of causes, consequences and solutions. *PLoS One.* 2018;13(8):e0203316.
39. Rest KD, Hirsch P. Insights and decision support for home health care services in times of disasters. *Cent Eur J Oper Res.* 2021:1-25.
40. The National Association for Homecare and Hospice. Emergency Preparedness Packet for Home Health Agencies. https://www.nahc.org/wp-content/uploads/2017/10/EP_Binder.pdf2008.

Acknowledgment

This project was supported by the CrisisReady Initiative at Harvard and Direct Relief. Dr. Caleb Dresser's work was also supported by the Climate and Health Foundation.

Authors

Emma Webb, BA, PA/MPH Candidate, Tufts University School of Medicine.
 Lakshman Balaji, BDS, MPH, Biostatistician at Beth Israel Deaconess Medical Center.
 Larry Nathanson, MD, Assistant Professor of Emergency Medicine at Harvard Medical School, Director of Emergency Medicine Informatics at Beth Israel Deaconess Medical Center.
 Satchit Balsari, MD, MPH, Assistant Professor of Emergency Medicine at Harvard Medical School.
 Caleb Dresser, MD, MPH, Instructor of Emergency Medicine at Harvard Medical School.

Disclaimer

The views expressed herein are those of the authors and do not necessarily reflect the views of Beth Israel Deaconess Medical Center or Harvard Medical School.

Correspondence

Caleb Dresser, MD, MPH
 Department of Emergency Medicine
 1 Deaconess Road
 Boston, MA 02215
cdresser@bidmc.harvard.edu

Asthma Exacerbations Attributable to Ozone Air Pollution in New England

NICHOLAS J. NASSIKAS, MD; KEITH SPANGLER, PhD; GREGORY A. WELLENIUS, ScD

ABSTRACT

BACKGROUND: Ground-level ozone (O_3) is an air pollutant and known trigger for asthma exacerbation. We sought to estimate the number of summertime emergency department (ED) visits for asthma exacerbations attributable to ozone in each county in New England (Rhode Island, Massachusetts, Connecticut, New Hampshire, Maine, and Vermont) in 2010.

METHODS: We performed a health impact assessment using BenMAP. We used population and incidence rates in New England, daily maximum 8-hour O_3 levels, and a concentration-response function derived from the epidemiological literature to quantify ozone-attributable asthma ED visits.

RESULTS: We estimate that in 2010 there were 4,612 (95% CI 2192, 6866) excess ED visits for asthma exacerbation attributable to summertime ozone across New England. Rates of ozone-attributable asthma ED visits were highest in Connecticut and Massachusetts.

CONCLUSIONS: There was a substantial number of ozone-attributable asthma ED visits in New England in 2010 with geographic heterogeneity across states and counties.

KEYWORDS: climate change, ozone, asthma, emergency department visit

INTRODUCTION

Asthma affects 1.3 million people in the New England region of the United States.¹ Exposure to pollen, dust, infections, and air pollutants can exacerbate symptoms of the disease. One air pollutant – tropospheric ozone (O_3), or ground-level O_3 – is a well-described trigger for asthma exacerbations.² Tropospheric O_3 is formed by the reaction of nitrogen oxides and volatile organic compounds in the presence of sunlight and heat and, unlike stratospheric ozone, which provides protection from ultraviolet radiation from the sun, ground-level ozone is harmful to human health. Tropospheric O_3 concentrations tend to be highest during the summer months when temperatures are higher and sun exposure is prolonged. One study estimated that approximately 8–21%

of asthma emergency room visits in the United States are associated with O_3 exposure,³ although the incidence varies geographically. Although ozone is recognized as a contributor to unhealthy air and an important trigger of asthma exacerbations across New England,⁴ the burden of ozone air pollution on asthma across all states in New England has not been established. Accordingly, we sought to estimate the burden of disease in terms of summertime emergency department (ED) visits for asthma exacerbations attributable to ozone in every county in Rhode Island, Massachusetts, Connecticut, New Hampshire, Maine, and Vermont in 2010.


METHODOLOGY

We performed a health impact assessment to estimate the number of ozone-attributable asthma exacerbations in New England among all ages using the United States Environmental Protection Agency Benefits Mapping and Analysis Program (BenMAP), similar to prior studies.⁵ This approach requires multiple data inputs including estimates of ozone concentrations, baseline incidence rates of asthma-related ED visits, population counts, and a defined concentration-response function providing estimates of the association between ozone and asthma exacerbation. Conditional on these inputs (described below), we then quantified ozone-attributable asthma exacerbation ED visits on a county-level as follows (Equation 1):

$$Y = Y_0 \cdot (1 - e^{-\beta \cdot \Delta O_3}) \cdot Pop$$

where Y is the number of ozone-attributable asthma exacerbation ED visits, Y_0 is the baseline incidence rate for asthma, ΔO_3 is the concentration of daily maximum 8-hour O_3 , Pop is the population (age 0-99), and β is the risk coefficient on a natural log scale for asthma ED visits associated with daily O_3 exposure. County-level age-adjusted baseline incidence rates for asthma ED visits in 2010 are obtained from the National Environmental Public Health Tracking Network.⁶⁻⁸ Daily maximum 8-hour ozone concentrations during the summertime, defined as May 1 to September 30, 2010 in New England are based on air pollution monitor stations within each state. BenMAP calculates an inverse-distance weighted average of nearby monitors to determine county-level ozone concentrations, an approach called the Voronoi Neighbor Averaging algorithm.⁹ County-level population numbers are

Figure 1. Panel A: Average summertime (May 1–September 30), county-level ozone concentrations (parts per billion) in New England states in 2010. Panel B: Age-adjusted baseline asthma emergency department visit (EDV) rates per 10,000 population in 2010 on a county-level.⁸ Panel C: Rates of ozone-attributable asthma ED visits per 10,000 people in New England counties in 2010.



based on the 2010 U.S. Census. The risk coefficient is based on the meta-analysis by Ji et al. (2011) that reports a 2.90% (95% CI 1.33, 4.50) increase in risk of ED visit for asthma exacerbation for every 10 part per billion (ppb) increase in daily maximum 8-hour O₃.¹⁰

RESULTS

In the New England states, warm season averages of daily maximum 8-hour O₃ concentrations between May and September 2010 ranged from 34–47 ppb and were highest in New London County and Middlesex County in Connecticut (Figure 1). Aroostook County in Maine had the lowest concentrations of O₃.

We estimate that in 2010 there were 4,612 (95% CI 2192, 6866) excess ED visits for asthma exacerbation attributable to summertime ozone exposure in New England. The average rate of excess asthma-related ED visits was 3.2 (95% CI 1.52, 4.75) per 10,000 population. Connecticut and Massachusetts had the highest rates of ozone-attributable asthma ED visits per 10,000 population, with 3.65 (95% CI 1.74, 5.41) and 3.40 (95% CI 1.62, 5.06) respectively, while Vermont had the lowest at 1.98 (95% CI: 0.94, 2.96) (Figure 1). Rates of ozone-attributable asthma ED visits per 10,000 population varied by county, ranging from 1.5 to 5.9 (Table 1). In terms of absolute numbers rather than rates, Massachusetts had the highest absolute number of ozone-attributable asthma ED visits on a state-level in

2010 and Vermont had the lowest at 2,226 (95% CI 1058, 3314) and 123 (95% CI 59, 185), respectively.

DISCUSSION

Ozone concentrations and the attributable asthma ED visits varied geographically across New England. The health effects of ozone on asthma vary in part due to differences in baseline asthma incidence rates between counties. Indeed, a few counties such as Washington County, Maine, have high rates of asthma ED visits at baseline that would contribute to relatively higher rates of ozone-related asthma ED visits despite relatively lower summertime ozone concentrations.

Table 1. Counties in New England with the highest and lowest rates of summertime ozone-attributable asthma ED visits in 2010.

Counties with highest number of ozone-attributable asthma ED visits		Counties with lowest number of ozone-attributable asthma ED visits	
County, State	Asthma ED visit rate per 10,000 (95% CI)	County, State	Asthma ED visit rate per 10,000 (95% CI)
Nantucket County, MA	5.89 (2.8,8.76)	Chittenden County, VT	1.49 (0.7,2.22)
Washington County, ME	4.79 (2.27,7.17)	Orange County, VT	1.61 (0.76,2.4)
Hampden County, MA	4.41 (2.1,6.57)	Lamoille County, VT	1.62 (0.76,2.41)
Dukes County, MA	4.27 (2.03,6.35)	Washington County, VT	1.8 (0.85,2.7)
New London County, CT	4.23 (2.02,6.28)	Caledonia County, VT	1.81 (0.86,2.7)
Tolland County, CT	4.17 (1.99,6.19)	Addison County, VT	1.97 (0.93,2.94)
Aroostook County, ME	4.11 (1.94,6.16)	Grand Isle County, VT	1.98 (0.94,2.96)
Somerset County, ME	3.0 (1.42,4.49)	Sagadahoc County, ME	2.16 (1.02,3.22)
Berkshire County, MA	3.92 (1.86,5.85)	Cheshire County, NH	2.17 (1.03,3.25)
Hartford County, CT	3.91 (1.86,5.81)	Windsor County, VT	2.18 (1.03,3.26)

Differences in ozone concentrations also explain the geographic heterogeneity of the health impacts on asthma and likely stem from the distribution of precursor emissions of nitrogen oxides and volatile organic compounds as well as sunlight and heat. While ozone concentrations in New England in 2010 were, on average, below the current Ozone National Ambient Air Quality Standard, set at 70ppb¹¹ there remained a substantial number of ED visits attributable to ozone. Prior studies have shown that reducing ozone concentrations below the national standard would lead to fewer deaths and fewer respiratory hospitalizations.¹²

In the future, climate change will have significant implications for asthma in New England, in part due to the influence of heat on ozone formation.¹³ The Northeast is projected to have the fastest warming in the contiguous US, increasing 3°C by 2050 under some models.¹⁴ There is also emerging evidence describing a synergistic effect of heat on air pollution and human health.¹⁵ While our study estimates ozone-attributable asthma ED visits based on ozone concentrations in 2010, temperature increases in the future due to climate change will lead to more ozone formation while precursor emission levels are expected to decrease,¹⁶ and may inform adaptation and mitigation strategies.

One of the strengths of this study is the spatial scale. We report ozone-attributable asthma ED visits on a state and county level. Understanding the local burden of ozone on asthma can provide useful information for policy-makers. Another strength is the use of local baseline incidence rates for asthma ED visits rather than applying a national or state level incidence rate to a county-level, since rates vary geographically.

There are multiple limitations to this study. Ozone concentrations are based on monitor station data, which are not always located in every county in this study and thus may not reflect the ozone concentrations on finer spatial resolutions. Similar to prior studies, we use an interpolation method within BenMAP to estimate ozone concentrations based on surrounding air pollution monitors.¹⁷ Another limitation is the use of a single concentration-response function describing the relationship between ozone concentrations and asthma ED visits in all ages. Prior studies have reported differences based on sex and age in the concentration response relationship between ozone and asthma.¹⁸

Future work in this field would benefit from examining age-specific ozone-attributable ED visits for asthma exacerbation especially among vulnerable populations, such as children and elderly adults. Future research should also examine how mitigation strategies to address climate change affect ozone-attributable asthma ED visits.

CONCLUSIONS

Ozone air pollution is associated with thousands of asthma exacerbations requiring ED visit in New England. Rates of ozone-attributable asthma ED visits in New England varied by state and county in 2010, with the highest rates found in Connecticut and Massachusetts. These differences across space are likely due to a combination of varying demographic health characteristics and pollution levels, suggesting that efforts to reduce ozone-attributable ED visits should include improvements both to asthma treatments and air quality.

References

1. Adult Current Asthma Prevalence by State or Territory (2019) [Internet]. 2019 [cited June 8, 2021].
2. Zheng XY, Ding H, Jiang LN, et al. Association between Air Pollutants and Asthma Emergency Room Visits and Hospital Admissions in Time Series Studies: A Systematic Review and Meta-Analysis. *PloS one*. 2015;10(9):e0138146.
3. Anenberg SC, Henze DK, Tinney V, et al. Estimates of the Global Burden of Ambient PM_{2.5}, Ozone, and NO₂ on Asthma Incidence and Emergency Room Visits. *Environ Health Perspect*. 2018;126(10):107004.
4. Strosnider HM, Chang HH, Darrow LA, Liu Y, Vaidyanathan A, Strickland MJ. Age-Specific Associations of Ozone and Fine Particulate Matter with Respiratory Emergency Department Visits in the United States. *American journal of respiratory and critical care medicine*. 2019;199(7):882-90.
5. Izquierdo R, Garcia Dos Santos S, Borge R, et al. Health impact assessment by the implementation of Madrid City air-quality plan in 2020. *Environ Res*. 2020;183:109021.
6. U.S. EPA. Environmental Benefits Mapping and Analysis Program: Community Edition (BenMAP-CE) User Manual and Appendices. Research Triangle Park, North Carolina. April 2017.
7. HCUP Databases. Healthcare Cost and Utilization Project (HCUP), Agency for Healthcare Research and Quality. Rockville, MD. April 2021. www.hcup-us.ahrq.gov/databases.jsp.
8. Centers for Disease Control and Prevention. National Environmental Public Health Tracking Network [cited 2021 July 22]. Available from: www.cdc.gov/ephracking.
9. Sacks JD, Lloyd JM, Zhu Y, et al. The Environmental Benefits Mapping and Analysis Program - Community Edition (BenMAP-CE): A tool to estimate the health and economic benefits of reducing air pollution. *Environ Model Softw*. 2018;104:118-29.
10. Ji M, Cohan DS, Bell ML. Meta-analysis of the Association between Short-Term Exposure to Ambient Ozone and Respiratory Hospital Admissions. *Environ Res Lett*. 2011;6(2).
11. United States Environmental Protection Agency. Review of the Ozone National Ambient Air Quality Standards, 40 CFR Part 50. 2020;85(251).
12. Berman JD, Fann N, Hollingsworth JW, et al. Health benefits from large-scale ozone reduction in the United States. *Environ Health Perspect*. 2012;120(10):1404-10.
13. Nassikas N, Spangler K, Fann N, et al. Ozone-related asthma emergency department visits in the US in a warming climate. *Environ Res*. 2020;183:109206.
14. Karmalkar AV, Bradley RS. Consequences of Global Warming of 1.5 degrees C and 2 degrees C for Regional Temperature and Precipitation Changes in the Contiguous United States. *PloS one*. 2017;12(1):e0168697.
15. Anenberg SC, Haines S, Wang E, Nassikas N, Kinney PL. Synergistic health effects of air pollution, temperature, and pollen exposure: a systematic review of epidemiological evidence. *Environ Health*. 2020;19(1):130.

16. Fann NL, Nolte CG, Sarofim MC, Martinich J, Nassikas NJ. Associations Between Simulated Future Changes in Climate, Air Quality, and Human Health. *JAMA Netw Open*. 2021;4(1):e2032064.
17. Carvour ML, Hughes AE, Fann N, Haley RW. Estimating the Health and Economic Impacts of Changes in Local Air Quality. *American journal of public health*. 2018;108(S2):S151-S7.
18. Sheffield PE, Zhou J, Shmool JL, Clougherty JE. Ambient ozone exposure and children's acute asthma in New York City: a case-crossover analysis. *Environ Health*. 2015;14:25.

Acknowledgments

We would like to thank Dr. Deborah Pearlman, Ashley Fogarty, and Samara Viner-Brown at the Rhode Island Department of Health for their help with obtaining background asthma data.

This work was presented previously in poster format at the American Thoracic Society International Conference on May 21, 2018 in San Diego, California.

Authors

Nicholas J. Nassikas, MD, Division of Pulmonary, Critical Care, and Sleep Medicine, Brown University, Providence, RI; Division of Pulmonary, Critical Care, and Sleep Medicine, Beth Israel Deaconess Medical Center, Boston, MA.

Keith Spangler, PhD, Department of Environmental Health, Boston University, Boston, MA.

Gregory A. Wellenius, ScD, Department of Environmental Health, Boston University, Boston, MA.

Disclosures

Dr. Wellenius has previously served as a visiting scientist at Google, LLC (Mountain View, CA) and has received consulting income from the Health Effects Institute (Boston, MA).

Correspondence

Nicholas J. Nassikas, MD
330 Brookline Ave, Boston, MA 02215
nnassika@bidmc.harvard.edu

Increased Temperatures Are Associated with Increased Utilization of Emergency Medical Services in Rhode Island

KATELYN MORETTI, MD, MS; BENJAMIN GALLO MARIN, AB, MD'23; LUKE B. SOLIMAN, MTS, MD'23;
NICHOLAS ASSELIN, DO; ADAM R. ALUISIO, MD, MSC, DTM&H

ABSTRACT

BACKGROUND: Increasing temperatures negatively impact health and increases demands on healthcare systems. However, this has been poorly studied in Rhode Island (RI). Here we characterize the impact of heat on emergency medical services (EMS) utilization in RI.

METHODS: The Rhode Island National Emergency Services Information System V3 dataset was merged with data from the National Center for Environmental Information of the National Oceanic and Atmospheric Administration from the summers of 2018 and 2019. The outcome of daily mean EMS runs were compared against the exposure increasing daily temperatures, measured as daily maximum, minimum and daily average °F, using Poisson regressions. Patient characteristics were included across temperature models.

RESULTS: Increasing daily temperatures were associated with increasing EMS encounters. The adjusted incident rate ratio (IRR) for mean daily EMS encounters by increasing maximum daily temperature was 1.006 (95% CI 1.004–1.007, Table 3). This resulted in a projected 17.2% increase in EMS runs on days with a maximum temperature of 65°F compared to days with a maximum temperature of 95°F. The adjusted IRR for mean daily EMS encounters by the daily minimum temperature was 1.004 (1.003–1.006) and the adjusted IRR for the mean daily EMS encounters by the daily average temperature was 1.006 (1.005–1.008).

CONCLUSIONS: Increasing minimum, maximum, and average daily temperatures were associated with increasing EMS utilization across Rhode Island in the summers of 2018 and 2019. Further research into these trends may help with planning and resource allocation as summer temperatures continue to rise.

KEYWORDS: Heat, Emergency Medical Services, Rhode Island

INTRODUCTION

Heat detrimentally affects human health, leading to increased morbidity and mortality.¹ As carbon emissions

continue to rise, the effects of heat on health will continue to worsen.² Specifically, increasing ambient temperature has been linked to worsened health outcomes, including hospitalizations and deaths across a range of disease-specific causes, including pure heat-related illness to cardiovascular, respiratory and renal disease.^{3–6} Despite an international commitment to limiting global temperature rise to 2° Celsius as outlined in the Paris Climate Agreement, carbon emissions and temperatures continue to rise.² Over the last 20 years, heat-related deaths in patients 65 years or older have increased 53.7%. This resulted in 296,000 global heat deaths in 2018, 19,000 of which were in the United States.² Preparation and mitigation strategies for population health from climate change will require a clear understanding of the current climate change impacts on healthcare delivery.

Within the state of Rhode Island, only one study has explored the impact of heat on Emergency Department (ED) visits demonstrating increasing ED volume.⁷ Logically, we expect this strain to increase across all aspects of the care continuum. However, no study has evaluated the impact of heat on EMS utilization across the state. Understanding changing demands in the summer may allow for improved planning and resource management as summers become hotter and longer. In this study, the correlation between increasing daily summer temperatures and emergency medical service utilization are described.

METHODS

Inclusion Criteria

All patient encounters with a Rhode Island agency emergency medical service (EMS) occurring in the state of Rhode Island between the months of June–August 2018 and 2019 were included for analysis.

Database Formation

The Rhode Island National Emergency Medical Services Information System (NEMSIS) V3 dataset was merged with the National Centers for Environmental Information of the National Oceanic and Atmospheric Administration (NOAA).⁸ The NEMSIS V3 dataset is managed by the Center of Emergency Medical Services of the State of Rhode Island Department of Health and collects pertinent ambulance-run data for all agencies across the state of Rhode Island. It

includes patient level data, including demographics, primary impression as well as the date, time and city of the encounter.

Duplicates were removed as defined by patient encounters that matched on date and time, age, race, gender, and complete address. Given that multiple EMS agencies may respond initially to the same call, a large number of duplicates were anticipated and observed. In addition, encounters not occurring within the state of Rhode Island were excluded. The NOAA database provides public access to the historical weather data and information in the United States. Daily maximum, minimum and average daily temperatures were extracted from the Providence, Theodore Francis Green State Airport weather station for dates of interest: June–August 2018 and June–August 2019 and merged by date with the NEMSIS dataset to provide temperature data on the date of each EMS call. Given possible effects of winter weather on EMS utilization, analysis was limited to months June–August. IRB approval was obtained through the RI Department of Health for access to the NEMSIS database.

Variable classifications

Patient-level data is ascertained from forms collected on each EMS run with each EMS agency responding completing a form. Demographic data extracted included: age, gender, race. Clinical data included: patient acuity and initial impression by EMS providers. Location information included: date and time of the response and address. The NOAA dataset contains no patient information and are available publicly. Age was categorized as <18 years, 18–65 and >65 years and race dichotomized as white versus non-white, consistent with previous heat research in Rhode Island.⁷ Acuity was classified by EMS providers within the NEMSIS database as green, yellow, red or black (deceased) consistent with the Simple Triage and Rapid Treatment (START) triage method.⁸ EMS primary impression were classified by physiologic system or topic including cardiac, renal, gastrointestinal disease, trauma.

Statistical Analysis

Given previous research demonstrating an increased mortality rate with maximum temperatures above 85°F in the state of Rhode Island,⁷ cohort characteristics including year (2018 vs 2019), age, gender, race and acuity were compared with chi squared analysis dichotomized by encounters occurring on days with a maximum temperature below 85°F versus those occurring on days with a maximum temperature equal to or above 85°F. Mean EMS runs dichotomized by daily temperature at 85°F were compared with a two-sample t test.


The outcome of daily mean EMS runs was modeled against the exposure of increasing daily temperatures using Poisson regression. Variables for the model were chosen using forward selection and a $P=0.2$. Variables considered included: the city of the response, year, patient age, gender, acuity

and EMS' first impression. Patient acuity, first impression and year were included in the maximum daily temperature model. Patient acuity, first impression, cohort year and age were included in the minimum daily temperature model and year and first impression were included in the average daily temperature model.

RESULTS

Within the NEMSIS database, 53,028 encounters were found for June–August 2018 and June–August 2019. After removal of duplicates and encounters not occurring within the state of Rhode Island, a final cohort of 49,548 EMS encounters remained (**Figure 1**). The majority (3,328) also matched on primary impression. Nine encounters had differing primary impressions (**Supplemental Table 1**).

Figure 1. Dataset creation with removal of duplicates from the EMS dataset. *For a list of the differing primary impressions see Supplemental table 1.



Supplemental table 1. Nine patients matched on date and time, age, race, gender and primary address with differing primary impressions. The encounter with impression 2 was removed from analysis.

Patient	Impression 1	Impression 2 (removed)
1	Traumatic injury	Injury of head
2	Injury of lower leg	Traumatic Injury, unspecified
3	Pain, unspecified	Injury of food
4	Fever	Missing
5	Injury of shoulder and upper arm	Injury of head
6	Injury of foot	Pain, unspecified
7	Injury of wrist, hand and fingers	Injury of lower back
8	Hypertension	Headache
9	Injury of neck	Injury of lower back.

Table 1. EMS response and patient characteristics for days below and above 85°F in Rhode Island, June–August 2018 and 2019.

	Maximum daily temperature < 85 °F N=28,168 (%)	Maximum daily temperature ≥ to 85 °F N=21,378 (%)	P value
Mean daily EMS runs	284.7 (95% CI 280.3–289.0)	300.7 (95% CI 295.9–305.5)	<0.0001
Year			
2018	12,782 (52.0)	11,788 (48.0)	<0.0001
2019	15,372 (61.6)	9,579 (38.4)	
Age			
<18	1,596 (5.7)	1,189 (5.6)	0.58
18-65	14,892 (52.9)	11,401 (53.4)	
>65	11,650 (41.4)	8,767 (41.1)	
Gender			
Male	13,253 (47.1)	10,188 (47.7)	0.33
Female	14,900 (52.9)	11,177 (52.3)	
Missing	10 (0.04)	10 (0.05)	
Race			
White	7,072 (25.1)	5,403 (25.3)	0.22
Non-white	1,013 (3.6)	774 (3.6)	
Missing	20,083 (71.3)	15,201 (71.1)	
Acuity			
Green	14,789 (52.5)	11,309 (52.9)	0.30
Yellow	4,753 (16.9)	3,643 (17.0)	
Red	964 (3.4)	680 (3.2)	
Black	1 (0.0)	3 (0.01)	
Missing	7,661 (27.2)	5,743 (26.9)	
Top Five Primary Impressions			
Other	6,079	4,788	0.30
Cardiac	3,719	2,814	
Trauma	3,587	2,682	
Psychiatric	3,518	2,602	
Gastrointestinal	2,478	1,879	

Mean daily EMS runs were statistically higher on days with a maximum temperature above 85°F (285 vs. 300, p=<0.0001 **Table 1**). There was no statistical difference in patient characteristics: age, gender, race or acuity between those with an EMS encounter on days less than or greater than 85°F (**Table 1**).

The mean maximum, mean minimum and mean average daily temperatures between 2018 and 2019 were similar (**Table 2**). However, there were more days > 85°F in 2018 than 2019 (46% vs 37%) and a greater percentage of EMS runs on days that were over 85°F in 2018 versus 2019.

Increasing daily temperatures were associated with increasing EMS encounters for the state of Rhode Island in the summers of 2018 and 2019 (**Figure 2**). The adjusted incident rate ratio (IRR) for mean daily EMS encounters by increasing maximum daily temperature

Table 2. Daily temperature characteristics for the overall cohort and by year for June–August.

	Mean maximum daily temperature in °F (95% CI)	Mean average daily temperature in °F (95% CI)	Mean minimum daily temperature in °F (95% CI)
Overall	82.2 (81.2–83.3)	72.6 (71.7–73.5)	63.9 (62.9–64.9)
2018	82.9 (82.8–83.0)	73.1 (71.7–74.4)	64.4 (62.9–65.8)
2019	81.9 (80.4–83.3)	72.2 (71.0–73.4)	63.5 (62.3–64.8)

Figure 2. Mean daily EMS Responses by the maximum, minimum and average daily temperatures (°F) across the state of Rhode Island for June–August 2018 and 2019




Table 3. Maximum daily temperature model controlled for patient acuity, first impression and year. Minimum daily temperature model controlled for patient acuity, first impression, year and age. Average daily temperature model controlled for year and first impression.

	Unadjusted Incident Rate Ratio Mean Daily EMS Encounters (95% CI)	Adjusted Incident Rate Ratio Mean Daily EMS Encounters (95% CI)
Maximum Daily Temperature	1.0053 (1.0052–1.0054)	1.006 (1.004–1.007)
Minimum Daily Temperature	1.0037 (1.0036–1.0038)	1.004 (1.003–1.006)
Average Daily Temperature	1.0053 (1.0052–1.0053)	1.006 (1.005–1.008)

was 1.006 (95% CI 1.004–1.007, **Table 3**). Thus, according to this model, we would expect an average of 250 EMS encounters on summer days with a maximum temperature of 65°F. While, on days with a maximum temperature of 95°F, we would expect an average call volume of 293, a 17.2% increase.

The adjusted IRR for mean daily EMS encounters by the daily minimum temperature was 1.004 (1.003–1.006) and the adjusted IRR for the mean daily EMS encounters by the daily average temperature was 1.006 (1.005–1.008).

CONCLUSIONS

Increasing minimum, maximum and average daily temperatures were associated with increasing EMS utilization across the state of Rhode Island in the summers of 2018 and 2019. Past literature has shown an association between increased ambient temperature and patient morbidity and mortality resulting in greater utilization and strain of the healthcare system.^{5,9,10} In Rhode Island (RI), rising summer daily temperatures have been linked to worsened emergency department strain, with close to a 24% increase in Emergency Department (ED) visits.⁷

Heat has been associated with increased acute diseases as well as exacerbations of chronic illnesses. Specifically, heat has been associated with increased incidence of cardiac arrest,⁶ subarachnoid hemorrhage,¹¹ acute kidney injury,¹² hyponatremia,¹³ pulmonary arterial hypertension¹⁴ and ischemic heart disease¹⁵ to name a few. However, in this study, EMS' primary impression of the patient's illness did not change for days less than or greater than 85°F. In addition, the patients demographics including age, race acuity level did not change.

There was a greater percentage of patients seen on days over 85°F in 2018 than in 2019 most likely because there were more days over 85°F in 2018 as compared to 2019.

LIMITATIONS

Associations of EMS runs, heat and patient acuity or race were limited by large amounts of missing patient data.

However, it is most likely missing at random and therefore represents a non-differential error. Thus, its impact on the association observed between heat and increasing EMS utilization is presumed to be minor. Given that heat impacts vulnerable populations disproportionately (SOURCE), vulnerable races may demonstrate a disproportionate increase in EMS utilization as temperature increases. However, data were insufficient to examine this relationship and future studies are needed to further examine this relationship. In addition, final hospital diagnosis was not available for patient encounters. The proxy of EMS' primary impression was used but may not fully capture the impacts on health outcomes in the population. Further research linking EMS records to hospital records including an evaluation of the final diagnosis would be beneficial.

Heat puts strains across all strata of emergency services, including EMS. As temperatures continue to rise, understanding the extent of increased demands may allow for appropriate staffing, early warning systems and increased public mitigation strategies such as cooling centers.

Additional heat considerations or measurements may also be important to examine. For example urban heat islands (UHIs) are areas of urbanization that, secondary to heat absorptive surfaces such as concrete and asphalt, result in surface temperatures above the average surface temperature in that geographic area.¹⁶ Urban heat islands and heatwaves, neither examined in this study, have also been identified as exacerbating heat effects and may play an important role in the increased EMS usage during hot days seen here. Heatwave events, or prolonged periods of abnormally high temperatures, have resulted in specific negative health impacts including an increase in heat-related mortality rates⁹ As our climate warms, heatwaves are increasing in frequency and associated morbidity is worsening.² In addition, UHIs are areas of urbanization that, secondary to heat absorptive surfaces such as concrete and asphalt, result in surface temperatures above the average surface temperature in that geographic area.¹⁷ Patients living in UHIs are especially susceptible to heat waves and the deleterious health effects of increasing ambient temperatures.¹⁶ Future research will further characterize EMS utilization during heatwaves and within UHIs. This understanding will better inform future preparedness and mitigation strategies.

References

1. Campbell S, Remenyi TA, White CJ, Johnston FH. Heatwave and health impact research: A global review. *Health Place* 2018;53:210–8.
2. Watts N, Amann M, Arnell N, et al. The 2020 report of The Lancet Countdown on health and climate change: responding to converging crises. *Lancet Lond Engl* 2021;397(10269):129–70.
3. Vaidyanathan A, Saha S, Vicedo-Cabrera AM, et al. Assessment of extreme heat and hospitalizations to inform early warning systems. *Proc Natl Acad Sci* 2019;116(12):5420–7.
4. Huang G, Zhou W, Cadenasso ML. Is everyone hot in the city?

- Spatial pattern of land surface temperatures, land cover and neighborhood socioeconomic characteristics in Baltimore, MD. *J Environ Manage* 2011;92(7):1753–9.
5. Riley K, Wilhalme H, Delp L, Eisenman DP. Mortality and Morbidity during Extreme Heat Events and Prevalence of Outdoor Work: An Analysis of Community-Level Data from Los Angeles County, California. *Int J Environ Res Public Health* 2018;15(4).
 6. Kang S-H, Oh I-Y, Heo J, et al. Heat, heat waves, and out-of-hospital cardiac arrest. *Int J Cardiol* 2016;221:232–7.
 7. Kingsley SL, Eliot MN, Gold J, Vanderslice RR, Wellenius GA. Current and Projected Heat-Related Morbidity and Mortality in Rhode Island. *Environ Health Perspect* 2016;124(4):460–7.
 8. Riza'i A, Ade WRA, Albar I, Sulitio S, Muharris R. Teaching Start Triage: A Comparison of Lecture and Simulation Methods. *Adv Sci Lett* 2018;24(9):6890–2.
 9. Knowlton K, Rotkin-Ellman M, King G, et al. The 2006 California heat wave: impacts on hospitalizations and emergency department visits. *Environ Health Perspect* 2009;117(1):61–7.
 10. Ragetti MS, Vicedo-Cabrera AM, Flückiger B, Röösli M. Impact of the warm summer 2015 on emergency hospital admissions in Switzerland. *Environ Health Glob Access Sci Source* 2019;18(1):66.
 11. Lee S, Guth M. Associations between Temperature and Hospital Admissions for Subarachnoid Hemorrhage in Korea. *Int J Environ Res Public Health* 2017;14(4).
 12. Thongprayoon C, Qureshi F, Petnak T, et al. Impact of Acute Kidney Injury on Outcomes of Hospitalizations for Heat Stroke in the United States. *Dis Basel Switz* 2020;8(3).
 13. Sailer CO, Winzeler B, Nigro N, Bernasconi L, Mueller B, Christ-Crain M. Influence of Outdoor Temperature and Relative Humidity on Incidence and Etiology of Hyponatremia. *J Clin Endocrinol Metab* 2019;104(4):1304–12.
 14. Jehn M, Gebhardt A, Liebers U, et al. Heat stress is associated with reduced health status in pulmonary arterial hypertension: a prospective study cohort. *Lung* 2014;192(4):619–24.
 15. Huang J, Zeng Q, Pan X, Guo X, Li G. Projections of the effects of global warming on the disease burden of ischemic heart disease in the elderly in Tianjin, China. *BMC Public Health* 2019;19(1):1465.
 16. Heaviside C, Macintyre H, Vardoulakis S. The Urban Heat Island: Implications for Health in a Changing Environment. *Curr Environ Health Rep* 2017;4(3):296–305.
 17. Oke TR. The energetic basis of the urban heat island. *Q J R Meteorol Soc* 1982;108(455):1–24.

Acknowledgments

Funding was provided by Brown Emergency Physicians, Inc.

Authors

- Katelyn Moretti, MD, MS, Department of Emergency Medicine, Warren Alpert Medical School of Brown University, Providence, RI.
- Benjamin Gallo Marin, AB, MD'23, Department of Emergency Medicine, Warren Alpert Medical School of Brown University, Providence, RI.
- Luke B. Soliman, MTS, MD'23, Department of Emergency Medicine, Warren Alpert Medical School of Brown University, Providence, RI.
- Nicholas Asselin, DO, Department of Emergency Medicine, Warren Alpert Medical School of Brown University, Providence, RI.
- Adam R. Aluisio, MD, MSc, DTM&H, Department of Emergency Medicine, Warren Alpert Medical School of Brown University, Providence, RI.

Correspondence

Katelyn Moretti, MD
55 Claverick St. Providence, RI 02903
katelyn_moretti@brown.edu

Potential Effects of Climate Change on Tick-borne Diseases in Rhode Island

HOWARD S. GINSBERG, PhD; JANNELLE COURET, PhD, MEM; JASON GARRETT, BSN, MPH; THOMAS N. MATHER, PhD; ROGER A. LEBRUN, PhD

ABSTRACT

Human cases of tick-borne diseases have been increasing in the United States. In particular, the incidence of Lyme disease, the major vector-borne disease in Rhode Island, has risen, along with cases of babesiosis and anaplasmosis, all vectored by the blacklegged tick. These increases might relate, in part, to climate change, although other environmental changes in the northeastern U.S. (land use as it relates to habitat; vertebrate host populations for tick reproduction and enzootic cycling) also contribute. Lone star ticks, formerly southern in distribution, have been spreading northward, including expanded distributions in Rhode Island. Illnesses associated with this species include ehrlichiosis and alpha-gal syndrome, which are expected to increase. Ranges of other tick species have also been expanding in southern New England, including the Gulf Coast tick and the introduced Asian longhorned tick. These ticks can carry human pathogens, but the implications for human disease in Rhode Island are unclear.

KEYWORDS: ticks, climate change, Lyme disease, Babesiosis, Ehrlichiosis

INTRODUCTION

Ticks are major disease vectors in north temperate regions worldwide, transmitting a wide variety of pathogen types. Although mosquitoes are recognized as vectors of widespread tropical diseases such as malaria and dengue, they are responsible for far fewer cases of disease than ticks in North America. In Rhode Island, mosquitoes are responsible for an average of only about one human case of arboviral disease per year.¹ In contrast, annual cases of tickborne ailments in Rhode Island run into the thousands (<https://health.ri.gov/data/diseases/>). Climate change would be expected to result, overall, in a poleward shift in the distributions of tick-borne diseases.² However, numerous other environmental and socioeconomic factors that affect vector-borne disease transmission are also changing. This complicates direct predictions of climate change effects on future disease distribution patterns, and only a few examples have been demonstrated convincingly for tick-borne pathogens.² In this brief review

we identify the major tick-borne diseases that currently occur in Rhode Island, describe the underlying transmission dynamics, and consider the likely effects of climate change. We also discuss tick species that have recently spread into the region and the pathogens they might bring with them.

MAJOR TICK-BORNE DISEASES IN RHODE ISLAND

The major tick disease vector in Rhode Island is the blacklegged tick (sometimes called the deer tick), *Ixodes scapularis*, which transmits the spirochete that causes Lyme disease. Several tick-borne pathogens cause human disease in Rhode Island, with Lyme disease being far the most common (Table 1). Reported cases of Lyme disease in the state are in the vicinity of 1,000 per year, but this is undoubtedly a substantial underestimate. A recent CDC estimate using insurance claims data indicates that the annual number of cases diagnosed in the US from 2010–2018 averaged about 476,000, compared to the average of only about 35,000 cases per year reported to the agency.³

The blacklegged tick is concentrated in woodlands with leaf litter and is most abundant in the southern portion of Rhode Island, but numbers fluctuate substantially from year

Table 1. Major tick-borne diseases in Rhode Island

Tick vector	Disease	Pathogen	Mean # cases per yr. (2015–2019)*
Blacklegged tick, deer tick (<i>Ixodes scapularis</i>)	Lyme disease	<i>Borrelia burgdorferi</i>	1,004.2
	Anaplasmosis	<i>Anaplasma phagocytophilum</i>	141.4
	Babesiosis	<i>Babesia microti</i>	165.8
	Powassan Encephalitis	Powassan virus, Deer tick virus	0.8
Lone star tick (<i>Amblyomma americanum</i>)	Ehrlichiosis	<i>Ehrlichia chaffeensis</i> , <i>E. ewingi</i>	57.0
American dog tick (<i>Dermacentor variabilis</i>)	Spotted Fever Group Rickettsiosis	<i>Rickettsia rickettsii</i> , <i>R. spp.</i>	6.0

* Data in Table 1 and Figure 1 from CDC (<https://www.cdc.gov/nndss/data-statistics/infectious-tables/index.html>)

to year such that dense populations extend throughout the state in some years.⁴ In addition to Lyme disease, which is caused by a spirochete, the blacklegged tick is also the primary vector of the two other major tick-borne pathogens in Rhode Island, the rickettsia that causes anaplasmosis and the protozoan that causes babesiosis (Table 1). This tick can also transmit pathogens less commonly reported in Rhode Island, including Powassan encephalitis virus, and the relapsing fever spirochete *Borrelia miyamotoi*.⁵

Two other tick species are important vectors of human pathogens in Rhode Island, the lone star tick, *Amblyomma americanum*, which transmits the rickettsiae that cause ehrlichiosis and possibly those that cause Rocky Mountain Spotted Fever, and the American dog tick, *Dermacentor variabilis*, the primary vector of Rocky Mountain Spotted Fever (Table 1). These two tick species can also transmit the bacterium *Francisella tularensis*, which causes tularemia, but this pathogen is often transmitted via infectious fluids through handling or even breathing aerosols from infected animals rather than from tick bite. These diseases are far less common than some of those vectored by *I. scapularis*, but the distributions of these tick species are changing,⁶ and effects of climate change are potentially important. Resources for identification of tick species, and information sources relevant to disease transmission and status in Rhode Island are readily available online (<https://web.uri.edu/tickencounter/>).

CLIMATE CHANGE AND TRANSMISSION PATTERNS OF TICK-BORNE PATHOGENS

Lyme disease, Anaplasmosis, Babesiosis

Rhode Island is near the center of the current geographic distribution of Lyme disease, and is likely to remain an epicenter of Lyme and other major tick-borne diseases in the foreseeable future. In the eastern and central U.S., Lyme disease is most common in the northern states, and relatively uncommon in the south (<https://www.cdc.gov/lyme/data-surveillance/maps-recent.html>). This pattern results primarily from latitudinal trends in host-seeking behavior and host associations of larval and nymphal ticks. Northern *I. scapularis* wait on leaves and twigs at the leaf litter surface and attach to passing hosts, while southern ticks remain below the litter surface when host-seeking.⁷ Therefore, people frequently encounter northern ticks in their woodland habitats, but there is relatively little human exposure in the south.⁸ Furthermore, northern larval and nymphal ticks feed primarily on small mammals such as rodents and shrews, which are excellent reservoirs for the Lyme spirochete, while southern ticks feed mostly on lizards, which are relatively poor reservoirs, so prevalence of spirochetal infection is much lower in southern than in northern ticks.⁹ Some of these latitudinal differences might well be related to north-south climatic differences,¹⁰ and so might be affected by climate change. Some modeling studies predict that climate

change-related effects on factors such as tick development and phenology are likely to result in increases in Lyme disease incidence in the northeast,^{11,12} as well as increased spread northward into Canada. Indeed, cases of Lyme disease have been increasing in the U.S., and northward spread into Canada has been documented.¹³ However, predictions for changes in current endemic areas vary with model assumptions and local complexities,¹⁴ so these effects remain uncertain for Rhode Island. Human cases of Lyme disease reported to the CDC have been increasing in Rhode Island (Figure 1A). Furthermore, evidence from various sources suggest that populations of *I. scapularis*, and human cases of associated diseases, have been increasing in the northeast,^{6,15} and that these changes result, at least in part, from the effects of climate change.²

Epidemiological trends in cases of anaplasmosis and babesiosis are expected to be similar to those for Lyme disease, because they have the same tick vector species, *I. scapularis*, and the same primary reservoir species, the white-footed mouse (*Peromyscus leucopus*). As such, patterns of epizootiology are similar for these pathogens, except that transmission efficiency is apparently greater for *Borrelia burgdorferi* than for *Anaplasma phagocytophila*¹⁶ or *Babesia microti*,¹⁷ so infection prevalence of these pathogens in ticks is lower than that of Lyme spirochetes, and there are fewer human cases (Figure 1B). Thus, anaplasmosis and babesiosis are likely to remain important in Rhode Island,

Figure 1A.





Figure 1B.



and though cases are expected to increase, they will likely remain less prevalent than Lyme disease.

Ehrlichiosis, alpha-gal syndrome

The lone star tick, *A. americanum*, is the vector of the rickettsiae that cause ehrlichiosis.¹⁸ Growing evidence indicates that lone star tick bites are also responsible for alpha-gal syndrome (also called red meat allergy or tick bite meat allergy), an allergic response to the carbohydrate galactose-alpha-1,3-galactose, which is found in mammalian meat.¹⁹ Populations of this tick species are spreading northward²⁰ and ehrlichiosis cases have been increasing in Rhode Island (<https://health.ri.gov/data/diseases/Ehrlichiosis.pdf>). Lone star ticks were formerly southern in distribution, with dense populations from central New Jersey and southward, but populations increased substantially on Long Island, NY, in the late 1980s and 1990s, and they are now established in Connecticut.²¹ Climate change might contribute to the northward spread of this species, but increasing populations of its primary hosts, including white-tailed deer, coyotes, and wild turkeys, are also important.²⁰ This tick has previously been rare on the mainland of Rhode Island, although a dense population has existed on Prudence Island in Narragansett Bay, at least since 1990.²² Populations have increased substantially since 2016 on Conanicut Island, and established populations (all three life stages) are now being detected at mainland sites, especially in coastal communities on both sides of Narragansett Bay and the south coast (TNM, personal observation). Thus, lone star ticks are spreading northward, likely related to changes in climate and in host distributions. Rhode Island is within the potential range of this species predicted by some models, but not others.²³ Given the recent local increases in *A. americanum* populations and the persistent population on Prudence Island, it is reasonable to expect increases in Rhode Island cases of ehrlichiosis and alpha-gal syndrome in the future.

Spotted fever group rickettsioses

The primary vector of *Rickettsia rickettsii*, etiologic agent of Rocky Mountain Spotted Fever (RMSF) in North America, is the American dog tick, *D. variabilis*, which is responsible for modest numbers of human cases in Rhode Island (Table 1). This species is distributed widely in the eastern U.S., and climate change might well result in larger populations in the northern U.S. and Canada.⁶ Most RMSF cases currently occur in a band from the Carolinas to Oklahoma,²⁴ so climate change could result in greater numbers of cases in our area. This pathogen is maintained vertically in *D. variabilis* (passed from mother to offspring), and outbreaks tend to be focal, with local amplification presumably resulting from horizontal transmission by ticks among mammal hosts.²⁵ However, these dynamics are poorly understood, so it is difficult to predict the likely effects of a changing climate on the epidemiology of RMSF.

Another spotted fever group rickettsia, *R. parkeri*, is transmitted by the Gulf Coast tick, *Amblyomma maculatum*. This tick was formerly found only in the southern states, but it has recently expanded its range into the mid-Atlantic region,²⁶ and an established population has recently been reported in Connecticut.²⁷ Human infection with *R. parkeri* has not yet been detected in Rhode Island, but given the northward range expansion of the vector, this possibility is worthy of attention in the future.

TICK SPECIES LIKELY TO SPREAD INTO RHODE ISLAND

Northward spread of two tick species has already been discussed: *A. americanum*, which is already present in Rhode Island and is likely to increase in the future, and *A. maculatum*, which has not yet been detected in the state, but which is spreading into the region. Both species present potential implications for human health, and their range expansion might be related, at least in part, to climate change.

A third tick species that was introduced into the United States, presumably from its native range in Asia, has recently appeared in the region, and the implications for human health are currently unclear. The Asian longhorned tick, *Haemaphysalis longicornis*, was first detected in North America in 2017. Examinations of earlier collections indicate that this species was present in the United States several years before that detection. It has now been detected in 16 states, and on dozens of animal species (https://www.aphis.usda.gov/animal_health/animal_diseases/tick/downloads/longhorned-tick-sitre.pdf). Populations are known to exist on Block Island, and *H. longicornis* has recently been collected on the Rhode Island mainland (TNM, personal observation). The North American populations are parthenogenetic and can rapidly increase to high densities. This tick is known to vector numerous pathogens in Asia, including Severe Fever with Thrombocytopenia Syndrome (SFTS) virus,²⁸ and has become an introduced pest of cattle in Australia. It serves as a vector of the cattle pathogen *Theileria orientalis* Ikeda in Virginia.²⁹ To date, no human pathogens have been isolated from *H. longicornis* in the United States, except for one *B. burgdorferi* isolate from Pennsylvania.³⁰ Lab studies suggest that *H. longicornis* is not a competent vector of *B. burgdorferi*³¹ or of *A. phagocytophilum*,³² but that this tick species can acquire and transmit the RMSF pathogen *R. rickettsii*.³³ The potential role of this species as a vector of human pathogens in the U.S. remains unclear. Fortunately, this species does not readily bite humans, although cases of attachment to humans have been reported in the United States.³⁴ Rhode Island is within the predicted distribution of this species,³⁵ and populations are likely to increase in the future.

CONCLUSIONS

Incidence levels of tick-borne diseases generally respond to changes in environmental conditions such as habitat, vertebrate host populations, and climate, as well as to factors that influence tick encounter by people. Human cases of several tick-associated diseases in Rhode Island, Lyme disease chief among them, but also anaplasmosis, babesiosis, ehrlichiosis, and alpha-gal syndrome, are likely to increase in the future. Northward range expansions of some tick species into Rhode Island are likely, but the implications for human health are unclear.

References

- Ginsberg HS, Gettman A, Becker E, Bandyopadhyay A, LeBrun RA. Environmental management of mosquito-borne viruses in Rhode Island. *RI Med J*. 2013;96(7):37-41.
- Ogden NH, Beard CB, Ginsberg HS, Tsao JI. Possible effects of climate change on ixodid ticks and the pathogens they transmit: what has been predicted and what has been observed. *J Med Entomol*. 2021;58(4):1536-1545.
- Kugeler CJ, Schwartz AM, Delorey MJ, Meade PS, Hinckley AF. Estimating the frequency of Lyme disease diagnoses, United States, 2010-2018. *Emerg Inf Dis*. 2021;27(2):616-619.
- Berger KA, Ginsberg HS, Dugas KD, Hamel LH, Mather TN. Adverse moisture events predict seasonal abundance of Lyme disease vector ticks (*Ixodes scapularis*). *Parasites & Vectors*. 2014;7:181.
- Fiorito TM, Reece R, Flanigan TP, Silverblatt FJ. *Borrelia miyamotoi* polymerase chain reaction positivity on a tick-borne disease panel in an endemic region of Rhode Island. *Inf Dis Clin Practice*. 2017;25(5):250-254.
- Sonenshine DE. Range expansion of tick disease vectors in North America: Implications for spread of tick-borne disease. *Int J Env Res Pub Hlth*. 2018;15:478.
- Arsnoe I, Tsao JI, Hickling GJ. Nymphal *Ixodes scapularis* questing behavior explains geographic variation in Lyme borreliosis risk in the eastern United States. *Ticks Tick-borne Dis*. 2019;10:553-563.
- Stromdahl EH, Hickling GJ. Beyond Lyme: aetiology of tick-borne human diseases with emphasis on the southeastern United States. *Zoon Pub Hlth*. 2012;59:48-64.
- Ginsberg HS, Hickling GJ, Burke RL, Ogden NH, Beati L, LeBrun RA, Arsnoe IM, Gerhold R, et al. Why Lyme disease is common in the northern U.S., but rare in the south: the roles of host choice, host-seeking behavior, and tick density. *PLoS Biol*. 2021;19(1):e3001066.
- Ginsberg HS, Albert M, Acevedo L, Dyer MC, Arsnoe IM, Tsao JI, Mather TN, LeBrun RA. Environmental factors affecting survival of immature *Ixodes scapularis* and implications for geographical distribution of Lyme disease: the climate/behavior hypothesis. *PLoS ONE* 2017;12(1):e0168723.
- Levi T, Keesing F, Oggenfuss K, Ostfeld RS. Accelerated phenology of blacklegged ticks under climate warming. *Phil Trans R Soc B*. 2015;370:20130556.
- Couper LI, MacDonald AJ, Mordecai EA. Impact of prior and projected climate change on US Lyme disease incidence. *Glob Change Biol*. 2021;27:738-754.
- Kulkarni MA, Narula I, Slatculescu AM, Russell C. Lyme disease emergence after invasion of the blacklegged tick, *Ixodes scapularis*, Ontario, Canada, 2010-2016. *Emerg Inf Dis*. 2019;25(2):328-332.
- Ostfeld RS, Brunner JL. Climate change and *Ixodes* tick-borne diseases of humans. *Phil Trans R Soc B*. 2015;370:20140051.
- Menis M, Whitaker BI, Wernecke M, Jiao Y, Eder A, Kumar S, Xu W, Wei Y, et al. Babesiosis occurrence among United States Medicare beneficiaries, ages 65 and older, during 20006-2017: overall and by state and county of residence. *Open Forum Inf Dis*. 2021;8(2).ofaa608. <https://doi.org/10.1093/ofid/ofaa608>.
- Levin ML, Ross DE. Acquisition of different isolates of *Anaplasma phagocytophilum* by *Ixodes scapularis* from a model animal. *Vector-borne Zoon Dis*. 2004;4:53-59.
- Mather TN, Telford SR, Moore SI, Spielman A. *Borrelia burgdorferi* and *Babesia microti*: efficiency of transmission from reservoirs to vector ticks (*Ixodes dammini*). *Exp Parasitol*. 1990;70(1):55-61.
- Mixson TR, Campbell SR, Gill JS, Ginsberg HS, Reichard MV, Schulze TL, Dasch GA. Prevalence of *Ehrlichia*, *Borrelia*, and rickettsial agents in *Amblyomma americanum* (Acari: Ixodidae) collected from nine states. *J Med Entomol*. 2006;43(6):1261-1268.
- Young I, Prematunge C, Pussegoda K, Corrin T, Waddell L. Tick exposures and alpha-gal syndrome: a systematic review of the evidence. *Ticks Tick-borne Dis*. 2021;12:101674
- Molaei G, Little EAH, Williams SC, Stafford KC. Bracing for the worst – range expansion of the lone star tick in the northeastern United States. *N Engl J Med*. 2019;381(23):2189-2192.
- Stafford KC, Molaei G, Little EAH, Paddock CD, Karpathy SE, Labonte AM. Distribution and establishment of the lone star tick in Connecticut and implications for range expansion and public health. *J Med Entomol*. 2018;55(6):1561-1568.
- Mather TN, Mather ME. Intrinsic competence of three ixodid ticks (Acari) as vectors of the Lyme disease spirochete. *J Med Entomol*. 1990;27(4):646-650.
- Springer YP, Jarnevich CS, Barnett DT, Monaghan AJ, Eisen RJ. Modeling the present and future geographic distribution of the lone star tick, *Amblyomma americanum* (Ixodida: Ixodidae) in the continental United States. *Am J Trop Med Hyg*. 2015;93(4):875-890.
- Biggs HM, Behravesh CB, Bradley KK, Dahlgren FS, Drexler MA, Dumler JS, Folk SM, Kato CY, et al. Diagnosis and management of tickborne rickettsial diseases: Rocky Mountain Spotted Fever and other spotted fever group rickettsioses, ehrlichiosis, and anaplasmosis – United States. *Morb Mort Wkly Rept*. 2016;65(2):1-45.
- Macaluso KR, Paddock CD. Tick-borne spotted fever group rickettsioses and *Rickettsia* species. Pp. 211-250. In Sonenshine DE, Roe RM (eds.), *Biology of Ticks*, vol. 2, 2nd edition. 2014. Oxford University Press, NY.
- Nadolny RM, Gaff HD. Natural history of *Amblyomma maculatum* in Virginia. *Ticks Tick-borne Dis*. 2018;9:188-195.
- Molaei G, Little EAH, Khalil N, Ayres BN, Nicholson WL, Paddock CD. Established population of the Gulf coast tick, *Amblyomma maculatum* (Acari: Ixodidae), infected with *Rickettsia parkeri* (Rickettsiales: Rickettsiaceae), in Connecticut. *J Med Entomol*. 2021;58(3):1459-1462.
- Zhao L, Li J, Cui X, Jia N, Wei J, Xia L, Wang H, Zhou Y, et al. Distribution of *Haemaphysalis longicornis* and associated pathogens: analysis of pooled data from a Chinese field survey and global published data. *Lancet Planet Hlth*. 2020;4:e320-29.
- Dinkel KD, Herndon DR, Noh SM, Lahmers KK, Todd SM, Ueti MW, Scoles GA, Mason KL, et al. A U.S. isolate of *Theileria orientalis*, Ikeda genotype, is transmitted to cattle by the invasive Asian longhorned tick, *Haemaphysalis longicornis*. *Parasites & Vectors*. 2021;14:157.
- Price KJ, Graham CB, Witmier BJ, Chapman HA, Coder BL, Boyer CN, Foster E, Maes SE, et al. *Borrelia burgdorferi* sensu stricto DNA in field-collected *Haemaphysalis longicornis* ticks, Pennsylvania, United States. *Emerg Inf Dis*. 2021;27(2):608-611.
- Breuner NE, Ford SL, Hojgaard A, Osikowicz LM, Parise CM, Rizzo MFR, Bai Y, Levin ML, et al. Failure of the Asian longhorned tick, *Haemaphysalis longicornis*, to serve as an experi-

- mental vector of the Lyme disease spirochete, *Borrelia burgdorferi* sensu stricto. *Ticks Tick-borne Dis.* 2020;11:101311.
32. Levin ML, Stanley HM, Hartzler K, Snellgrove AN. Incompetence of the Asian longhorned tick (Acari: Ixodidae) in transmitting the agent of human granulocytic anaplasmosis in the United States. *J Med Entomol.* 2021;58(3):1419-1423.
 33. Stanley HM, Ford SL, Snellgrove AN, Hartzler K, Smith EB, Krapivnaya I, Levin ML. The ability of the invasive Asian longhorned tick *Haemaphysalis longicornis* (Acari: Ixodidae) to acquire and transmit *Rickettsia rickettsii* (Rickettsiales: Rickettsiaceae), the agent of Rocky Mountain Spotted Fever, under laboratory conditions. *J Med Entomol.* 2020;57(5):1635-1639.
 34. Wormser GP, McKenna D, Piedmonte N, Vinci V, Egizi AM, Backenson B, Falco RC. First recognized human bite in the United States by the Asian Longhorned tick, *Haemaphysalis longicornis*. *Clin Inf Dis.* 2020;70:314-316.
 35. Rochlin I. Modeling the Asian longhorned tick (Acari: Ixodidae) suitable habitat in North America. *J Med Entomol.* 2019;56(2):384-391.

Acknowledgments

The authors thank Michael Gosciminski, MT, MPH, and Utpala Bandy, MD, MPH, for constructive comments on an early draft of the manuscript. This work was supported by the US Geological Survey, RI Department of Health, and the University of Rhode Island. Thank you to all the RIDOH Center for Acute Infectious Disease Epidemiology (CAIDE) staff and CAIDE scholars who participated in the RI tick-borne surveillance systems, contributing to the 2015-2019 surveillance data presented in this article. This publication was supported by the Cooperative Agreement NOFO number CDC-RFA-CK19-1904 entitled 2019 Epidemiology and Laboratory Capacity for Prevention and Control of Emerging Infectious Diseases (ELC), funded by the Centers for Disease Control and Prevention. Its contents are solely the responsibility of the authors and do not necessarily represent the official views of the Centers for Disease Control and Prevention or the Department of Health and Human Services. Any use of trade, firm, or product names is for descriptive purposes only and does not imply endorsement by the US government.

Authors

- Howard S. Ginsberg, PhD, is a Research Ecologist with the U.S. Geological Survey, Eastern Ecological Science Center. He serves as Unit Leader of the USGS Rhode Island Field Station, and Professor in Residence at the University of Rhode Island.
- Jannelle Couret, PhD, MEM, is an Assistant Professor in the Department of Biological Sciences, University of Rhode Island.
- Jason Garrett, BSN, MPH, is a Public Health Epidemiologist with the Rhode Island Department of Health.
- Thomas N. Mather, PhD, is a Professor in the Department of Plant Sciences & Entomology, University of Rhode Island.
- Roger A. LeBrun, PhD, is Carnegie Professor of Life Sciences Emeritus, Department of Plant Sciences & Entomology, University of Rhode Island.

Correspondence

Howard S. Ginsberg, PhD
 U.S. Geological Survey, Eastern Ecological Science Center
 University of Rhode Island, Woodward Hall – PSE
 Kingston, RI 02881
hginsberg@usgs.gov

Wasting No Time: Implementation and the Climate Impact of a Solid Waste Stream Process Intervention in a Large Academic Emergency Department

KATELYN MORETTI, MD, MS; REBECCA KARB, MD, PhD; ROGER DURAND; LEO KOBAYASHI, MD; ALISON S. HAYWARD, MD, MPH

INTRODUCTION

Climate change is one of the most critical issues facing our planet, and healthcare systems play a major role in generating the greenhouse gases that lead to accelerated climate change.¹ In fact, healthcare is responsible for 10% of all greenhouse gas emission in the United States² and generates 4.3 million tons of waste annually.³ Moreover, the emergence of the COVID-19 pandemic has created myriad challenges including the increased production of biomedical waste.⁴

Meaningful solutions to reduce the impact of human activity on climate change must include efforts by healthcare systems to reduce waste and decrease the production of greenhouse gases. In particular, biohazard waste, also known as regulated medical waste (RMW), requires autoclaving and shredding and results in up to fifteen times as much greenhouse gas equivalents emitted as compared to regular solid waste.^{5,6} In addition, RMW processing is seven times more costly to healthcare systems as compared to regular solid waste processing.

RMW is typically defined as items saturated in blood or potentially infectious bodily fluids, sharps, and syringes and is generally collected in red biohazard bags and sharps containers. RMW comprises just a fraction of all waste produced within a medical system, yet idiosyncratic institutional practices (such as the sole use of red bags for all medical waste disposal) can adversely affect the environment at a disproportionate level. In order to mitigate this particular problem in the Emergency Department (ED) at Rhode Island Hospital (RIH), a multi-disciplinary departmental “Green Team” planned and implemented a pilot intervention, i.e., the introduction of a regular solid waste disposal “clear bag” option with accompanying educational efforts and materials on proper waste sorting.

Similar efforts have been successful in other healthcare settings: Inova Fairfax Hospital saved nearly \$200,000 in annual waste disposal fees through better segregation of waste and a concerted effort to educate and engage staff;⁷ a study by Garcia et al. reported a 2-million pound decrease in biohazard waste with a corresponding savings of \$696,000 in annual costs through a similar hospital-wide initiative at Brookdale University Hospital and Medical Center.⁸ However, we are not aware of initiatives that have been specific to the ED – this may be due to the positioning of the ED as a unique and high-risk environment with respect to waste

Figure 1. Regulated medical waste and tan regular solid waste bins.



disposal, given the nature of its clinical operations and associated challenges.

In this context, the Green Team aimed to decrease its ED greenhouse gas footprint as well as operational costs by introducing new tan waste bins with clear bags for regular solid waste disposal (alongside the existing red waste bins and red bags for biohazard waste, **Figure 1**). In addition, educational materials were created and disseminated to ED staff by the Green Team to promote proper waste disposal one month prior to rollout of the project. In parallel with these processes, the team planned and conducted an objective analysis of the environmental impact and fiscal savings of this change within the ED. This manuscript describes the project’s design, conduct, findings, and measured impact.

METHODS

Baseline Analysis

Green Team members met with institutional personnel, including environmental service staff to understand the existing waste stream processes. After meetings and direct observation, the project team conducted a baseline analysis by weighing the ED waste stream (comprising RMW and regular solid waste) over 7 days and then 14 days at two time points prior to implementation of the project’s pilot intervention.

Education

To ensure that all ED technicians, nurses, advanced practice providers (APPs) and physicians understood proper disposal processes as well as the implications of correct waste

disposal, the Green Team created educational materials for distribution. The Green Team developed posters to be placed on waste bins and in utility rooms, as well as slide shows with a brief overview of the planned intervention to display at education sessions. All materials were reviewed and approved by the institution's department of infection control. In-person education sessions were held at faculty meetings, resident conferences, APP meetings, and nursing and technician huddles; emails detailing the project were sent to all staff. Education sessions were also held with all environmental services staff who were briefed on waste disposal process updates as well as how to report RMW that was erroneously discarded in the regular solid waste stream (solid waste within the RMW stream was not reported). Instances of incorrect disposal of RMW were reported via an online reporting form for review by the Green Team and the institutional safety officer.

The multi-disciplinary composition of the Green Team facilitated staff buy-in and adherence to the new waste stream process. Dissemination of project goals and new waste disposal guidelines to staff began one month prior to implementation of the new process.

Project Solid Waste Stream Process Intervention

One month following the second baseline weighing, 105 tan bins with clear bags were added to ED patient rooms and areas (Figure 1). (Pre-existing tan bins which had been lined with red bags received clear bags.) Laminated signs reminding staff of appropriate waste disposal guidelines were affixed to the lids of the tan bins within patient rooms (Figure 2). A third ED waste stream weighing (for RMW only) was completed over a 3-week intervention period. (Figure 3)

Data Analysis


Daily ED red bag weights for the pre-intervention period (RMW and regular solid waste) and for the intervention period (RMW only) were calculated with correction for ED census (patients who left prior to full medical evaluation were excluded from the census). Given that no interventions were implemented to change provider behaviors or institutional processes with respect to medical care and the resultant waste, the quantity of overall, total census-adjusted ED daily waste was presumed to remain the same, such that any red bag weight reductions in the intervention period were ascribed to the diversion of regular solid waste into the new clear bags and tan bins. A one-sided t-test was employed to compare the pre-intervention and intervention periods' red bag weights.

Greenhouse gas equivalents were calculated using the M+WasteCare Calculator.⁵ The calculator used the amounts

Figure 2. Educational poster attached to tan bins placed in the ED for regular solid waste handling.



Figure 3. ED waste weights (lbs) per patient before and during introduction of a solid waste stream process intervention.



of pollutants in each step of the disposal process and converted them to carbon dioxide equivalents (greenhouse gas equivalents). Emissions factors were used for all calculations, including waste transportation to landfill and emissions associated with the landfill. The difference in greenhouse gas equivalent tons per year (TPY) for autoclaved RMW vs regular solid waste was calculated and applied to the observed and annual projected reductions (based on the

2020 ED census) in RMW production. RMW was classified as autoclaved on-site prior to being landfilled. Cost savings were estimated using the reduction in daily ED red bag waste weights between the two periods and with adjustment for accompanying increase in solid waste processing costs.


RESULTS

During the pre-intervention measurement periods, the average red bag waste produced per ED patient was 3.0 lbs (95% CI 2.5-3.5). Over the intervention period, the average red bag waste produced per ED patient was 2.3 lbs (95% CI 1.8-2.8, $p=0.02$), resulting in a net 0.7 lb reduction in biohazard waste per ED patient. Assuming no change in overall waste production per patient, regular solid waste constituted 23.3% of all waste generated during the intervention period. Using the 2020 ED census of 84,221 encounters, the projected annual RIH ED diversion of biohazard waste for 2021 will be 62,323 lbs (31.2 tons). The projected reduction in ED waste greenhouse gas equivalents is 942 TPY; the projected savings in ED waste processing costs is \$11,841 per year. As of three months after the intervention period, there were no reported incidents of RMW within the regular solid waste stream.

DISCUSSION

Institution of a regular solid waste stream within a large academic ED resulted in significant reductions in greenhouse gas emissions. Projections indicate that in just the

Figure 4. Educational poster distributed after institution of the solid waste stream process intervention to inform employees of the project's impact.



subsequent year, emissions prevented will be equivalent to over 2 million miles driven by a passenger car, 1,980 barrels of oil, or over 11 tanker trucks worth of gas (Figures 4).¹⁰ Furthermore, the project's pilot intervention was found to be fiscally beneficial for the hospital system, with return on investment from start-up costs reached within nine months.

While the 23% reduction in actual RMW by the intervention represents a significant decrease for our ED, waste audits of other EDs suggest room for improvement. For example, a recent waste audit of a large academic ED in Boston demonstrated RMW was 10.7% total waste⁹, and a similar audit of a community ED in Rhode Island found just 3% RMW. We expect greenhouse gas savings to improve over time with continued education and institutional adoption of eco-friendly waste disposal practices.

Limitations

Given the process by which hospital waste is measured at our institution, it was not possible to isolate and weigh only the regular solid waste component. Thus, calculations assumed the same total generation of waste on a per-patient basis. A formal waste audit was not performed to calculate the proportions of appropriate waste versus inappropriate waste within the RMW stream – a formal audit of the RMW stream in the future would allow for tailored education and further interventions to continue to reduce RMW generated.

Due to the architectural configuration and operational layout of the ED, certain patient care areas such as the 12-bed critical care patient space were not included in the project – there is significant healthcare waste produced in this area, and there exists the potential for substantial additional greenhouse gas reductions and cost savings. Future research will examine the amount and type of waste generated from this area as well as potential interventions to reduce impact.

The M+WasteCare calculator⁵ uses known formulas for carbon and other pollutant generation from medical waste. Variables added to the calculator include distance to landfills, type and frequency of waste transport. There may be slight variations in the actual amount of greenhouse gases produced based on these variables. However, because the variables were constant across the measured periods, the impact of the intervention should not have been affected.

CONCLUSIONS

Proper sorting of RMW and regular solid waste within EDs represents a straightforward, economical, and impactful environmental intervention. While risk to the healthcare system exists in the form of waste misclassification, from this intervention, this appears to be minimal and outweighed by the advantages. In light of the COVID-19 pandemic and the related increase in biomedical waste, conscientious disposal practices will be even more important to the environmental sustainability of our healthcare systems.

References

1. The 2020 report of The Lancet Countdown on health and climate change: responding to converging crises - The Lancet. [cited 2021 Apr 21]; Available from: [https://www.thelancet.com/journals/lancet/article/PIIS0140-6736\(20\)32290-X/fulltext](https://www.thelancet.com/journals/lancet/article/PIIS0140-6736(20)32290-X/fulltext)
2. Eckelman MJ, Sherman JD. Estimated Global Disease Burden From US Health Care Sector Greenhouse Gas Emissions. *Am J Public Health*. 2018;108(Suppl 2):S120–2.
3. Howard J. Less waste how-to guide. Practice Greenhealth. 2019. [Accessed November 17, 2020]. Available at: https://practicegreenhealth.org/sites/default/files/2019-03/hhi-booklet-waste-bu_v3.0_full.pdf
4. Rume T, Islam SDU. Environmental effects of COVID-19 pandemic and potential strategies of sustainability. *Heliyon*. 2020;6(9): e04965.
5. M+WasteCare Calculator [Internet]. [cited 2020 Jan 6]; Available from: <https://wastecare.mazzetti.com/calculator>
6. Health-care waste. World Health Organization [Internet]. [cited 2020 Jan 6]; Available from: <https://www.who.int/news-room/fact-sheets/detail/health-care-waste>
7. Regulated medical waste | Practice Greenhealth [Internet]. [cited 2020 Jan 6]; Available from: <https://practicegreenhealth.org/topics/waste/regulated-medical-waste>
8. Garcia R. Effective cost-reduction strategies in the management of regulated medical waste. *Am J Infect Control*. 1999;27(2): 165–75.
9. Hsu S, Thiel CL, Mello MJ, Slutzman JE. Dumpster Diving in the Emergency Department. *West J Emerg Med*. 2020;21(5):1211–1217. doi:10.5811/westjem.2020.6.47900
10. US EPA O. Greenhouse Gas Equivalencies Calculator [Internet]. US EPA. 2015; available at: <https://www.epa.gov/energy/greenhouse-gas-equivalencies-calculator>

Acknowledgment

The authors acknowledge the multidisciplinary members of the Anderson Emergency Department Green Team for their pivotal role in designing and implementing this project.

The authors also acknowledge Ananya Suram for her design and creation of Figure 4.

Authors

- Katelyn Moretti, MD, MS, Assistant Professor of Emergency Medicine, Alpert Medical School of Brown University; Rhode Island Hospital, Providence, RI.
- Rebecca Karb, MD, PhD, Assistant Professor of Emergency Medicine, Alpert Medical School of Brown University; Rhode Island Hospital, Providence, RI.
- Roger Durand, Director, ESD & Textile Services at Lifespan Rhode Island Hospital.
- Leo Kobayashi, MD, Professor of Emergency Medicine, Alpert Medical School of Brown University; Rhode Island Hospital, Providence, RI.
- Alison S. Hayward, MD, MPH, Assistant Professor of Emergency Medicine, Alpert Medical School of Brown University; Rhode Island Hospital, Providence, RI.

Correspondence

Katelyn Moretti, MD, MS
katelyn_moretti@brown.edu

Trends in Waste Production at a Community Hospital during the COVID-19 Pandemic

KYLE DENISON MARTIN, DO, MA, MPH, DTM&H; JANE J. CHEN, BS, MD'24; JAMIE THORNDIKE, BS, MD'24; WINSTON MCCORMICK, BS, MD'23; JOHN ROTA, BA; BRIAN BERG, DO; ANNIE DULSKI, DO

ABSTRACT

INTRODUCTION: As of September 2021, the COVID-19 pandemic has led to 42,500,000 cases and 680,000 deaths in the United States. In Rhode Island, there have been 170,000 cases and 2,820 deaths. Investigating resource utilization and waste production during disease outbreaks can inform efforts at disaster preparedness. The purpose of this study was to examine trends in waste production during the COVID-19 pandemic.

METHODS: This is a descriptive study examining trends in waste production during the COVID-19 pandemic. The study was conducted at a suburban community hospital in Rhode Island. Data was collected on regulated medical waste (RMW) and linen use from October 2019–July 2021. Adjusted patient days (APD) values were calculated using hospital census and revenue data. Total weight and weight/APD were calculated for each month of the study period. Data was then compared with overall COVID-19 cases and hospitalizations in Rhode Island. This data was gathered from the Rhode Island Department of Health (RIDOH) COVID Response Data Hub.

RESULTS: Regulated Medical Waste (RMW) by total weight was lowest in April 2020, when the hospital census and adjusted patient days (APD) were at their lowest. In contrast, linen use remained largely consistent with pre-pandemic levels during the initial months of the pandemic despite a decrease in hospital census. The highest linen weight/APD value (23.32 lbs/APD) was in April 2020. Both RMW and linen use (weight/APD) decreased during the study period. Linen use was highest during months with increased COVID-19 cases and hospitalizations.

CONCLUSIONS: This study examined trends in waste production at a community hospital during the COVID-19 pandemic. Linen use was highest during months of increased COVID-19 cases and hospitalizations, while RMW production decreased. There was a particular increase in linen use in April 2020, when the pandemic was in its initial phases.

KEYWORDS: COVID-19, waste, medical waste, pandemic, disaster preparedness

INTRODUCTION

As of September 2021, the COVID-19 pandemic has led to 42,500,000 cases and 680,000 deaths in the United States.¹ In Rhode Island, there have been 170,000 cases and 2,820 deaths.^{2,3} Sophisticated hospital based patient care requires an enormous investment in resources, but results in waste production. Understanding how waste is produced can inform disaster preparedness efforts for future pandemics.

Most waste from the ED is considered municipal solid waste (MSW) and can be disposed in a landfill. It requires no additional treatment prior to transport. Disposal of regulated medical waste (RMW), which includes most bodily fluids such as blood and sputum suspected to carry infectious pathogens, may potentially increase when a new and unfamiliar disease emerges. The disposal of RMW requires strict adherence to specific procedures and, as such, is expensive and burdensome to the environment. Variables that affect the carbon footprint of waste production and removal include the type of waste, the method and site of disposal, and the means of conveyance.

The global response to waste production underscores the necessity of planning during a pandemic. In Romania, government-imposed restrictions on the disposal of waste from health care and designated quarantine facilities led to illegal dumping of medical waste.⁴ In Indonesia, medical waste increased during the initial months of the pandemic; however, there was a lack of appropriate facilities for the disposal of this waste.⁵

Understanding trends in RMW during outbreaks of novel disease can guide preparation among health systems and government officials and can inform disaster preparedness efforts for future pandemics. As such, the purpose of this study was to examine trends in waste production at a suburban community hospital during the COVID-19 pandemic. Investigators focused particularly on RMW and linen use during the study period.

METHODS

This is a descriptive study examining trends in waste production during the COVID-19 pandemic. The study was conducted at a suburban community hospital in Rhode Island with 346 inpatient beds and an annual Emergency Department (ED) volume of approximately 70,000 patient

encounters. The hospital has an Intensive Care Unit (ICU) and Progressive Care Unit (PRG) that provide services for critically ill patients. During the COVID-19 pandemic, the number of beds in these units varied based upon patient volume, infection control policies and care requirements. The hospital is home to several graduate medical training programs, including residencies in Family Medicine, Internal Medicine and Emergency Medicine.

Data was collected on regulated medical waste (RMW) and linen use from October 2019–June 2021. This time period was selected since it included six-months prior to the beginning of the COVID-19 pandemic. RMW and linen were selected as categories for analysis because of their pertinence to infectious control. RMW included all items placed in red biohazard bags. This did not include sharps. It did not include waste that could be categorized as RMW inappropriately placed in non-RMW containers. Linens included patient gowns and bedding as well as reusable isolation gowns. The hospital in this study utilizes reusable isolation gowns for patients who meet certain infection control criteria. These gowns are laundered after use and subsequently returned into circulation.

Environmental Services (EVS) gathers information on various categories of hospital waste, including sharps, linens, municipal solid waste (MSW), and RMW. These materials are weighed daily and recorded monthly. Adjusted patient days (APD) were calculated using hospital census and revenue data per criteria established by the Centers for Medicare & Medicaid Services (CMS). Weight per APD (lbs/APD) was then calculated for RMW and linens. These data points were plotted and compared to overall cases and hospitalization due to COVID-19 in Rhode Island. This information was gathered from the Rhode Island Department of Health (RIDOH) COVID Response Data Hub.

RESULTS

RMW reached its lowest total weight (5,726 lbs) in April 2020 and highest total weight (14,052 lbs) in June 2020. RMW weight/APD hit its low in January 2021 (0.71 lbs/APD) and high in June 2020 (1.54 lbs/APD). The overall trend of RMW weight mirrored that of APD during the study period (Figures 1 & 2).

By total weight, RMW was at its lowest levels during April and May 2020 when the hospital census was lowest. In fact, RMW decreased by about 50% from pre-pandemic levels during these months. RMW increased as hospital volumes grew during the COVID-19 pandemic.

Linen reached its lowest total weight (119,078 lbs) in May

Figure 1. Total Weight (lbs) of Regulated Medical Waste (RMW) by Month




Figure 2. Adjusted Patient Days (APD) by Month





Figure 3. Trend in Linen Weight Per Adjusted Patient Day (lbs/APD)



2020 and its highest total weight (205,563 lbs) in December 2020. Linen weight/APD hit a low (11.91 lbs/APD) in May 2021 and peaked (23.32 lbs/APD) in April 2020 (Figure 3).

When averaged over three-month periods from Oct 2019–June 2021, RMW weight per APD showed a steady decline while linen weight per APD increased during the period of April–June 2020 followed by a steady decline (Table 1).

Rhode Island has experienced several waves of COVID-19 infection, with total cases reaching their peak in April 2020, December 2020 and March 2021 (Figure 4). Hospitalizations due to COVID-19 reached their highest levels in April 2020, December 2020 and April 2021 (Figure 5). The largest number of patients hospitalized in the ICU was in April 2020.

Table 1. Average RMW and Linen Weight per APD (lbs/APD) by Three-Month Period

	RMW (lbs/APD)	Linen (lbs/APD)
Oct–Dec 2019	1.20	14.99
Jan–March 2020	1.21	15.83
April–June 2020	1.11	17.53
July–Sept 2020	1.10	15.71
Oct–Dec 2020	1.02	15.08
Jan–March 2021	0.96	14.57
April–June 2020	0.95	12.00

DISCUSSION

This study identified several trends in RMW and linen use during the COVID pandemic. RMW by total weight was lowest in April 2020, when the hospital census and adjusted patient days (APD) were at their lowest. In contrast, linen use remained largely consistent with pre-pandemic levels during the initial months of the pandemic despite a decrease in hospital census. This led to higher weight/APD values, including the highest linen weight/APD value (23.32 lbs/APD) in April 2020. However, linen use (weight/APD) decreased after this initial rise.

Linen use was highest during months with increased

Figure 4. Number of Daily New Positive Cases of COVID-19 in Rhode Island,¹⁰ April 1, 2020–September 15, 2021.





Figure 5. Number of Hospitalizations Due to COVID-19 in Rhode Island¹¹



COVID-19 cases and hospitalizations. This is most pronounced in April 2020 at the beginning of the pandemic. This was also when ICU hospitalizations due to COVID-19 were at their highest level in Rhode Island. Overall linen use (by total weight) was highest in December 2020, when COVID-19 cases and hospitalizations reached their highest levels in Rhode Island.

Findings from this study are similar to those from other sites, including Wuhan, China, where the COVID-19 pandemic originated. During the initial phase of outbreak in Wuhan (January–April 2020), health care waste per 1,000 persons increased from 3.64 kg/day to 27.32 kg/day.⁶ To accommodate this increase in waste production, disposal capacity in the region was increased from 50 tons/day to 280 tons/day. Researchers in Malaysia found that clinical waste (which includes any item potentially contaminated with infectious material) increased by 27% during the initial months of the COVID-19 pandemic.⁷ This increase was mainly attributed to the disposal of PPE, such as gloves, facemasks and gowns. In Romania, waste from medical facilities peaked in the early months of the pandemic and then tapered over the following months.⁵

Several studies from Taiwan have examined waste use during 2003 SARS outbreak. A 2,000-bed hospital in northern Taiwan found that daily infectious waste generation tripled during the outbreak.⁸ Interestingly, due to a decrease in hospital census, the mean daily overall waste production decreased 19.2%–25.3%. Taiwan saw an overall increase in infectious waste of about 4,000 tons (from 14,648 to 19,350 tons) following SARS.⁹

There are several limitations to this study, most notably that it was conducted at a single site. Data from this study came from a suburban community hospital in Rhode Island. Resource utilization might be different at a rural or urban hospital site. This is notably important in the case of COVID-19 since the pandemic affected different regions of the United States in very different ways. Some hospitals might have only had a few overall cases of disease, whereas others might have been overwhelmed to the point where they needed to transfer patients requiring ICU level-of-care to another facility. Hospital infection control policies likely also influenced resource utilization. A hospital that uses disposable gowns for PPE might have different patterns of use when compared to a hospital that uses reusable gowns. Another limitation of this study is aggregate data on COVID-19 hospitalizations on a state level was used for comparison. Hospital-level data and department-level data (particularly from the ED and ICU) would be more useful. Finally, a field hospital provided care to patients with mild-to-moderate cases of COVID-19 during the study period. Resource utilization at the field hospital was not included in this study.

This study examined the overall trends of RMW and linen use during the COVID-19 pandemic. The dramatic increase in linen use (weight/APD) during April 2020 is of

particular interest. Several factors could have played a role in this occurrence, including infection control policies, staff compliance with isolation precautions, lack of knowledge of disease transmission, or fear amongst hospital employees of a new and unknown disease. Trends suggest that there may be an association with increased linen use and ICU hospitalizations. However, further investigation is needed to clarify this association. The gradual decrease in linen use (weight/APD) since April 2020 is also of interest. Are hospitals becoming more efficient with resource utilization or have staff developed noncompliance with infection control protocols as the COVID-19 pandemic drags on? Qualitative studies could provide insight. Overall, while this study highlighted trends that hint at correlation and causation, further research is needed to clarify these relationships.

CONCLUSION

This study examined trends in waste and resource utilization at a community hospital during the COVID-19 pandemic. While largely hypothesis-generating, it hints at several conclusions, including a relationship between linen use and ICU hospitalizations. Findings from this study can contribute to further understanding of resource utilization during outbreaks of disease and inform health system and government protocols for pandemic preparedness.

References

- Centers for Disease Control and Prevention (CDC). "COVID-19." Updated on Sept 24, 2021. Accessed on Sept 24, 2021. <https://www.cdc.gov/coronavirus/2019-ncov/index.html>
- Rhode Island Department of Health. "Rhode Island COVID-19 Response Data." Updated on Sept 27, 2021. Accessed on Sept 24, 2021. <https://ri-department-of-health-covid-19-data-rihealth.hub.arcgis.com>
- The New York Times. "Tracking Coronavirus in Rhode Island: Latest Map and Case Count." Updated on Sept 27, 2021. Accessed on Sept 24, 2021. <https://www.nytimes.com/interactive/2021/us/rhode-island-covid-cases.html>
- Mihai F. "Assessment of COVID-19 Waste Flows During the Emergency State in Romania and Related Public Health and Environmental Concerns." *Int J Environ Res Public Health*. July 2020;17(15):5439.
- Anwar F. "The Medical Waste in Jakarta is Predicted to Reach 12,750 Ton Because of Corona Outbreak." *Detik Health*. 2020. Accessed on Sept 24, 2021. <https://health.detik.com/berita-detikhealth/d-4986675/limbah-medis-di-jakarta-diprediksi-capai-12750-ton-karena-wabah-corona>.
- Yang L, Yu X, Wu X, et al. "Emergency response to the explosive growth of health care wastes during COVID-19 pandemic in Wuhan, China." *Resour Conser Recycl*. Jan 2021;164: 105074.
- Agamuthu P, Barasarathi J. "Clinical waste management under COVID-19 scenario in Malaysia." *Waste Manag Res*. June 2021;39(1_suppl): 18-26.
- Chiang CF, Sung FC, Chang FH, et al. "Hospital waste generation during an outbreak of severe acute respiratory syndrome in Taiwan." *Infect Control Hosp Epidemiol*. May 2006; 27(5): 519-522.
- Huang M, Lin J. "Characteristics and management of infectious industrial waste in Taiwan." *Waste Management*. Nov 2008. 28(11):2220-2228

10. Rhode Island Department of Health. "Rhode Island COVID-19 Cases Data." Updated on Sept 27, 2021. Accessed on Sept 24, 2021. <https://ri-department-of-health-covid-19-case-data-rihealth.hub.arcgis.com>.
11. Rhode Island Department of Health. "Rhode Island COVID-19 Hospitalization Data." Updated on Sept 27, 2021. Access on Sept 24, 2021. <https://ridoh-covid-19-response-hospital-data-rihealth.hub.arcgis.com>

Authors

Kyle Denison Martin, DO, MA, MPH, DTM&H, Academic Faculty, Department of Emergency Medicine, Kent Hospital, Warwick, RI.

Jane J. Chen, BS, MD'24, Warren Alpert Medical School of Brown University, Providence, RI.

Jamie Thorndike, BS, MD'24, Warren Alpert Medical School of Brown University, Providence, RI.

Winston McCormick, BS, MD'23, Warren Alpert Medical School of Brown University, Providence, RI.

John Rota, BA, General Manager, Department of Environmental Services, Kent Hospital, Warwick, RI.

Brian Berg, DO, Emergency Medicine Resident, Kent Hospital, Warwick, RI.

Annie Dulski, DO, Academic Faculty, Department of Emergency Medicine, Kent Hospital, Warwick, RI.

Correspondence

Kyle Denison Martin, DO, MA, MPH, DTM&H
Kent Hospital
455 Toll Gate Rd
Warwick RI 02886
kdenisonmartin@brown.edu

Trash Talk in the ED: Takeaways from Waste Audits at New England Hospitals

KYLE DENISON MARTIN, DO, MA, MPH, DTM&H; WINSTON MCCORMICK, BS, MD'23;
JULIA CAPACCI, DO, MS; KATELYN MORETTI, MD, MS

KEYWORDS: sustainability, environment, climate change & health, waste

Recent investigations at Kent Hospital, a suburban community hospital in Rhode Island, and Massachusetts General Hospital (MGH), a Level 1 trauma center, highlight the importance of health care waste.^{1,2} These investigations involve a simple concept: sorting through trash can help identify ways to cut costs and reduce environmental impact.

The waste audits – both conducted in the Emergency Department (ED) – followed a similar protocol. Environmental Services staff collected waste from the ED for a 24-hour period. Waste was then sorted into various categories, including paper, plastic, food waste, glass, metal, electronic material and unused items. Most waste from the ED is considered municipal solid waste (MSW) and requires no additional treatment and can be disposed in a landfill. Regulated Medical Waste (RMW), on the other hand, requires specific protocols prior to disposal. It is expensive and often requires energy intensive processes such as autoclave sterilization or incineration.

Attention was given to whether Regulated Medical Waste (RMW) and sharps were appropriately disposed (i.e., in designated containers). Investigators also examined if non-RMW was inappropriately placed in RMW containers. All categories of waste were weighed and direct pollutant emissions were calculated using the M+ WasteCare Calculator.³ The waste audits at the two facilities revealed some important takeaways:

(1) Inappropriate disposal of Regulated Medical Waste (RMW) is expensive and inefficient

While the amount of waste generated per-patient encounter was similar at Kent and MGH, estimated carbon emissions were 10 times higher at MGH. The difference in emissions was driven by differences in RMW. The waste audits identified 71.67 kg in RMW containers at MGH compared with 4.67 kg at Kent. Disposing of this waste accounted for 70% of overall waste emissions at MGH. This is because there are specific requirements for disposing of RMW that tend to be energy intensive. RMW needs to be rendered safe prior to disposal in a landfill. Hospitals can perform on-site sterilization or utilize a specialized hauling service to have waste processed off-site.

Only 15% of waste disposed in RMW containers at MGH met criteria for RMW. Eighty-five percent of material in RMW containers could have been transported to a landfill but, instead, was processed using the energy-intensive disposal methods reserved for RMW. Behavioral characteristics are partly responsible for inappropriate use of the RMW containers. Improvements in ED staff knowledge regarding RMW criteria and ED design simplifying RMW bin location could lead to more appropriate disposal.

(2) ED waste is filled with unused items

Audits at Kent and MGH demonstrated that unused items tossed in the garbage are a significant issue in the ED. More than 170 unused items were identified in the Kent audit, consisting of 5.2% of total waste. These items included unused boxes of gloves, surgical face masks, suture material and medications (Table 1). The audit at MGH identified 201 unused items, including 76 bundled and unused tourniquets. Both audits revealed resuscitation supplies still in storage packaging. Unused items tossed in the garbage are pure waste. There are monetary and environmental costs to producing these items, transporting them to the hospital and then disposing of them in a landfill.

It is unclear as to why there are so many unused items in ED waste. Infection control policies may play a role. Unused items may be considered “contaminated” if present in a room under isolation precautions. In addition, procedure kits often contain redundant or useless items, leading to the disposal of unused items. Evidence of unused resuscitation supplies in the Kent and MGH audits suggests that resuscitation scenarios are particularly prone to waste. There is likely also a behavioral component to the disposal of unused items. In a busy ED, it is easier to toss unused supplies into the garbage rather than placing them back in a drawer or supply closet.

Potential solutions for the problem of unused items include only bringing necessary supplies into the room when caring for patients on isolation precautions and designing “in-house” procedure kits with items pre-approved by ED staff. Kits for certain procedures (such as laceration repairs) could be sterilized in-house and then returned into circulation. ED administration could incentivize re-stocking of unused supplies. Specific ED staff could also be tasked with a re-stocking role. As with RMW disposal, ED design should

Table 1. A List of Unused Items Identified during the Kent ED Waste Audit

2 unopened boxes of non-sterile gloves
2 partially finished boxes of non-sterile gloves
3 unopened sets of sterile gloves
8 unopened female catheterization kits
6 unopened Yankauer suction handles
8 unopened suction tubing sets
1 unopened sterile drape
2 unopened 4x4 sponges
4 unused (and folded) adult diapers
7 unopened urine sample collection kits
13 unused bedpans
1 unopened Tegaderm
1 unopened C-collar
2 unused bags of IV fluids
12 unopened normal saline flushes
3 unopened IV catheters
10 unopened pairs of socks
1 partially finished box of procedural masks
2 sterile large cotton swap applicators
1 sterile solution bowl
4 unopened electrode sets
1 unused suction container
1 unopened ambu-bag
34 unopened alcohol prep pads
6 unopened small gauze packets
6 unopened ChloroPrep swab kits
1 unopened pill of Lorazepam
13 unused emesis basins
2 unopened Iodine prep pads
6 unused large pink irrigation containers
26 unused Castille soap towelettes
1 unused bag of IV antibiotics
Multiple unopened food items

also be taken into account. The layout of patient rooms and the ED should make it as easy as possible for staff to restock unused items instead of tossing them in the trash.

(3) How and where we process waste matters

Waste disposal sites and their proximity to the hospital are important to consider. Gunderson Health System in Wisconsin treats 90% of RMW through an in-house sterilization system powered by steam from the heating and cooling system at one of its buildings.^{4,5} After sterilization, waste is transported just four miles to a waste-to-energy facility. Previously, waste was shipped to a landfill 1,250 miles away.

CONCLUSION

Waste audits can help make EDs and health systems more cost-efficient and environmentally friendly. While health-care providers pledge *primum non nocere* – do no harm – pollutants and wasteful practices adversely affect our patients. We need to continue to examine hospital waste habits and embrace solutions that provide healthcare without harm.

References

1. Hsu S, Thiel CL, Mello MJ, et al. Dumpster diving in the Emergency Department: quantity and characteristics of waste at a level 1 trauma center. *West J Emerg Med.* 2020;21:1211-1217.
2. Hsu S, Banskota S, McCormick W, et al. Utilization of a waste audit at a community hospital emergency department to quantify waste production and estimate environmental impact. *The Journal of Climate Change and Health.* October 2021;4: 10041.
3. <http://wastecare.mazzetti.com/>.
4. Kelly M. Reducing the Emergency Department's Footprint: Greening the ED. *Annals of Emergency Medicine.* September 2021;78(3):15A-18A.
5. Khamayzer A. The profitable hospital system with sustainability in its DNA. *GreenBiz.com.* Published on April 10, 2017. <https://www.greenbiz.com/article/profitable-hospital-system-sustainability-its-dna>

Authors

- Kyle Denison Martin, DO, MA, MPH, DTM&H, Warren Alpert Medical School of Brown University, Providence, RI; Department of Emergency Medicine, Kent Hospital, Warwick, RI.
- Winston McCormick, BS, MD'23, Warren Alpert Medical School of Brown University, Providence, RI.
- Julia Capacci, DO, MS, Department of Emergency Medicine, Kent Hospital, Warwick, RI.
- Katelyn Moretti, MD, MS, Warren Alpert Medical School of Brown University, Providence, RI.

Correspondence

Kyle Denison Martin, DO, MA, MPH, DTM&H
Kent Hospital
455 Toll Gate Rd, Warwick RI 02886
kdenisonmartin@brown.edu

The Rhode Island Climate Change and Health Program: Building Knowledge and Community Resilience

RACHEL CALABRO, MS; CAROLINE HOFFMAN, MPH

ABSTRACT

Climate change acts as a risk multiplier, meaning vulnerable populations bear a disproportionate burden of its effects. Improving climate resiliency is a key strategy to help the Rhode Island Department of Health meet its overarching goals of addressing the socio-economic and environmental determinants of health for all Rhode Islanders. The Climate Change and Health Program focuses on both the immediate health impacts of climate change and building resiliency. Part of the US Centers for Disease Control and Prevention's Climate Ready States and Cities Initiative, the Program has partnered with community groups and other state and local agencies to bring technical assistance, educational resources, and funding to support community resilience to the challenges presented by the already changing climate. Specific projects discussed include the extreme heat communications plan and outdoor worker campaign; community-driven resiliency projects in response to flooding and natural hazards, and improving resilience in senior citizen housing.

KEYWORDS: climate change, resilience, public health, extreme heat, climate mitigation

INTRODUCTION

Climate change, health, and equity are inherently intertwined.¹ The impact of race and socio-economic status on health has been evident in the disparate outcomes seen across populations during the COVID-19 pandemic.² These same communities bear a disproportionate burden of the effects of climate change.¹ Because climate change is a risk multiplier, improving climate resiliency is a key strategy to help the Rhode Island Department of Health (RIDOH) meet its overarching goals of addressing the socio-economic and environmental determinants of health for all Rhode Islanders, especially those from the most vulnerable demographics. As incidences of heatwaves and flooding increase, RIDOH focuses on addressing immediate health impacts and building resiliency among Rhode Islanders.³

MAKING RHODE ISLAND A CLIMATE READY STATE

The RIDOH Climate Change and Health Program began as part of the US Centers for Disease Control and Prevention's (CDC) Climate Ready States and Cities Initiative. This initiative is part of the CDC's efforts to train medical and public health professionals to better understand the impact of climate change on health.⁴ The CDC provides resources for the professional community that include research papers, maps, and links to webinars from the American Public Health Association covering climate-related topics such as heart and lung health, children's health, mental health, and allergies and asthma.

Two notable data resources include the Heat and Health Tracker and localized climate change and health data. The Heat and Health Tracker is an online tool to help emergency and public health planners prepare for and respond to extreme heat events by providing local-level heat and health data.⁵ Localized data for climate change and health, including future scenarios for heat and precipitation to the year 2099, can be found on the CDC's National Environmental Public Health Tracking Network portal.⁶

The Rhode Island Climate Change and Health Program began in 2013 and continues to focus on health effects from urban heat, flooding, severe weather and sea level rise, food and water borne diseases, vector-borne diseases, and poor air quality. (See **Figure 1** for more information about the health effects of climate change.) The program has partnered with community groups and other state and local agencies to bring technical assistance, educational resources, and funding to support community resilience to these challenges.

ADDRESSING EXTREME HEAT IN RHODE ISLAND

Extreme heat has become an increasing threat across Rhode Island as the average temperature has already risen three degrees in the last century.³ During the climate normal period (1981 to 2010), there was an average of 9.3 days equal to or above 90 degrees Fahrenheit (F) in the Providence Metro Area each year.⁷ There is variability in this number year-to-year. While there was an average of 11 such days each year from 2010 to 2014, there were 19 such days in 2021 (see **Figure 2**). It could increase to as many as 27 days at or above 90 degrees F per year by mid-century with slow action to reduce greenhouse gas emissions.⁸

Figure 1. Climate change causes a range of direct and indirect impacts on human health. This figure depicts the most significant climate change impacts (rising temperatures, more extreme weather, rising sea levels, and increasing carbon dioxide levels), their effect on exposures, and the subsequent health outcomes that can result from these changes. (US Centers for Disease Control and Prevention, 2021.)





Figure 2. In the Providence Metro Area, there are an average of five more days per year above 90 degrees than there were in 1970. (Image provided by Climate Central.)



Extreme heat events are a leading cause of weather-related injury and can result in the worsening of existing illnesses. Extreme heat can cause increases in cardiovascular events leading to hospitalization.⁹ When Rhode Island sees multiple days of temperatures above 80 degrees F, emergency room visits begin to rise.¹⁰

Alarmingly, a recent study shows that one quarter of the US population had heat-related symptoms during the summer 2020.¹¹ This was largely due to a lack of cooling resources and the social isolation that was part of the COVID-19 pandemic. The interaction between the two


health crises expanded the scope and range of vulnerability to heat across the country. This research showed that when COVID-19 disrupted social networks, people were less able to seek comfort and suffered more from the symptoms of heat stress.

The effects of heat are not felt evenly across communities. Evidence shows that low-income communities with little tree cover and those that were formerly redlined are up to 12.6 degrees F hotter than surrounding neighborhoods.¹² Redlined areas are those that were coded as “high risk” by the Federal Housing Administration starting in the 1930s. Often based on race, these neighborhoods were denied government loans and services due to land use and other concerns. In Rhode Island, we see similar disparities across neighborhoods in Providence and surrounding urban areas, especially when it comes to heat and tree cover.

To create and implement more effective interventions, the neighborhoods and communities with the highest vulnerability and exposure need to be identified. Collaborating with researchers at CAPA Strategies, based at Portland State University, the Program in partnership with the Department of Environmental Management’s Division of Forest Environment and American Forests mapped ambient air temperatures and humidity in Providence, Pawtucket, Central Falls, and East Providence in July 2020.¹³ See Figure 3. The results showed that many Providence neighborhoods warmed by 10 degrees F more than others, indicating substantially higher exposure to heat. Many areas also stayed warm at night, maintaining temperatures above 70 degrees F and depriving residents of critically important time for the body to recover from heat stress experienced during the day.

To combat the effects of extreme heat in Rhode Island, the Program developed a heat communications plan and outreach for specific vulnerable populations. Implementation

Figure 3. Modeled afternoon temperatures in Providence, Pawtucket, Central Falls, and East Providence, based on data from CAPA Strategies Heat Watch Rhode Island Program, July 29, 2020.



of the extreme heat communications plan includes sending email notices to primary care providers before heat events and using social media to inform the public about heat safety and heat alerts issued by the National Weather Service. Other interventions have included trainings for outdoor workers about the dangers of heat exposure and a *Beat the Heat* campaign with both television and radio advertisements. Grants have been provided to local emergency management officials for portable misting stations; to the Providence Housing Authority and the Town of Barrington for educating seniors about the dangers of heat and natural disasters, and to youth groups for heat-related outreach campaigns.

COMMUNITY-DRIVEN RESILIENCE FRAMEWORK

One way that the Program works to build overall resilience in a community is by building up program supports and social networks. These networks help to create social cohesion and places for residents to turn during an emergency.^{14,15} It is important to empower the communities most affected by climate change to design resiliency programs that address the realities of their lived experience. One way that RIDOH builds community support is through the Health Equity Zone (HEZ) initiative. The HEZ initiative is a place-based approach that brings communities together to address systemic changes at the local level. Each HEZ identifies the unique social, economic, and environmental factors that are preventing people from being as healthy as possible.¹⁶

The Climate Change and Health Program worked with HEZs in Providence, Pawtucket, Central Falls, and Newport to support community resilience to the effects of climate change. Through community workshops, the Program helped residents assess their strengths and vulnerabilities associated with climate change and identify strategies to reduce climate hazards. After the workshops, each team surveyed their own communities to identify strategies and develop a community-led intervention.

The HEZ in the Olneyville neighborhood of Providence chose to create a film about the historic 2010 floods. (The film is available to view at: <https://vimeo.com/359888817/6c75fdddf>). The film aims to raise awareness of the dangers presented by the increasing frequency of natural disasters. It works to transform the threat of a major disaster into a tangible reality so that residents feel a greater sense of urgency regarding emergency preparation. The film was screened at multiple events and will continue to be used in schools and with community groups.

BUILDING RESILIENCE IN RESPONSE TO NATURAL GAS OUTAGE

During a gas outage in January 2019, National Grid was forced to shut down a portion of its gas distribution system

to over 7,000 customers in Newport for more than one week. Evacuations forced people from their homes, and information was limited and inconsistent. This led to a feeling within the community that their needs were not being met by the established emergency management procedures and was fueled by underlying inequities that are often revealed in times of crisis.

During this time, the Newport HEZ became an important part of the effort to share information and reach vulnerable community members. The HEZ leaders realized that there was not a well-defined emergency response plan, nor was there the capacity locally to carry a plan out.

After the event, community conversations about climate change provided opportunities for residents to receive disaster-preparedness training and supplies and to establish relationships with public officials. Creating connections between agencies, officials, and communities increased cohesion and will hopefully lead, both directly and indirectly, to better health outcomes by ensuring that local needs are addressed and plans are in place before disaster strikes.

EMPOWERING COMMUNITY PARTNERS FOR RESILIENT CLIMATE MITIGATION

In Pawtucket and Central Falls, the nonprofit group Groundwork RI partnered with Southside Community Land Trust and Farm Fresh RI's Harvest Kitchen for a six-week summer youth program. These community partners employed and trained 24 youth on how to grow, process, and cook their own local produce. The youth also worked to engage residents and gather data regarding resilient climate mitigation strategies. By the end of the summer, they completed 25 green home assessments, planted 21 trees at Baldwin Elementary school in Pawtucket, installed nine raised garden beds at resident homes, and collaborated with the City of Pawtucket to install and deliver seven residential rain barrels to reduce flooding and utility bills.

IMPROVING EXTREME WEATHER RESILIENCY IN SENIOR HOUSING

Older adults are vulnerable to extreme weather because they often have limited mobility and must shelter in place. This is especially true for those who live in independent senior housing or assisted living. To help make these spaces more resilient to natural disasters and other emergencies such as power loss, the Program partnered with the Yale New Haven Health System Center for Emergency Preparedness and Disaster Response to support long-term care, assisted living, and independent living senior housing facilities in preparing for climate-related disasters through energy resiliency audits and the development of all-hazards emergency plans that emphasize sheltering in place. These important emergency preparedness actions reduce risk to senior citizens

by limiting disruptions like power outages or flooding that can force the evacuation of medically vulnerable people. By increasing the overall emergency preparedness levels of the facilities that serve them and allowing them to shelter in place, seniors are safer. Resources from the project include a facility self-assessment tool, shelter-in-place plan templates, staff trainings, and webinars.¹⁷

PROGRAM RESOURCES AND WEBSITE

Resources are available from the Program for a variety of topics, including extreme heat, air quality, and climate literacy. Partnerships with other RIDOH programs such as the Center for Acute and Infectious Disease Epidemiology have allowed the development of educational tools related to vector-borne diseases like Lyme disease, Eastern equine encephalitis (EEE), and West Nile virus, along with emerging diseases like Zika virus. Links to our reports, brochures, and videos are available on the Program webpage.

The Program has also focused on providing resources to help teachers bring environmental health and climate change into their classrooms. Partnering with school nurse teachers, the two programs have provided thousands of copies of the *Tick Workbook for Kids* to classrooms across the state.

CONCLUSION

The effects of the changing climate in Rhode Island are far-reaching in terms of the scope of the systems impacted, types of impacts, and people who are impacted. At RIDOH, we are continuing these efforts and remain committed to supporting climate change mitigation strategies, preparing for the human health effects, and working closely with community partners to help improve the resilience of all Rhode Islanders.

References

1. Reidmiller DR, Avery CW, Easterling DR, Kunkel KE, Lewis KLM, Maycock TK, Stewart BC. Impacts, risks, and adaptation in the United States: Fourth national climate assessment, volume II. USGCRP. 2018;1515 pp. doi: 10.7930/NCA4.2018.
2. Dorn AV, Cooney RE, Sabin ML. COVID-19 exacerbating inequalities in the US. *Lancet*. 2020;395(10232):1243-1244. doi: 10.1016/S0140-6736(20)30893-X
3. Runkle J, Kunkel K, Easterling D, Stewart B, Champion S, Stevens L, Frankson R, Sweet W. Rhode Island state climate summary. NOAA Technical Report. 2017;NESDIS 149-RI:4.
4. National Center for Environmental health. CDC's climate-read states & cities initiative. US Centers for Disease Control and Prevention. Web. 2020. https://www.cdc.gov/climateandhealth/climate_ready.htm.
5. Climate & Health Program. Heat & health tracker. US Centers for Disease Control and Prevention. Web. 2021. <https://ephtracking.cdc.gov/Applications/heatTracker/>.
6. National Environmental Public Health Tracking Network. Data explorer. US Centers for Disease Control and Prevention. Web. <https://ephtracking.cdc.gov/DataExplorer/>.
7. National Weather Service Boston/Norton, MA. The Providence RI climate summary for the year of 2020. Climatological Report (Annual). 2021. Web. <https://forecast.weather.gov/product.php?site=BOX&product=CLA&issuedby=PVD>
8. Dahl K, Spanger-Sieffried E, Licker R, Caldas A, Abatzoglou J, Mailloux N, Cleetus R, Udvardy S, Deplet-Barreto J, Worth P. Killer heat in the United States: Climate choices and the future of dangerously hot days. Union of Concerned Scientists. 2019. <https://www.ucsusa.org/resources/killer-heat-united-states-0>.
9. Halaharvi H, Schramm PJ, Vaidyanathan A. Heat exposure and cardiovascular health: A summary for health departments. US Centers for Disease Control and Prevention. 2020.
10. Kingsley SL, Eliot MN, Gold J, Vanderslice RR, Wellenius GA. Current and projected heat-related morbidity and mortality in Rhode Island. *Environ Health Perspect*. 2016;124(4):460-7.
11. Wilhelmi OV, Howe PD, Hayden MH, O'Lenick CR. Compounding hazards and intersecting vulnerabilities: experiences and responses to extreme heat during COVID-19. *Environ. Res. Lett*. 2021;16(8): 084060. <https://iopscience.iop.org/article/10.1088/1748-9326/ac1760/meta>
12. Hoffman JS, Shandas V, Pendleton N. The effects of historical housing policies on resident exposure to intra-urban heat: A study of 108 US urban areas. *Climate*. 2020; 8(1):12. <https://doi.org/10.3390/cli8010012>.
13. Heat Watch. Providence, East Providence, Pawtucket & Central Falls, Rhode Island: Heat watch report. CAPA Strategies. Web. 2020;1:26.
14. Kawachi I, Berkman L. Social cohesion, social capital, and health. *Social Epidemiology*. 2000;17(7):290-319.
15. Lin N. Building a network theory of social capital. *Social Capital*. 1999; 22(1):28-51.
16. Health Equity Zones. Health equity zones. Rhode Island Department of Health. Web. 2018:4 pp. <https://health.ri.gov/publications/brochures/HealthEquityZones.pdf>.
17. Health Equity Zones Resilience Project. Community resilience. Rhode Island Department of Health. Web. 2021. https://health.ri.gov/programs/detail.php?pgm_id=1105.

Acknowledgment

The Rhode Island Climate Change and Health Program is supported by funds made available from the Centers for Disease Control and Prevention, Office for State, Tribal, Local and Territorial Support, under Grant #NUE1EH001317.

Disclosure

The content, proposals, and opinions within this document are those of the authors and do not necessarily represent the official position of or endorsement by the Centers for Disease Control and Prevention.

Authors

Rachel Calabro, MS, Climate Change and Health Program Manager, Rhode Island Department of Health.

Caroline Hoffman, MPH, Senior Public Health Promotion Specialist, Rhode Island Department of Health.

Correspondence

rachel.calabro@health.ri.gov

Climate Change and Health in New England: A Review of Training and Policy Initiatives at Health Education Institutions and Professional Societies

CALEB DRESSER, MD, MPH; EMILY GENTILE, BS; RACHAEL LYONS, BS; KALI SULLIVAN, BS; SATCHIT BALSARI, MD, MPH

ABSTRACT

BACKGROUND: There has been increasing interest in climate change among healthcare professionals, but it is unclear to what extent resources on this topic are available to students and clinicians in New England.

METHODS: Structured review of publicly available information regarding climate change and health activity at schools of medicine, public health, and physician assistant studies and in state medical and physician assistant societies in New England.

RESULTS: Of 39 programs reviewed, 18 (46%) had at least one climate-related initiative. Six universities accounted for 87% of climate change and health initiatives in the region. Three out of 12 state professional associations had committees or position statements addressing climate change.

CONCLUSION: There is substantial activity related to climate change and health in New England, but it is concentrated in a small number of locations. Opportunities exist to improve access to education on this topic and increase involvement of health professional associations.

KEYWORDS: climate change, health education, organized medicine, physician assistant, public health

There is increasing interest in curricula, courses, and training programs on climate change and health education.^{10,11,12,13} Medical Students for a Sustainable Future, a student-driven organization founded in 2019, advocates for action and education on climate change and health; in the two years since its creation, it has expanded to 105 medical schools, including chapters in five of the six New England states.¹⁴ New England now has free online climate and health courses,^{15,16} a physician fellowship in Climate and Human Health,¹⁷ and a steady stream of lectures and symposia on the topic.^{18,19}

However, many health care workers remain unaware of how healthcare systems contribute to climate change and how climate change threatens their patients and healthcare institutions, or have difficulty engaging with this issue.^{20,21} While it is clear that study of the intersection of climate change and health is increasingly prioritized at major academic centers in the region, it is less clear how available resources on this topic are to the bulk of students and practicing clinicians in the region.

The purpose of this review was to assess the extent of climate and health activities including educational offerings and policy statements at medical, physician assistant, and public health education institutions and within health professional societies in the New England region.

INTRODUCTION

Climate change is a threat to health in New England. Climate-related hazards include heat, extreme rainfall, flooding, vector-borne disease, hurricanes, and sea level rise.¹ Rising awareness of climate change has been accompanied by increased recognition of its implications for patients and healthcare institutions.²

The past decade has seen increasing engagement with climate change issues in the regional healthcare community. Multiple academic institutions host centers working at the intersection of human health, healthcare, and climate change.^{3,4,5} There have also been efforts to address healthcare-related greenhouse gas emissions^{6,7} and medical waste,⁸ improve resilience to climate health hazards,⁹ and communicate climate change and health issues to clinicians, policymakers and the general public.^{2,3}

METHODS

This was a structured review of publicly available material describing educational offerings, academic centers, student organizations or committees, and position statements from educational institutions offering graduate degrees in medicine, public health, and physician assistant studies, state medical societies, and state physician assistant societies in New England. All institutions granting graduate degrees in Medicine, Public Health, or Physician Assistant Studies within the six New England states (Maine, New Hampshire, Vermont, Massachusetts, Connecticut, Rhode Island) were included. State medical associations and physician assistant associations for each of the six New England states were also included. Lists of institutions were obtained from the Association of American Medical Colleges (AAMC), the American Association of Colleges of Osteopathic Medicine (AACOM), the Accreditation Review Commission on Education for the Physician Assistant (ARC-PA), the Physician

Assistant Education Association (PAEA), and the Council on Education for Public Health (CEPH).

Due to the wide variety of website designs and online platforms, ascertainment of the presence of publicly available information on items of interest could not be performed reliably via institutional website navigation. Instead, a methodology employing standardized search terms was implemented via internet search engine. Search terms consisted of “[Name of Institution]” + “climate” + “class” OR “course” OR “elective” OR “center” OR “curriculum” OR “statement” OR “student organization” OR “student group” for educational institutions, and “[Name of Association]” + “climate” + “committee” OR “position” OR “statement” OR “policy” for state associations. The first 30 results were reviewed for relevance to these climate and health activity areas. For educational institutions, activity areas included the following categories: an elective course in climate change and health, the inclusion of climate change topics in the core curriculum, an institutionally-recognized student organization focused on climate change, a center focused on climate change and health, and the existence of an institutional position statement on climate change and health. For associations, activity areas included the existence of a committee or sub-committee focused on climate change and health and the existence of a formal position statement or other policy statement regarding climate change.

Results were tabulated, and summary statistics regarding specific areas of activity, institutional subsets, and geographic regions were computed. All searches were performed in the standard public Google search engine, and reflect the top results displayed via Google search engine prioritization as of July 2021.²² Analysis was performed in R v3.1.0.²³ All materials reviewed in this study were publicly available and were accessed remotely via public-facing websites. This study was exempt from IRB review as it did not involve human subjects research.

RESULTS

A total of 51 separate institutions or associations were included in the analysis. These consisted of 11 institutions offering degrees in medicine, 13 institutions offering graduate degrees in public health, and 15 institutions offering degrees in physician assistant studies within the six New England states, as well as 6 state medical associations and 6 state physician assistant associations (Table 1).

Of the 39 degree granting institutions, 18 (46%) had at least one climate change and health initiative, 9 (23%) had two or more, and 21 (54%) had none. Institutions with education or advocacy activities related to climate change and human health were located in Massachusetts (n=8), Connecticut (n=5), New Hampshire (n=3) and Rhode Island (n=2), principally in coastal cities (Figure 1).

Table 1. Climate and health activities at educational programs in New England

	Medical Schools	Physician Assistant Programs	Public Health Programs
Elective	2	0	4
Position Statement	0	1	1
Core Curriculum	1	0	4
Student Group	7	1	4
Center	2	2	2
None of the Above	3	12	6
Total Programs	11	15	13

Figure 1. Climate Change and Health Initiatives in New England




Figure 2. Climate Change and Health Initiatives at Educational Institutions




Table 2. Climate and health initiatives at educational programs in New England

Institution	Degree Program	Climate Change and Health Initiatives
Boston University School of Medicine	Medicine	The Climate Action Group, a student, staff, and faculty organization engaged in “environmental advocacy through a variety of climate initiatives including climate change education for health professionals, improving campus sustainability and reducing the University’s use of nonrenewable energy, and exploring the role of climate change in impacting human health and wellbeing.”
Frank H. Netter MD School of Medicine at Quinnipiac University	Medicine	Human Health and Climate Change medical student group.
Geisel School of Medicine at Dartmouth	Medicine	Chapter of Medical Student for a Sustainable Future whose goal is to “integrate climate change competencies into the medical education curriculum.”
Harvard Medical School	Medicine	In addition to an active student group, an academic center focused on climate change and health activities hosts workshops and training programs, a physician fellowship that includes research and public communication, and courses on climate change and health. Additional climate and health initiatives at the university level and at individual teaching hospitals exist but did not meet inclusion criteria. A course on climate change and health is known to the authors but was not identified via structured search methodology.
The Warren Alpert Medical School of Brown University	Medicine	Students can take an elective on Climate Change and Health that provides “an overview of the wide-ranging health impacts of climate change as well as the impact of healthcare on the environment.”
UMass Chan Medical School	Medicine	A chapter of Medical Students for a Sustainable Future is active. Students created a Climate Change and Medicine elective. The core curriculum is being updated to include climate change topics.
University of Connecticut School of Medicine	Medicine	Active student group; Medical Students for a Sustainable Future has been active on sustainability at UConn Health organization meetings
Yale School of Medicine	Medicine	There is an active student group, and students have access to an academic center on climate change and health that houses a wide range of initiatives including courses, a student associate program, and a certificate program in climate and health.
Boston University	Physician Assistant	The Climate Action Group, a student, staff, and faculty organization engaged in “environmental advocacy through a variety of climate initiatives including climate change education for health professionals, improving campus sustainability and reducing the University’s use of nonrenewable energy, and exploring the role of climate change in impacting human health and wellbeing.”
MGH Institute of Health Professions	Physician Assistant	Students have access to a Center for Climate Change, Climate Justice, and Health; the core curriculum may include climate change in future years. Official Institute materials describe climate change as “a threat to the health of individuals, families, communities, and populations worldwide.”
Yale School of Medicine	Physician Assistant	Students have access to an academic center on climate change and health that houses a wide range of initiatives including courses, a student associate program, and a certificate program in climate and health.
Boston University School of Public Health	Public Health	In addition to student involvement in the Climate Action Group (above), the BU School of Public Health offers an MS in Climate and Health, a Program on Climate and Health, a climate change elective course, and includes climate change topics in the core curriculum.
Brown University School of Public Health	Public Health	Students have the opportunity to take an elective titled “Climate Change and Human Health”, which is also a substantial content area in other courses and faculty research. The School of Public Health urges “policy action to mitigate the negative health impacts of climate change.”
Dartmouth Geisel School of Medicine MPH Program	Public Health	A course on “Public Health Impacts of Climate Change” is included in the second-year curriculum and student organization is active on climate change issues.
Harvard T.H. Chan School of Public Health	Public Health	An academic center focused on climate change and health activities hosts workshops and training programs, a physician fellowship that include research and public communication, and courses on climate change and health. There is also organized student activity. Additional climate and health initiatives at the university level did not meet inclusion criteria. A course on climate change and health is known to the authors but was not identified via structured search methodology.
Simmons University Public Health Program	Public Health	Core curriculum includes climate change topics as part of an Environmental Health & Justice course.
University of New Hampshire	Public Health	Curriculum includes a Climate Change and Health course that teaches “an environmental epidemiology framework for analyzing the direct and indirect impacts of climate variability to public health as well as appropriate public policies.”
Yale School of Public Health	Public Health	A degree concentration in climate change and health is offered. There is an active student group, and students have access to an academic center on climate change and health that houses a wide range of initiatives including courses, a student associate program, and a certificate program in climate and health.

Table 3. Number of climate and health activities aggregated by academic affiliation of educational institutions.

Institutional Academic Affiliation	Total Climate and Health Activities
Yale University	6
Harvard University	6
Boston University	5
Dartmouth	4
University of Massachusetts	3
Brown University	3
All Others	4

A total of 31 climate change and health initiatives were identified, of which 12 (39%) were student groups, 6 (19%) were electives, 6 (19%) were climate change and health centers, 5 (16%) involved inclusion of climate change in the core curriculum, and 2 (6%) were position statements (Figure 2). These initiatives are summarized in Table 2. A total of 27 (87%) of the climate change and health initiatives were located at sites affiliated with one of six large research universities (Harvard University, Yale University, Boston University, Dartmouth, the University of Massachusetts, and Brown University), each of which hosted at least three initiatives; the remaining 4 initiatives were located at Simmons University, the University of New Hampshire, Quinnipiac, and the University of Connecticut (Table 3).

Climate change and health initiatives varied substantially across the programs reviewed. At several sites, the sole on-campus offering was a chapter of Medical Students for a Sustainable Future¹⁴ or another student organization; these were typically set up by students with limited institutional support and were focused on advocacy including requests for inclusion of climate change topics in the core or elective curriculum. In some cases, these efforts were successful; for example, at the UMass Chan Medical School, students recruited external lecturers to create an elective in Climate Change and Medicine,²⁴ and efforts to include climate change in the core curriculum are ongoing.²⁵ Academic centers dedicated to climate change and human health tended to offer a range of activities that included research programs, CME courses, elective courses, fellowships in Climate and Human Health for physicians, and a range of climate change and health policy and advocacy material.^{3-5,15-17,19}

Of the 12 state-level medical and physician assistant associations reviewed, three medical associations (Maine, Massachusetts, and Vermont) had position statements regarding climate change. Two statements included language on the implications of climate change for the health of patient populations, one called for reductions in greenhouse gas emissions, and one called for investment in climate change adaptation as a means to protect health. All three position

statements were issued in 2017 or later. Two state medical associations (Maine and Massachusetts) had committees whose goals included addressing climate change issues.

DISCUSSION

This review describes climate change and health activity in a set of health education institutions and professional societies in New England. The burgeoning number of educational offerings, their uneven distribution, and the lack of public action from most regional medical and physician assistant societies are of significance to educators, students, advocates, and policymakers.

Climate change and health has clearly gained attention as an educational topic and object of organized student and professional activity. While this subject was largely absent from discussion outside specialist circles for many years, nearly half of the educational institutions assessed in this review now host some form of climate change and health education or advocacy activity, and the past four years have witnessed the first statements on climate change from medical societies in the region.

However, most climate change and health activity identified in this review was concentrated at large research universities; the majority of medical, public health, and physician assistant education institutions in the region do not publicly describe any climate-related activities. The concentration of initiatives at coastal, urban institutions may impede clinicians and students in rural or interior locations from accessing these resources and presents an important opportunity for growth. Climate change will have substantial effects on health throughout New England during the decades in which clinicians who are being trained today can expect to practice,¹ and it is important that opportunities for climate change education and action reach clinicians training and practicing in communities throughout the region.

Students, faculty, and administrators interested in implementing climate change and health curricula or other offerings at their institutions now have access to a wide variety of materials to support their efforts. Organizations including Medical Students for a Sustainable Future,²⁶ Healthcare Without Harm,²⁷ several academic centers,^{3,4} and the Medical Society Consortium on Climate and Health²⁸ offer model curricula, educational content, templates for action, networking opportunities, remote courses, and other resources. Recent academic work provides needs assessments, guidelines on program design, and case studies of climate change and health education and advocacy in action.^{2,11-13,29-32} For residents, fellows, and practicing clinicians, resources ranging from career planning tools³³ to federal guidance on the specific health impacts of climate change are also available.^{1,34}

There are abundant opportunities for professional societies to increase their engagement; only three of the twelve professional societies reviewed were active on climate change

issues. Policy statements can provide an authoritative voice in support of climate and health education, climate smart healthcare systems, and climate resilient communities. Affiliation with national organizations such as the Medical Society Consortium on Climate and Health is another avenue for action – one that the medical societies in Vermont, Rhode Island, and Massachusetts have already taken.³⁵

While uneven, the overall trajectory of climate change and health activity in New England is one of rapid expansion and improving capabilities. In the brief period since completion of data collection for this review, a new regional fellowship program focused on climate and health advocacy has become available.³⁶ In addition, the climate and health activity of nursing programs, hospitals, and other organizations that were not reviewed in this study is substantial; examples include the Nurse's Climate Challenge,³⁷ hospital programs,³⁸ state government programs,^{39,40} the launch of the *Journal of Climate Change and Health*⁴¹, the inclusion of the climate crisis as key topic area in the *New England Journal of Medicine*,⁴² and special issues of regional publications focused on climate change.^{43,44} While many of these initiatives are associated with large research universities that are already active in this area, they provide a model for future engagement with this issue throughout New England.

Students, trainees, and healthcare professionals now have access to an increasing variety of education and policy materials related to climate change and health. As New Englanders face escalating health risks from hurricanes, extreme rainfall, heat waves, vector-borne disease, sea level rise, and other hazards, it is important that these education and policy resources be extended to benefit health professionals and patients throughout the region.

LIMITATIONS

As a review of publicly available material rather than a survey of educators and administrators, our methodology assesses information that is accessible to the general public and intentionally mimics the approach students, prospective students, and healthcare workers may take when seeking information on offerings at their institutions, but also has several limitations. All information was gathered from public internet sources, and as such may have missed initiatives that were not described on the internet or were behind firewalls, password protection, or other barriers. Some relevant programs may not have been prioritized by search engines using our search criteria; the term “environment” was not used in structured searches as it led to a large number of results unrelated to climate change. As searches were restricted to health education institutions, some potentially relevant initiatives did not meet inclusion criteria. As a result, initiatives including an elective course on climate change and health,¹⁵ a chapter of Medical Students for a Sustainable Future,¹⁴ and a university-level center working on

environmental issues including health impacts of climate change⁴⁵ were not identified by structured searches or included in our analysis.

CONCLUSION

Climate change threatens patients, healthcare systems, and future clinicians. This review reveals uneven access to education and policy guidance on the subject of climate change and health; while some academic centers have many climate-related initiatives, the majority of the institutions reviewed do not publicly describe any offerings at all, and only a quarter of the medical and physician assistant associations reviewed are publicly engaged with this issue. Health professionals, administrators, and students have an opportunity to make education on this topic more accessible and to advocate for climate change action through state medical and physician assistant associations.

References

1. USGCRP, 2018: Impacts, Risks, and Adaptation in the United States: Fourth National Climate Assessment, Volume II [Reidmiller, D.R., C.W. Avery, D.R. Easterling, K.E. Kunkel, K.L.M. Lewis, T.K. Maycock, and B.C. Stewart (eds.)]. U.S. Global Change Research Program, Washington, DC, USA, 1515 pp. doi: 10.7930/NCA4.2018.
2. Salas RN, Friend TH, Bernstein A, Jha AK. Adding A Climate Lens To Health Policy In The United States. *Health Aff (Millwood)*. 2020 Dec;39(12):2063-2070. PMID: 33284694.
3. C-CHANGE Center for Climate, Health and the Global Environment. Harvard T.H. Chan School of Public Health. 2021. Accessed September 10, 2021. URL: <https://www.hsph.harvard.edu/c-change/>
4. Yale Center on Climate Change and Health. Yale School of Public Health. 2021. Accessed September 10, 2021. URL: <https://ysph.yale.edu/climate/>
5. Program on Climate and Health. Boston University School of Public Health. 2021. Accessed September 10, 2021. URL: <https://sites.bu.edu/climateandhealth/>
6. Boston Hospitals Cut Energy Use by 6% Despite Continued Facility Growth. Healthcare Without Harm. December 11, 2014. Accessed September 10, 2021. URL: <https://noharm-uscanada.org/articles/press-release/us-canada/boston-hospitals-cut-energy-use-6-despite-continued-facility-growth>
7. Sherman J, Le C, Lamers V, Eckelman M. Life cycle greenhouse gas emissions of anesthetic drugs. *Anesth Analg*. 2012 May;114(5):1086-90. doi: 10.1213/ANE.0b013e31824f6940. Epub 2012 Apr 4. PMID: 22492186.
8. Hsu S, Thiel CL, Mello MJ, Slutzman JE. Dumpster Diving in the Emergency Department. *West J Emerg Med*. 2020 Aug 24;21(5):1211-1217. doi: 10.5811/westjem.2020.6.47900. PMID: 32970577; PMCID: PMC7514403.
9. Municipal Vulnerability Preparedness Program. Commonwealth of Massachusetts. 2021. Accessed September 10, 2021. URL: <https://www.mass.gov/municipal-vulnerability-preparedness-mvp-program>
10. Howard, B. Climate change in the Curriculum. AAMC News. October 10, 2019. URL: <https://www.aamc.org/news-insights/climate-change-curriculum>
11. Ryan EC, Dubrow R, Sherman JD. Medical, nursing, and physician assistant student knowledge and attitudes toward climate change, pollution, and resource conservation in health care. *BMC Med Educ*. 2020;20(1):200. Published 2020 Jun 23. doi:10.1186/s12909-020-02099-0
12. Kotcher J, Maibach E, Miller J, Campbell E, Alqodmani L, Maiero M, Wynn A. Views of health professionals on climate change and health: a multinational survey study. *Lancet Planet Health*.

- 2021 May;5(5):e316-e323. doi: 10.1016/S2542-5196(21)00053-X. Epub 2021 Apr 8. PMID: 33838130; PMCID: PMC8099728.
13. Lemery J, Balbus J, Sorensen C, Rublee C, Dresser C, Balsari S, Calvello Hynes E. Training Clinical And Public Health Leaders In Climate And Health. *Health Aff (Millwood)*. 2020 12; 39(12):2189-2196. PMID: 33284695.
 14. Medical Students for a Sustainable Future: Partnerships. 2021. Accessed September 10, 2021. URL: <https://ms4sf.org/partnerships/>
 15. The Health Effects of Climate Change. Harvard University. 2021. Accessed September 10, 2021. URL: <https://online-learning.harvard.edu/course/health-effects-climate-change?delta=0>
 16. Climate Change and Health Certificate. Yale School of Public Health. 2021. Accessed September 10, 2021. URL: <https://ysph.yale.edu/cchcert/>
 17. Climate and Human Health Fellowship. Beth Israel Deaconess Medical Center. 2021. Accessed September 10, 2021. URL: <https://www.bidmc.org/medical-education/medical-education-by-department/emergency-medicine/emergency-medicine-fellowship-programs/climate-and-human-health-fellowship>
 18. Climate Change & Health Symposium 2021: The Intersection of Climate Change, COVID-19, and Structural Racism. MGH Institute of Health Professions. 2021. Accessed September 10, 2021. URL: <https://www.mghihp.edu/climate2021>
 19. The Climate Crisis and Clinical Practice Symposium. Harvard T.H. Chan School of Public Health. 2019. Accessed September 10, 2021. URL: <https://www.hsph.harvard.edu/c-change/events/the-climate-crisis-and-clinical-practice-symposium/>
 20. Hathaway J, Maibach EW. Health implications of climate change: a review of the literature about the perception of the public and health professionals. *Curr Environ Health Rep*. 2018; 5: 197-204
 21. Hubbert B, Ahmed M, Kotcher J, Maibach E, Sarfaty M. Recruiting health professionals as sustainability advocates. *Lancet Planet Health*. 2020; 4: e445-e446
 22. How results are automatically generated. Google. 2021. Accessed September 10, 2021. URL: <https://www.google.com/search/howsearchworks/how-search-works/ranking-results/>
 23. R Core Team (2019). R: A language and environment for statistical computing. R Foundation for Statistical Computing, Vienna, Austria. URL <https://www.R-project.org/>.
 24. Medical Students Organize Climate Elective. UMass Chan Medical School. 2021. Accessed September 12, 2021. URL: <https://www.umassmed.edu/growinggreen/newsletter/student-climate-elective/>
 25. Climate Change Curriculum. UMass Chan Medical School. 2021. Accessed September 12, 2021. URL: <https://www.umassmed.edu/oume/initiatives/climate-change-curriculum/>
 26. Climate and Health Curriculum Reform Guide. Medical Students for a Sustainable Future. 2021. Accessed September 13, 2021. URL: <https://ms4sf.org/climate-and-health-curriculum-reform-guide/>
 27. Physician Network Resource Library. Healthcare Without Harm. 2021. Accessed September 13, 2021. URL: <https://sites.google.com/hcwh.org/physiciansnetwork/home>
 28. GCCHE Knowledge Bank. The Medical Society Consortium on Climate and Health. 2021. Accessed September 13, 2021. URL: <https://medsocietiesforclimatehealth.org/educate/health-educators/>
 29. Lemery J, Sorensen C, Balbus J, Newman L, Davis C, Reno E, Salas R, Hynes EC. Science Policy Training for a New Physician Leader: Description and Framework of a Novel Climate and Health Science Policy Fellowship. *AEM Educ Train*. 2019 Apr 1;3(3):233-242. doi: 10.1002/aet2.10323. PMID: 31360816; PMCID: PMC6637012.
 30. Limaye VS, Grabow ML, Stull VJ, Patz JA. Developing A Definition Of Climate And Health Literacy. *Health Aff (Millwood)*. 2020 Dec;39(12):2182-2188. PMID: 33284692; PMCID: PMC8428792.
 31. Cook C, Demorest S, Schenk E. Nurses and Climate Action. *AJN, American Journal of Nursing*: April 2019 - Volume 119 - Issue 4 - p 54-60 doi: 10.1097/01.NAJ.0000554551.46769.49
 32. Neal-Boylan L, Breakey S, Nicholas PK. Integrating Climate Change Topics Into Nursing Curricula. *J Nurs Educ*. 2019 Jun 1;58(6):364-368. doi: 10.3928/01484834-20190521-09. PMID: 31157907.
 33. Rublee C, Dresser C, Linstadt H, Dahl R, Moretti K. RAMS Roadmaps: Climate Change. Society for Academic Emergency Medicine. 2021. Accessed September 13, 2021. <https://www.saem.org/detail-pages/publication/rams-roadmaps-climate-change>
 34. Climate and Health. United States Centers for Disease Control and Prevention. 2021. Accessed September 13, 2021. <https://www.cdc.gov/climateandhealth/default.htm>
 35. Affiliates. The Medical Society Consortium on Climate & Health. 2021. Accessed September 10, 2021. URL: <https://med-societiesforclimatehealth.org/about/affiliates/>
 36. Climate Health Organizing Fellows Program. Cambridge Health Alliance Center for Health Equity Education and Advocacy. 2021. Accessed September 10, 2021. URL: <https://www.healthequity.challiance.org/climate-health>
 37. Nurses Climate Challenge. 2021. Accessed September 10, 2021. URL: <https://nursesclimatechallenge.org/>
 38. Center for the Environment and Health. Massachusetts General Hospital. 2021. Accessed September 10, 2021. URL: <https://www.massgeneral.org/environment-and-health>
 39. Climate and Health. Commonwealth of Massachusetts. 2021. Accessed September 10, 2021. URL: <https://www.mass.gov/climate-and-health>
 40. Climate Change and Health Program. State of Rhode Island. 2021. Accessed September 10, 2021. URL: https://health.ri.gov/programs/detail.php?pgm_id=174
 41. The Journal of Climate Change and Health. Elsevier. 2021. Accessed September 10, 2021. URL: <https://www.journals.elsevier.com/the-journal-of-climate-change-and-health>
 42. Climate Crisis and Health. The New England Journal of Medicine. 2021. Accessed September 13, 2021. URL: https://www.nejm.org/climate-crisis?query=main_nav_lg
 43. Climate Change and it's Impact on Public Health. *Worcester Medicine*, 72:2. May/June 2011. Accessed September 13, 2021. https://www.wdms.org/wp-content/uploads/2020/08/womed_may-jun11.pdf
 44. Climate Change. *Worcester Medicine*, 90:2. March/April 2021. Accessed September 13, 2021. URL: https://issuu.com/wdms/docs/wdms_worcestermedicine_marapr_v09digitaledition
 45. Human Health Programs. Harvard University Center for the Environment. 2021. Accessed September 14, 2021. URL: <https://environment.harvard.edu/human-health>

Acknowledgment

This project was supported by the CrisisReady initiative at Harvard and Direct Relief. Dr. Caleb Dresser's work was also supported by the Climate and Health Foundation.

Authors

Caleb Dresser, MD, MPH, Instructor of Emergency Medicine at Harvard Medical School.
 Emily Gentile, BS, MD candidate at University of Massachusetts Chan Medical School.
 Rachael Lyons, BS, MD candidate at University of Massachusetts Chan Medical School.
 Kali Sullivan, BS, MD candidate at Tufts University School of Medicine.
 Satchit Balsari, MD, MPH, Assistant Professor of Emergency Medicine at Harvard Medical School and Harvard T.H. Chan School of Public Health.

Disclosures

The authors have no relevant financial disclosures.

Disclaimer

The views expressed herein are those of the authors and do not necessarily reflect the views of the institutions with which they are affiliated.

Correspondence

Caleb Dresser, MD, MPH
 Department of Emergency Medicine
 1 Deaconess Road, Boston, MA 02215
cdresser@bidmc.harvard.edu

Beyond the Hazard Vulnerability Analysis: Preparing Health Systems for Climate Change

JOSHUA BAUGH, MD, MPP, MHCM; KATIE KEMEN, MBA; JOHN MESSERVY, AIA; PAUL BIDDINGER, MD

ABSTRACT

INTRODUCTION: Climate change is heightening both long-term adverse risks to human health and the immediate-term risk of injuries and illness following climate-related disaster events that are becoming more frequent and severe. In addition to its direct health effects, climate change poses new threats to the nation's health care infrastructure – with potential to negatively impact healthcare capacity amidst increasing demand – through risks of flooding, wind damage, heat stress, power outages, and other physical harm to facilities. The typical Hazard Vulnerability Analyses conducted annually by hospitals use historical data to assess risks; these analyses are likely now inadequate for future preparation due to the impact of climate change. This article describes one approach to how healthcare leaders can better assess both near-term and long-term risks due to climate change, to mitigate against unprecedented but foreseeable threats.

METHODS: In our large health system in the US Northeast, a process was undertaken to gather updated data and expert projections to forecast threats faced by each of our facilities in different climate-related disaster scenarios. Hazards examined in our setting included precipitation-based and coastal flooding events, heat waves, and high wind events, in addition to seismic events. Probabilities of occurrence and extents of different hazards were projected for the near term (2030) and the long term (2070). We then performed detailed vulnerability analyses for each facility with the predicted amount of rainfall, storm surge, heat stress, and windspeed, in collaboration with leaders at each facility. This was followed by a process to understand what would be needed to mitigate each vulnerability along with the associated costs. Ultimately, a cost/benefit analysis was performed – incorporating the relative likelihood and impact of different scenarios – to decide which improvement projects to embark on immediately, and what to defer and/or incorporate into future building plans.

RESULTS: In our system, all facilities were vulnerable to the effects of increased temperatures, and multiple hospitals were noted to be vulnerable to extreme precipitation, storm surge, and high winds. Specific damaging scenarios identified included flooding of basements and building in-

frastructure spaces, water entry through windows during high winds, and overheating of power systems during heat waves. Potential solutions included improved power redundancy for cooling systems, enhancements to roof and window systems, and the acquisition of deployable flood barriers. We identified four categories for prioritization of action based on projected impact: 1) priorities in need of urgent mitigation, 2) priorities in need of investigative study for medium-term mitigation, 3) priorities for planned capital improvement projects, and 4) priorities to integrate into new facility construction.

DISCUSSION: While the specific risks and vulnerabilities for each facility will differ according to its location and structural features, the approach we describe is broadly applicable. By forecasting specific risks, diagnosing vulnerabilities, developing potential solutions, and using a risk/benefit approach to decision making, hospitals can work toward protecting facilities and patients in the face of potential climate related natural disasters in an economically sound manner.

KEYWORDS: climate change, climate resilience, disaster medicine, hazard analysis, emergency preparedness

INTRODUCTION

Due to climate change, the frequency and intensity of dangerous weather patterns and natural disasters are known to be increasing throughout the world. These trends pose significant threats to human health both in the immediate and long term. The impact of climate-related changes on patient health include both direct injury and illness from discrete events such as hurricanes and fires, as well as exacerbation of chronic conditions like asthma and heart disease from increased levels of atmospheric carbon and higher average temperatures.¹ In addition to the increased demand on healthcare services that these trends will bring, there is another particularly difficult challenge for hospitals and other care facilities: at precisely the moment a weather event makes more patients ill or injured, that event may also directly interfere with the ability of healthcare facilities to function effectively.

This is, of course, the focus of most hospital-based

emergency preparedness initiatives and the rationale behind standard Hazard Vulnerability Analyses (HVA). The problem is, in the era of climate change, HVAs based on historical data often underestimate the true level of risk for health-care facilities. For example, hospitals in New York had, in fact, built flood barriers as part of their preparedness plans for extreme precipitation prior to 2012; yet, the flood depths produced by Hurricane Sandy were higher than those planned for and unfortunately overwhelmed these physical barriers, necessitating the evacuation of multiple facilities and resulting in extensive damage.² Similarly, multiple hospitals in Louisiana very recently required evacuation due to damage from high winds and precipitation during Hurricane Ida, a particularly challenging task amidst the power grid failures resulting from the storms.³ These types of events are harmful to patient care, harrowing for staff, and ultimately very expensive for the facilities involved.

The unprecedented weather patterns created by climate change require a new approach to emergency preparedness for healthcare systems, as risks previously thought impossible are likely to become reality in coming years. Yet, each healthcare entity cannot reasonably prepare for every possible extreme scenario; the cost would be overwhelming and would compromise other important hospital missions. An approach is therefore needed that accounts for the rapid pace of climate change and prepares facilities appropriately without bankrupting health systems. In this article, we describe the strategy that our large health system in the US Northeast undertook to increase the climate change resilience of our facilities.

METHODS

Setting

Our large integrated health system is located in New England in the United States, and includes facilities in urban, suburban, rural, and island locations. There are 10 individual hospitals in our system along with many other associated facilities, including rehabilitation centers, administrative buildings, and research complexes. Some of the hospitals are high volume quaternary care academic centers, while others are relatively small community hospitals with limited numbers of service lines. In total, our process included resilience analyses for over 30 individual buildings across the system.

Climate Projections

Our first step involved an assessment of climate change projections for each of our locations over short-term and longer-term time horizons (for the years 2030 and 2070, respectively). To do this, we engaged a consultant group with climate change expertise and leveraged recent city and state level reports on projected changes in weather patterns over time. For our locations, we focused on three climate-related threats: 1) Extreme heat (higher temperatures and prolonged

duration) 2) flooding (due to storm surge and/or high precipitation), and 3) high winds (likely in the setting of severe storms). We also decided to include a seismic assessment, although this is not known to be climate related. Fire and drought were not included as likely threats in our location, but these would certainly be necessary to plan for in other parts of the US.

For heat projections, we used a report commissioned by the city of Cambridge, Massachusetts with extrapolations for our various locations. To assess flood risk, we created probabilistic and consequence-based flood maps for each location. Probabilistic flood maps showed us flood extents and depths for 100-year and 500-year storms at our two future time horizons. Probabilistic maps were useful in assessing risk tolerance; for example, we might tolerate flooding that extended into a parking lot but not a building or we might tolerate a 5% annual chance of flooding in a garage but not a 1% annual chance of flooding in a clinical building. Consequence-based flood maps modeled worst-case scenario storms, which provided insight into the maximum level of protection our health system might need, or if that wasn't feasible, what kind of emergency preparedness plans we needed. This modeling was possible through use of the Boston-Harbor Flood Risk Model, commissioned by the Massachusetts Department of Transportation.⁴

Previous studies predict that wind speeds are expected to increase 5% for each 1 degree increase in sea surface temperature.⁵ Applying this principle to climatology and storm surge models, high level estimates of potential wind speed increases through 2070 were prepared. These projections were compared with the wind speed design criteria specified by the Massachusetts Building Code for the location and use at the time of construction.

To evaluate seismic risk, we used historical building codes for each of the facilities evaluated, assuming that each building had been constructed with only the seismic resiliency required by local code for its primary use. The probability of earthquakes was calculated based on historic data and an understanding of the geological conditions which help to predict epicenters, drivers, and propagation of seismic events.

Ultimately, we produced a series of scenarios categorized by likelihood and projected consequence, which could subsequently be applied to each facility.

Vulnerability Analysis

Next, we sought to examine the potential impact of various climate scenarios on each of our facilities, to discover vulnerabilities to different types and levels of threat. To do this, members of our project team met with structural engineers, facility managers, research, operations, and emergency preparedness leaders, as well as clinical leaders from every site. At these meetings, these local leaders were given the projections for various levels of heat, flooding, wind speed, and

seismic disturbance, and asked to assess the potential consequences of each for their respective areas of responsibility. At each site, the following specific structural/systems features were explicitly discussed:

Main grid power, emergency power, natural gas, medical gases, fuel oil, HVAC system, potable water, non-potable water, storm water removal, waste water removal, medical waste, information technology, communications, among others.

In addition, the following operational functions were also each discussed:

Patient transfers, staff availability, on-site accommodations for staff, capacity for patient surges, medical supply chain, lab and pharmacy operations, and food and nutrition availability, among others.

Multiple conversations occurred with each set of local leaders to quantify potential disruptions to each of these functions due to the various projected climate threats. Ultimately, a comprehensive list of vulnerabilities for all sites was generated, organized by likelihood of an event and the potential consequences of that event based on our models. Consequences from a vulnerability were organized into three categories: Major, Severe, and Catastrophic (See **Table 1** for definitions of these criteria by facility type).

Table 1. Facility Damage Assessment Criteria

	Major	Severe	Catastrophic
Outpatient clinical operations	Interruption of clinical activities for 1 week or less OR inability to provide essential services	Interruption of clinical activities for 1 month or less	Interruption of clinical activities for greater than 1 month OR complete loss of an outpatient care site
Inpatient clinical operations	Evacuation or closure of some major inpatient care areas (but not the whole facility) for < 3 days	Evacuation or closure of some major inpatient care areas (but not the whole facility) for < 14 days	Evacuation of all inpatient care areas OR closure of some major inpatient care areas for > 14 days
Research operations	Evacuation or closure of major research labs/ areas (but not all research areas) for < 3 days	Evacuation or closure of major research labs/ areas (but not all research areas) for < 14 days OR loss of unique animals, cell lines, specimens, data, etc	Evacuation of all research areas OR complete loss of a research facility
Support services/ areas	Damage or destruction of a support area that will cause moderate impact to facility operations	Damage or destruction of a support area that will cause major impact to facility operations	Damage or destruction of a support area that will cause severe impact to facility operations

Developing Interventions

Once a comprehensive list of vulnerabilities was created, we then worked with a team of facility engineers along with operational and emergency preparedness leaders to develop plans for what would be needed to reduce or eliminate each vulnerability. Some solutions involved structural changes, while others involved primarily operational changes. Common interventions included moving equipment from a basement to a location higher in the building and installing cooling mechanisms for backup power systems. These projects ranged widely in complexity and estimated cost. Through this process, an exhaustive list of over 300 potential projects was generated, with a description of the work requirements and estimated costs for each.

Assessing Interventions

The list of potential interventions was far too long and expensive for every project to be carried out in the near term, so a prioritization scheme was needed. Elements of the scoring system included whether failure to fix an issue would create a regulatory violation, which type of clinical area would be affected (e.g. emergency departments and operating rooms vs. outpatient clinic spaces), and how severe the disruption would be to local operations. Weighed against the costs of each project, the team chose to prioritize interventions into four categories:

- 1) Immediate priorities for urgent mitigation
- 2) Projects in need of investigative study for medium-term mitigation
- 3) Priorities for planned capital improvement projects
- 4) Priorities for integration into new facility construction

This rational blended approach facilitated a plan that would increase preparedness in the short term while also ensuring that climate resiliency would improve over time – in line with increasing levels of threat over time – as new construction projects were undertaken across the system. Where physical plant improvements were not feasible or would be delayed for later phases of implementation, emergency preparedness and operational leaders were tasked with developing emergency plans for the relevant climate-driven events.

RESULTS

Our systematic approach to risk assessment and intervention development identified over 300 potential vulnerabilities in our health system, with a range of consequences from moderate to catastrophic. We identified that climate change adjusted projections for heat, flooding, and wind would all strain our current facilities if improvements were not undertaken. See **Table 2** for examples.

Table 2. Examples of vulnerabilities created by each type of climate-related disaster in our settings, along with potential solutions in the near and longer terms.

Threats	Examples of Vulnerabilities	Examples of Solutions
Heat Waves	<ul style="list-style-type: none"> - Lack of cooling for back-up power sources - Lack of back-up power for cooling mechanisms - Potential regional brownouts - Potential need for hospital as shelter for the community 	<ul style="list-style-type: none"> - Adding chillers to emergency power - Creating street connections for supplemental emergency power & chillers - Energy efficiency initiatives to reduce burden on power sources, in current and future facilities
Flooding	<ul style="list-style-type: none"> - Critical infrastructure interruptions (transit, energy, telecom) - Supply chain disruptions - Personal impact to workforce - Loss of critical services on lower levels (utilities, pharmacy, food services) - Water contamination 	<ul style="list-style-type: none"> - Business continuity planning - Flood barriers - Utility hardening - Storm water management systems - Advocate and partner for public utility improvements - Incorporate flood projections into new building designs/relocations
High Winds	<ul style="list-style-type: none"> - Hazards from slate and ballasted roofs - Wind-driven rain entering through windows, under cladding and roofs - Equipment blowing off roofs 	<ul style="list-style-type: none"> - Roofing repairs and replacements - Wind resiliency improvements to current facades - Stronger equipment tie-downs - New facilities built to codes of areas with higher wind speeds

Nearly all facilities faced vulnerabilities amidst projected extended heat waves, particularly during summer months when a majority of days could be expected to exceed 90-degree temperatures by 2070. The most common problem encountered was how to keep buildings cool if local power systems or the grid were to fail, as most buildings did not have emergency power for cooling. Most buildings also faced vulnerabilities if wind speeds were to increase markedly, with particular risks to older slate and ballasted roofs. In addition, some equipment on building exteriors was deemed at risk due to inadequate strength of tie-downs, and many buildings faced risk of wind-driven water entry through windows and under roofs.

Flooding was also projected to be a major risk for many sites. Flooding on-site could directly damage critical services on buildings' lower levels including utilities, pharmacy, food services, and water supply. Flooding of surrounding areas could disrupt critical infrastructure (e.g. local transit, energy lines, and telecommunications), and interrupt usual supply chain routes. Such flooding could also impede the ability of staff to come to work (and return home). Most facilities were not fully prepared for the flood depths that could occur in future climate-adjusted models.

Some of the more immediate interventions undertaken in response to these vulnerabilities included adding chillers to emergency power, creating connections to the street for accessing supplemental emergency power, replacing roofs and windows, increasing wind resiliency of existing structures, strengthening equipment tie-downs, creating higher flood barriers, improving storm water management systems, and hardening of utilities against water damage. Longer-term interventions included considering flood projections into new building site choices, incorporating wind resilience into new building plans far beyond local code requirements, and advocating for improvements to local public utility infrastructure.

DISCUSSION

In this article, we present a systematic approach to assessing healthcare system vulnerabilities and prioritizing preparedness interventions in the setting of increasingly severe weather events resulting from climate change. Utilizing climate projections that largely already existed for our local area, we were able to create realistic scenarios for what each of our facilities could expect in coming years, and plan for levels of threat that would not be accounted for by usual Hazard Vulnerability Analyses based on historical data. Some of the projections were startling and resulted in construction principles we would otherwise have never employed; one of our Massachusetts hospitals now has a building enclosure, roof, and walls designed to the building code for wind levels in Miami, for example. Without a clear-eyed assessment of climate change scenarios, our healthcare systems risk more events like those seen in Hurricanes Katrina and Sandy. However, by using data often already available, health systems can plan for these types of severe events and become better prepared for when they occur more frequently in coming decades.

A key feature of our strategy was the blending of short- and long-term projections with a mix of time horizons for improvements. Not all projects could feasibly be completed immediately, but not all needed to be; some could safely wait a decade or two based on climate projections. This allowed many strategies to be built into future construction plans, which is more cost effective than renovating current structures. Plans created now can therefore help to ensure increased climate resiliency for many decades.

One important lesson from our experience was the need to engage local leaders across disciplines at each facility. Assembling climate projections and flood maps can be done in a conference room or behind a desk, but the vulnerability analysis cannot. Without engaging the people who understand the structure and operational processes of each facility

in depth, a vulnerability analysis cannot effectively reveal the most pertinent issues to be addressed. In addition, involving local leaders – including CFOs – helps to ensure buy-in for projects once improvement possibilities are identified.

While our analysis focused only on facility structures and day-to-day operations, there are certainly broader impacts of climate change that hospitals should consider moving forward. One major challenge will involve accommodating increasing healthcare demand due to the impact of climate change on patients. Others have discussed how both discrete weather events and overall climate trends are likely to increase the burden of both acute injury and vector-borne illness as well as many chronic diseases.¹ Many American hospitals already lack reserve capacity and struggle with overcrowding on a daily basis; future health system design must account for yet further increases in the number of patients requiring care in coming decades.⁶

Another important consideration is the impact of climate change on facilities beyond traditional hospitals. In our setting, we included research facilities as medical research is core to the mission of our health system; with extra-cold freezers and other unique needs, protecting research capability can necessitate additional types of strategies. We also assessed rehabilitation centers, and future work at local and broader levels should include skilled nursing and long-term care facilities. If these types of facilities are rendered unsafe during a disaster, their residents will end up in hospitals, only further stressing the healthcare system. For example, Florida hospitals received many patients during Hurricane Irma in 2017 after cooling systems in nursing homes failed.⁷ Helping to ensure that lower level of care settings have preparedness plans can help alleviate burdens on hospitals at times of great stress.

CONCLUSION

The potential impacts of climate change on our hospitals and health systems are daunting but must be faced head-on to ensure that patients can receive life-saving care when they need it most. While the specific risks and vulnerabilities for each hospital will differ according to location and structural features, the rational blended approach we describe here is broadly applicable. By forecasting specific risks, diagnosing vulnerabilities, developing potential solutions, and using a risk/benefit approach to decision making, hospitals can work toward protecting themselves and their patients in the face of potential climate-related natural disasters in an economically sound manner.

References

1. Sorensen CJ, Salas RN, Rublee C, Hill K, Bartlett ES, Charlton P, Dyamond C, Fockele C, Harper R, Barot S, Calvello-Hynes E. Clinical implications of climate change on US emergency medicine: Challenges and opportunities. *Annals of Emergency Medicine*. 2020 Aug 1;76(2):168-78.
2. Teperman S. Hurricane Sandy and the greater New York health care system. *Journal of Trauma and Acute Care Surgery*. 2013 Jun 1;74(6):1401-10.
3. Achenbach J, Johnson A, Dupree J. Hurricane Ida forces three damaged hospitals to evacuate patients. *Washington Post*. August 30, 2021.
4. Bosma K, Douglas E, Kirshen P, MacArthur K, Miller S, Watson C. MassDOT-FHWA Pilot Project Report: Climate Change and Extreme Weather Vulnerability Assessments and Adaptation Options for the Central Artery. 2015. https://www.mass.gov/files/documents/2018/08/09/MassDOT_FHWA_Climate_Change_Vulnerability_1.pdf.
5. Young IR, Zieger S, Babanin AV. Global trends in wind speed and wave height. *Science*. 2011 Apr 22;332(6028):451-5.
6. Baugh JJ, White BA, Biddinger PD, Raja AS, Wittbold KA, Sonis JD, Yun BJ. To solve our new emergency care crisis, let's start with the old one. *The American Journal of Emergency Medicine*. 2020 Oct 1;38(10):2000-1.
7. Dosa DM, Skarha J, Peterson LJ, Jester DJ, Sakib N, Ogarek J, Thomas KS, Andel R, Hyer K. Association between exposure to Hurricane IRMA and mortality and hospitalization in Florida nursing home residents. *JAMA network open*. 2020 Oct 1; 3(10):e2019460-.

Authors

Joshua Baugh, MD, MPP, MHCM, Department of Emergency Medicine, Massachusetts General Hospital; Center for Disaster Medicine, Massachusetts General Hospital; Harvard Medical School.

Katie Kemen, MBA, BSC Group, Inc.

John Messervy, AIA, Massachusetts General Brigham.

Paul Biddinger, MD, Department of Emergency Medicine, Massachusetts General Hospital; Center for Disaster Medicine, Massachusetts General Hospital; Harvard Medical School, Mass General Brigham.

Correspondence

Joshua Baugh, MD, MPP, MHCM

jbaugh@partners.org

Providence's Vulnerability to Floods: Impacts of Sea Level Rise, Stronger Storms, and Heavier Rainfall

ANDREW E. BINDER, BA; SELIM SUNER, MD, MS; H. CURTIS SPALDING, MPA; ERICH OSTERBERG, PhD

KEYWORDS: climate change, climate resilience, disaster medicine, hazard analysis, emergency preparedness, flooding

INTRODUCTION

The City of Providence, RI, and its healthcare infrastructure face growing risks from climate change. Environmental stressors including higher temperatures and atmospheric changes increase pressure on the existing social-environmental system. In a 2019 report, the Providence Emergency Management Agency (PEMA) and Local Hazard Mitigation Committee (LHMC) identified flood-related hazards as a major risk to Providence's built and natural environments and community health.¹ Hazard Vulnerability Analyses, (HVAs), conducted annually as part of each acute care hospitals' emergency preparedness, also consider the impact of flooding, hurricanes, and winter storms in their danger assessments.

Providence is located on the lowland shore of the northern fringe of Narragansett Bay, a coastal estuary environment with a 4-foot tidal range.² The city area also includes four river systems that are tidally affected. Providence's northern region is drained by the Moshassuck River, and the western-central region is drained by the Woonasquatucket River. Their confluence at downtown's Waterplace Park forms the tidal Providence River, which flows southward to join the Seekonk River at the head of Narragansett Bay. The city's coastal floodplain from the port area along Allens Avenue and river floodplains through the downtown Fox Point Hurricane Barrier (FPHB) and up the Moshassuck and Woonasquatucket River corridors, face risk of hazards related to inland, riverine, tidal, and storm-surge flooding.²

Since the establishment of the Providence Colony in 1636, centuries of industrial and residential development have modified the coastal landscape. Dense urbanization, river engineering, and waterfront barriers replaced the region's salt marshes, swamps, and river floodplains with impervious surface cover (>37% of the city), which has decreased the city's ability to infiltrate excess water into soil.^{2,3} This development pattern has compounded the magnitude and frequency of flooding events, especially urban flooding caused by stormwater runoff.²

The causes of flooding are multifactorial. Global surface temperatures have risen over the past 150 years, with a rapid change since 1970, due to increasing atmospheric concentrations of carbon dioxide (CO₂), methane (CH₄), nitrous oxide (N₂O), and fluorinated gases.⁴ Warming is certain to continue until at least mid-century under even the most aggressive emissions reduction scenarios, with significantly different temperature ranges beyond 2050, depending on the emissions path.⁴ Together with changing physical conditions, three major climatic impact drivers (CIDs) – rising sea levels, intensifying storms, and increasing precipitation rates – increase the likelihood of hazardous flooding in Providence.²

RISING SEA LEVELS

Thermal expansion of oceans, mass loss from glaciers and ice sheets, and transfer of water from terrestrial aquifers to the ocean have led to a global mean sea level (GMSL) rise of 7.9 inches between 1901–2018.⁴ Rhode Island's tide gauges have risen even higher.^{5,6,7} Observations from the National Oceanic and Atmospheric Administration's (NOAA) Providence Tidal Station show that Relative Sea Level (RSL) rose 9 inches from 1938–2020, a higher rate than the global average due, in part, to local land subsidence and ocean circulation.^{5,6} The rate will likely accelerate non-linearly through the 21st century. Under a high emissions scenario, Narragansett Bay's RSL is projected to rise an additional 3.7 inches by 2030, 8.3 inches by 2040, 13.8 inches by 2050, 2.5 feet by 2070, and 4 feet by 2100, according to the United States Army Corps of Engineers (USACE) and NOAA.⁶ The rising high tide level will cause the city's coastal floodplain to encroach inland toward downtown, expanding the area prone to flood events.

STORM INTENSITY

The Northeast is experiencing intensification of tropical cyclones and extratropical storms due to warmer sea surface temperatures and the poleward shift of North Atlantic storm tracks.^{2,4,8} These storms temporarily raise sea level – called 'storm surge' – due to several factors, including changes in atmospheric pressure, high winds and wave activity that push water up against a coastline. According to storm modelers at the University of Rhode Island, Narragansett Bay's

funnel-like shape further amplifies the height of a storm surge as it moves up the bay.² Examples of storm surges in Narragansett Bay include the Great New England Hurricane of 1938 (Category 3), Hurricane Carol (1954, Category 3), Hurricane Diane (1955, Category 1), Hurricane Bob (1991, Category 2), Superstorm Sandy (2012), and Tropical Storm Henri (2021). More intense coastal storms coupled with a significant rise in the RSL will generate greater storm surges surpassing historical benchmarks and extending further inland. Importantly, a future storm-driven 1-in-20 year flood on top of a two-foot SLR will have the same water level and depth as today's 100-year flood.⁷


PRECIPITATION RATES

The warming atmosphere causes higher evaporation rates, which increase humidity, rainfall totals, and frequency of rainfall events. Many regions of the contiguous United States have experienced this trend since the 1950s, but it is most pronounced in the Northeast.^{5,8} From 1895 to 2011, average annual precipitation in the Northeast increased 10%, and since 1958 extremely heavy precipitation increased 70%.⁹ In Rhode Island, the number of heavy downpours (downpour = top 1% heaviest rain events) has increased 104% since 1950, a higher increase than in any other state.¹⁰ This year's July was Rhode Island's third wettest July on record with 7.12 inches of rainfall, nearly double the historical average.¹¹ Only July of 1916 and 2009 were wetter, with 9.09 and 10.32 inches of rainfall respectively.¹¹ By the end of the century, the number of 24-hour events per year that produce at least 1 inch of precipitation is projected to increase in all emissions scenarios.^{4,10} This rainfall trend presents risk of inland and urban flooding which may be compounded by simultaneous coastal flood events and overwhelmed drain-age system infrastructure.

IMPACT ON PROVIDENCE

Taken together, the climatic impact drivers make sections of Providence vulnerable as seen in the flood map (see **Figure 1**).¹ Even one additional foot of sea level rise, predicted by mid-century, would multiply the impact of flooding, and thereby affect numerous facilities in an expanded flood zone. Critically affected infrastructural assets include commercial and industrial port facilities, drinking water supply systems, wastewater treatment plants and stormwater systems, along with roadways, the FPHB, and some residential neighborhoods. Healthcare facilities located south of the FPHB, near the port area, are particularly vulnerable to inundation or access disruption. These include Hasbro Children's Hospital, Rhode Island Hospital, and Women & Infants Hospital,

Figure 1. FEMA Flood Zones in Providence, RI.



as well as several smaller facilities.¹ PEMA's 2019 assessment of flood-related hazards show that these three hospitals/healthcare facilities would be significantly impacted by the storm surge of a Category 1 hurricane, while numerous additional facilities would be devastated by a Category 3 hurricane storm surge.^{1,2} As seen in Hurricanes Katrina and Harvey, flooding and associated storm winds and rain can also cause havoc from above by damaging protective structures such as roofing, and damage hospitals outside the flood zone. Destruction of critical hospital infrastructure, including generators, HVAC systems, water supply, and the medical supply chain can lead to additional downstream emergencies.^{12,13}

CONCLUSION

The risk of flooding in Providence will become more severe in the coming decades. HVAs are constructed using historical data, but as Baugh et al. emphasize, we live in an era of climate change, making past data unreliable in estimating an accurate level of risk for Providence's critical healthcare facilities.¹⁴ Coalitions including the Providence Resilience Partnership (PRP) have emphasized the need for a predictive flood model incorporating climate change, land-cover types, hurricane barrier dynamics, and diverse stakeholder needs, but a common resilience strategy does not yet exist.² It is important that the city of Providence, the Rhode Island Sea Grant, the RI Department of Health, RI Emergency Management Agency (RIEMA) and the city's hospitals develop a plan based on future projections, as the natural hazard of flooding poses a significant risk of harm to the city, its hospitals, and the frontline communities they serve.

References

1. Rhode Island Emergency Management Agency (RIEMA) "City of Providence Strategy for Reducing Risks from Natural, Human-Caused and Technologic Hazards: A Multi-Hazard Mitigation Plan, April 2019" <http://www.riema.ri.gov/planning/hazardmitigation/index.php>
2. Rubinoff P, Elwell M, Kennedy S, Hallisey N, Squillante L. Towards a Resilient Providence: Issues, Impacts, Initiatives, Information. Providence Resilience Partnership (PRP). February 2021 Combined_3_25_final-low-res.pdf (providenceresilience.org)
3. Narragansett Bay Estuary Program. 2017. State of Narragansett Bay and Its Watershed (Executive Summary, pages 3-6). Technical Report. Providence, RI.
4. IPCC, 2021: Climate Change 2021: The Physical Science Basis. Contribution of Working Group I to the Sixth Assessment Report of the Intergovernmental Panel on Climate Change [Masson-Delmotte, V., P. Zhai, A. Pirani, S. L. Connors, C. Péan, S. Berger, N. Caud, Y. Chen, L. Goldfarb, M. I. Gomis, M. Huang, K. Leitzell, E. Lonnoy, J. B. R. Matthews, T. K. Maycock, T. Waterfield, O. Yelekçi, R. Yu and B. Zhou (eds.)]. Cambridge University Press. In Press
5. Rhode Island Coastal Resources Management Council (CRMC). "Rhode Island Shoreline Change Special Area Management Plan," 2018. <http://www.beachsamp.org/>
6. Sea-Level Change Curve Calculator (Version 2019.21) https://cwbiapp.sec.usace.army.mil/rccslc/slc_calc.html
7. Coastal Risk for Rhode Island, USA, Climate Central, https://riskfinder.climatecentral.org/county/providence-county.ri.us?-comparisonType=county&forecastType=NOAA2017_int_p50&level=7&unit=ft&zillowPlaceType=postal-code. Accessed 10/7/21
8. Dupigny-Giroux, LA, Mecray EL, Lemcke-Stampone M, Hodgkins G, et al. A risk based assessment of the impacts of a changing climate in the U.S. Northeast. 2018: Northeast. In *Impacts, Risks, and Adaptation in the United States: Fourth National Climate Assessment, Volume II* in DR Reidmiller, CW Avery, DR Easterling, KE Kunkel, et al. (eds.) U.S. Global Change Research Program, Washington, DC, USA, pp. 669–742. doi: 10.7930/NCA4.2018.CH18
9. "What Climate Change Means For Rhode Island." United States Environmental Protection Agency. August 2016. EPA 430-F-16-041.
10. Across U.S., Heaviest Downpours On The Rise." Research Report by Climate Central. May 27th, 2015. <https://www.climatecentral.org/news/across-us-heaviest-downpours-on-the-rise-18989>. Accessed September 25, 2021
11. National Weather Service Records. NOAA National Centers for Environmental Information (NCEI) Climate Data Online (CDO). <https://www.weather.gov/wrh/Climate?wfo=box>
12. Hines E, Reid CE. Hurricane Harvey Hospital Flood Impacts: Accuracy of Federal Emergency Management Agency Flood Hazard Areas in Harris County, Texas. *Am J Public Health*. 2020 Apr;110(4):574-579. doi: 10.2105/AJPH.2019.305520. Epub 2020 Feb 20. PMID: 32078350; PMCID: PMC7067069.
13. Sheri Fink. Five Days at Memorial: Life and Death in a Storm-ravaged Hospital. New York: Crown Publishers, 2013.
14. Baugh J, Kemen K, Messervy J, Biddinger P. Beyond the Hazard Vulnerability Analysis: Preparing Health Systems for Climate Change. *Rhode Island Medical Journal*. 2021;104(9).

Authors

Andrew E. Binder, BA, Dartmouth College, Hanover, NH.

Selim Suner, MD, MS, Professor of Emergency Medicine, Surgery, and Engineering, Brown University; Director of Disaster Medicine and Emergency Preparedness, Alpert Medical School of Brown University.

H. Curtis Spalding, MPA, Professor of the Practice Institute at Brown for Environment and Society (IBES), Brown University; former Regional Administrator of EPA Region 1 New England, and Executive Director of Save the Bay, Providence, RI.

Erich Osterberg, PhD, Associate Professor of Earth Sciences, Dartmouth College, Hanover, NH.

Correspondence

Andrew.E.Binder.21@dartmouth.edu

A Case of Heat Stroke in the Era of Climate Change

FRED VARONE, MD; WILLIAM BINDER, MD

*From the Case Records of the Alpert Medical School
Residency in Emergency Medicine*

DR. FRED VARONE: Today's patient is an elderly male of unknown age (later found out to be 68 years old) who was brought into the Emergency Department (ED) by Emergency Medical Services (EMS) after being found unresponsive in a park. The patient was fully dressed and wearing a coat, lying face down, despite outside temperatures hovering at 99°F and > 90% humidity.

Initial vital signs were notable for a systolic blood pressure of 70 mm Hg, pulse 130s, and a rectal temperature of 43.3°C (110°F). The patient was noted to be unresponsive to painful stimuli and his skin was dry and warm to touch. Pupils were 3 millimeters bilaterally and sluggishly reactive. His cardiac exam revealed tachycardia with a regular rhythm. Breath sounds were clear although he was noted to have sonorous respirations.

The patient was given a bolus of crystalloid IV fluids and push dose phenylephrine for hypotension. He was intubated for airway protection via rapid sequence intubation using ketamine and rocuronium.

DR. AMY MATSON: What laboratory and imaging studies were obtained?

DR. VARONE: Notable labs are seen in **Table 1**. The patient's laboratory abnormalities are not unusual in heat stroke. Rhabdomyolysis and injury to the kidneys and liver are often noted by elevations in the creatinine phosphokinase (CPK), creatinine and liver function tests. Thrombocytopenia is also

common. Troponin is frequently elevated as well, reflecting a type 2 myocardial infarction.^{1,2,3,4,5}

There are no specific findings on radiographic imaging suggestive of heat stroke. When acute lung injury and ARDS are suspected, chest X-ray will show the typical bilateral infiltrates. In our patient, a chest X-ray after intubation showed the endotracheal tube in good position and no acute cardiopulmonary abnormalities. CT imaging of the brain and cervical spine without IV contrast was unremarkable for acute changes. CT imaging of the abdomen and pelvis with intravenous contrast revealed diffuse thickening of bowel loops with enhancing walls concerning for bowel ischemia, as well as bibasilar airspace disease.

DR. JOSEPH LAURO: With a temperature of 110°F, this patient has heat stroke. Is it unusual for a temperature to rise so high without exertion? Can you review the different types of heat stroke?

DR. VARONE: Heat-related illness exists on a spectrum of disease severity, with heat stroke being the most severe manifestation. Other classifications for heat-related illnesses include heat stress, heat injury, heat exhaustion, heat syncope, and heat cramps. The loss of fluid and salt in sweat can lead to muscle cramping and intravascular volume depletion leading to syncope. Heat exhaustion can be considered the precursor to heat stroke, with symptoms such as headache, dizziness, nausea, weakness, and a temperature that is typically between 36 and 40°C. While these are all readily reversible clinical conditions with cooling and hydration, they can progress into organ failure if untreated, leading to the syndrome of heat stroke.

Our patient has heat stroke, classically defined as high environmental heat exposure combined with CNS dysfunction and elevated core body temperature > 40°C (104°F).⁶ Additional definitions have been proposed which do not include temperature criteria (See **Table 2**).⁷

Heat stroke can be divided into classic heat stroke (CHS) and exertional heat stroke (EHS). CHS occurs more often in vulnerable populations, such as the elderly, as their ability to compensate is often impaired by comorbidities and medications. EHS, on the other hand, occurs in

Table 1. Heat Stroke Criteria

Lab	Initial presentation	Day 1 of hospitalization	Day 3 of hospitalization
Hemoglobin/Hematocrit	10.9/33	11.8/34	
Platelets	64 x 10 ⁹ /L	21 x 10 ⁹ /L	74 x 10 ⁹ /L
Creatinine	1.80 mg/dl	2.51 mg/dl	5.43 mg/dl
Troponin I	0.166 ng/ml	68.194 ng/ml	65.271 ng/ml
AST/ALT	125/36 IU/L	4,060/354 IU/L	949/302 IU/L
CPK	243 IU/L	6,638 IU/L	4,555 IU/L
Lactate	9.3 mEQ/L	4.6 mEQ/L	

Table 2. Comparison of different criteria for diagnosis of heat stroke

All definitions	<ul style="list-style-type: none"> • High Environmental Heat Exposure
Bouchama	<ul style="list-style-type: none"> • Core temperature > 40°C • CNS effects (coma, delirium, seizures)
JAAM	<ul style="list-style-type: none"> • CNS symptoms (Japan Coma Scale score of ≥ 2, cerebellar symptoms, convulsions/seizures) • Hepatic or renal dysfunction • DIC by JAAM criteria
JAAM-HS-WG	<ul style="list-style-type: none"> • GCS ≤ 14 • JAAM DIC score ≥ 4 • Creatinine or bilirubin sym 1.2 mg/dl

Glasgow Coma Scale, Japanese Association of Acute Medicine (JAAM), Japanese Association of Acute Medicine heat stroke committee working group (JAAM-HS-WG)

young, healthy individuals during strenuous physical activities (athletes, military personnel).^{7,8} Temperatures can rise substantially in both forms of heat stroke.

In patients with heat-related illness, heat loss is mediated by several different mechanisms. These include radiation, conduction, convection, and evaporation. In cool temperatures, the majority of body heat (55%–65%) is lost by radiation, or the emission of energy from the skin. Convection occurs through the transfer of heat from the body to a gas (air), while conduction occurs by the transfer of heat between two surfaces with different temperatures. In warm temperatures, evaporation is the primary means to dissipate heat, but it has limited efficacy when humidity rises above 60%.¹ Unfortunately, our patient was unable to dissipate heat due to his clothing and the high heat index (a combination of air temperature and relative humidity).

DR. ELIZABETH SUTTON: This patient's temperature reached 110°F. Is that usually survivable? What happens to the patient's organs?

DR. VARONE: While organ systems may start to exhibit dysfunction at 40°C (104°F), higher temperatures are survivable, and patients may have temperatures up to 43.3°C (109.9°F) and survive with good neurologic outcome.⁹ Some studies suggest that time to cooling may be more important than maximum temperature.^{3,10,11}

The cause of an elevated temperature is important. Fever is created when pyrogens (such as bacterial endotoxins or cytokines produced from cell damage) act on the hypothalamus to increase the natural temperature set point through increased prostaglandin production.⁷

In heat stroke, however, temperatures rise without an adjustment in the hypothalamic set point. In heat injury, cell membrane physiology is disrupted, leading to endothelial and organ cell death.^{2,12,13} Proteins start to denature around 40°C, leading to enzyme disruption, mitochondrial failure, and loss of oxidative phosphorylation. In response to injury, cells produce heat-shock proteins, which are protective against heat, hypoxia, and ischemia, but these quickly

can become overwhelmed in severe heat stroke.¹ The end result is cell death across multiple systems.¹⁴

The impact of hyperthermia on organ systems is significant. In the CNS, thermal injury can lead to cellular denaturing, neuronal death, and excitotoxicity. Cytokine release and resultant vascular damage, as well as local ischemia and systemic changes in blood flow can result in cognitive impairment, seizures, and cerebral injury.¹⁵

Gastrointestinal tract physiology is altered by hyperthermia and contributes to mortality. To dissipate heat, blood flow shunts from the GI tract to the skin, resulting in bowel ischemia, mucosal barrier disruption, and bacterial translocation. Bacteremia, circulating endotoxin and sepsis can ensue.^{1,3,13} Additional inflammatory mediators, including IL-1 and IL-6, are released from injured cells, causing a systemic inflammatory syndrome response (SIRS).^{1,7} Vascular endothelial cells also sustain damage, which can lead to a consumptive coagulopathy and systemic microthrombi.^{2,13}

Other organ systems are impacted. Myocardial infarction has been reported in up to 7% of admitted heat stroke patients.⁴ Multi-organ dysfunction and systemic manifestations including disseminated intravascular coagulation, rhabdomyolysis, hepatic necrosis, and acute kidney injury have all been reported.^{5,16}


DR. OTIS WARREN: Are heat-related illness occurring with greater frequency?

DR. WILLIAM BINDER: There has been an increase in the frequency of extreme heat events (EHEs) over the last several decades. Data analyzed between 1961 and 2019 for 50 metropolitan areas in the US shows an increase in heat wave frequency, duration, intensity, and an increase in the length of the heat wave season (see **Figure 1**).¹⁷

Heat related mortality has risen concordantly. While it is reported that approximately 700 individuals die annually in the US due to heat, it is felt that this number is grossly underestimated as comorbid diseases are often reported as a primary cause of death, thereby concealing environmental heat's role.^{1,18} The recent Lancet countdown on health and climate change suggests that in 2018, 19,000 deaths in the US were related to high environmental temperatures.¹⁹ Underscoring this, our patient's problem list and discharge summary in the electronic health record acknowledges NSTEMI, encephalopathy, and acute kidney injury, but makes no mention of heat stroke.

While data on heat-related mortality in Rhode Island is not available, average temperatures have risen over the past several decades. Between 1981–2010 there were an average of 9 days $\geq 90^\circ\text{F}$ in the Providence Metro area; this number increased between 2010 to 2014, and in 2021 there were 19 days $\geq 90^\circ\text{F}$. Rising summer daily temperatures have been linked to increased EMS utilization and ED visits.^{20,21}

Figure 1. Heat-wave frequency, duration, intensity, and length of season between 1961–2019 for 50 metropolitan areas in the US.



DR. VARONE: Anthropogenic climate change from increasing urbanization has resulted in heat islands, areas where the average temperature is higher than surrounding areas. Developed cities tend to have higher day and nighttime temperatures due to increased heat production, replacement of vegetation by concrete and other manmade materials, and disruption of natural airflow by large buildings, streets, and city geometry.²² Higher evening temperatures prevent humans from recovering from daytime heat exposure and injury. Analysis of the 2003 Paris heat wave shows a clear association between increased nighttime surface temperature in heat islands and risk of death, particularly in elderly people.²²

While cities only make up around 2% of the Earth's surface, they are responsible for 75% of the world's emissions.²³ In 2007 there were 19 cities with 10 million or more inhabitants; by 2018 there were 33 "mega-cities." Increasing urbanization will lead to more heat islands and more heat-related illness, particularly in developing countries where populations and urbanization are still accelerating.^{23,24}

DR. CATHERINE CUMMINGS: What was your management of this patient with hyperthermia?

DR. VARONE: Rectal temperatures should be checked in patients with suspected heat stroke. External measurements have been shown to correlate poorly with rectal temperature and underestimate the degree of hyperthermia.^{8,11} In one study of collapsed marathon runners, only two of 17 collapsed runners with true hyperthermia (rectal temperature $\geq 39.4^{\circ}\text{C}$) were identified correctly by temporal measurement.²⁵

Rapid cooling is the mainstay of treatment for heat stroke and improved outcomes are related to early recognition and treatment.^{6,8,10,11,12} Patients should be cooled at a rate of $0.1\text{--}0.2^{\circ}\text{C}$ per minute until evidence of clinical improvement, with careful attention paid to preventing overcooling.^{6,26} While there is no specific temperature goal and the patient's clinical picture should guide cooling, some studies have used specific temperature cutoffs to prevent overcooling ($38\text{--}39^{\circ}\text{C}$).^{6,10} Techniques used for cooling in the ED include cold packs and ice (conduction), and evaporative cooling using cold water and fans. Cold-water immersion, frequently used in the pre-hospital setting for patients with EHS, is often impractical in the ED and potentially harmful in older patients with CHS requiring monitoring. Invasive methods for cooling include bladder, gastric and peritoneal lavage, and cardiopulmonary bypass methods such as ECMO-based cooling.²⁶

Our patient received intravenous fluids, cold packs and ice, and evaporative cooling with fans. His temperature decreased to 39°C within 45 minutes of arrival to the ED.

Of note, antipyretics have no utility in hyperthermia due to environmental exposure, and some studies suggest they may be harmful in heat-related injury.²⁶

DR. NICK MUSISCA: What happened to this patient?

DR. VARONE: After intubation, orogastric and rectal tubes were placed. His ED course was complicated by runs of atrial fibrillation with rapid ventricular response, and cardioversion was attempted but was unsuccessful. His rate decreased with fluid administration and temperature control. He received benzodiazepines, to prevent shivering and agitation, piperacillin-tazobactam for ischemic bowel, and anti-epileptic medications (AEDs).

The patient was admitted to the medical intensive care unit and multiple medical services were consulted. Continuous electroencephalography was performed for 24 hours, and the patient did not demonstrate seizure activity. On hospital day 4 he was extubated, although he was confused and encephalopathic. On hospital day 10 he was able to provide health care workers his name, and social work was able to locate his daughter. She reported a history of chronic alcohol abuse and dementia - known risk factors for heat stroke. The patient improved to his baseline and was discharged to a skilled nursing facility on hospital day 19.

References

- Bouchama A, Knochel JP. Heat Stroke. *N Engl J Med*. 2002; 346(25): 1978-1988.
- Leon LR, Helwig B. Heat stroke: Role of the systemic inflammatory response. *J Appl Physiol*. 2010; 109: 1980-1988. doi:10.1152/jappphysiol.00301.2010.
- Yang M, Li Z, Zhao Y, Zhou F, et al. Outcome and risk factors associated with extent of central nervous system injury due to exertional heat stroke. *Medicine (Baltimore)*. 2017 Nov;96(44):e8417. doi: 10.1097/MD.0000000000008417. PMID: 29095276; PMCID: PMC5682795.
- Bathini T, Thongprayoon C, Chewcharat A, et al. Acute Myocardial Infarction among Hospitalizations for Heat Stroke in the United States. *J Clin Med*. 2020;9(5):1357. Published 2020 May 6. doi:10.3390/jcm9051357
- Johnson RJ, Sánchez-Lozada LG, Newman LS, Lanaspá MA, Diaz HF, Lemery J, Rodríguez-Iturbe B, Tolan DR, Butler-Dawson J, Sato Y, García G, Hernando AA, Roncal-Jimenez CA. Climate Change and the Kidney. *Ann Nutr Metab*. 2019;74 Suppl 3:38-44. doi: 10.1159/000500344. Epub 2019 Jun 14. PMID: 31203298.
- Ruble C, Dresser C, Guidice C, Lemery J, Sorensen C. Evidence-Based Heatstroke Management in the Emergency Department. *West J Emerg Med*. 2021; 22(2):186-195.
- Hifumi, T, Kondo, Y, Shimizu, K. Heat stroke. *J Intensive Care*. 2018;6(30). <https://doi.org/10.1186/s40560-018-0298-4>.
- Casa DJ, DeMartini JK, Bergeron MF, Csillan D, Eichner ER, Lopez RM, Ferrara MS, Miller KC, O'Connor F, Sawka MN, Yeargin SW. National Athletic Trainers' Association Position Statement: Exertional Heat Illnesses. *J Athl Train*. 2015;50(9): 986-1000. doi: 10.4085/1062-6050-50.9.07. Erratum in: *J Athl Train*. 2017 Apr;52(4):401. PMID: 26381473; PMCID: PMC4639891.
- Argaud L, Ferry T, Le QH, Marfisi A, Ciorba D, Achache P, Ducluzeau R, Robert D. Short- and long-term outcomes of heatstroke following the 2003 heat wave in Lyon, France. *Arch Intern Med*. 2007 Nov 12;167(20):2177-83. doi: 10.1001/archinte.167.20.i0170147. Epub 2007 Aug 13. PMID: 17698677.
- Demartini JK, Casa DJ, Stearns R, Belval L, Carago A, Davis R, Jardine J. Effectiveness of Cold Water Immersion in the Treatment of Exertional Heat Stroke at the Falmouth Road Race. *Med Sci Sports Exerc*. 2015; 47(2):240-245. doi: 10.1249/MSS.0000000000000409. PMID: 24983342.
- American College of Sports Medicine, Armstrong LE, Casa DJ, Millard-Stafford M, Moran DS, Pyne SW, Roberts WO. American College of Sports Medicine position stand. Exertional heat illness during training and competition. *Med Sci Sports Exerc*. 2007; 39(3):556-572. doi: 10.1249/MSS.0b013e31802fa199. PMID: 17473783.
- Casa DJ, Armstrong LE, Kenny GP, O'Connor FG, Huggins RA. Exertional heat stroke: new concepts regarding cause and care. *Curr Sports Med Rep*. 2012 May-Jun;11(3):115-23. doi: 10.1249/JSR.0b013e31825615cc. PMID: 22580488.
- Epstein Y, Yanovich R. Heatstroke. *N Engl J Med*. 2019; 380(25):2449-2459.
- Aruga T, Miyake Y. Pathophysiology of heat illness. *Nihon Rinsho*. 2012 Jun;70(6):940-6. Japanese. PMID: 22690597.
- Walter EJ, Carraretto M. The neurological and cognitive consequences of hyperthermia. *Crit Care*. 2016;20(1):199. Published 2016 Jul 14. doi:10.1186/s13054-016-1376-4
- Satirapoj B, Kongthaworn S, Choovichian P, Supasynndh O. Electrolyte disturbances and risk factors of acute kidney injury patients receiving dialysis in exertional heat stroke. *BMC Nephrol*. 2016;17(1): 55
- (<https://www.epa.gov/climate-indicators/climate-change-indicators-heat-waves>. Accessed September 24, 2021)
- Vaidyanathan A, Malilay J, Schramm P, Saha S. Heat-Related Deaths – United States, 2004-2018. *MMWR Morb Mortal Wkly Rep*. 2020;69:729-734. DOI: <http://dx.doi.org/10.15585/mmwr.mm6924a1>
- Watts N, Amann M, Arnell N, et al. The 2020 report of the Lancet Countdown on health and climate change: responding to converging crises. *Lancet*. 2021; 397(10269):129-70.
- Kingsley SL, Eliot MN, Gold J, Vanderslice RR, Wellenius GA. Current and Projected Heat-Related Morbidity and Mortality in Rhode Island. *Environ Health Perspect*. 2016;124(4):460467.
- Moretti K, Gallo Marin B, Soliman LB, Asselin N, Aluisio AR. Increased Temperatures are Associated with Increased Emergency Medical Services Utilization in Rhode Island. *Rhode Island Medical Journal*. 2021;104(9).
- Laaïdi K, Zeghnoun A, Dousset B, Bretin P, Vandentorren S, Giraudet E, Beaudeau P. The impact of heat islands on mortality in Paris during the August 2003 heat wave. *Environ Health Perspect*. 2012; 120(2): 254-259. doi:10.1289/ehp.1103532
- Madlener R, Sunak Y. Impacts of urbanization on urban structures and energy demand: What can we learn for urban energy planning and urbanization management? *Sustainable Cities and Society*. 2011;1(1): 45<https://doi.org/10.1016/j.scs.2010.08.006>.
- Vicedo-Cabrera AM, Scovronick N, Sera F, et al. The burden of heat-related mortality attributable to recent human-induced climate change. *Nat. Clim. Chang*. 2021;11:492–500. <https://doi.org/10.1038/s41558-021-01058-x>
- Ronneberg K, Roberts WO, McBean AD, Center BA. Temporal artery temperature measurements do not detect hyperthermic marathon runners. *Med Sci Sports Exerc*. 2008; 40(8):1373-5. doi:10.1249/MSS.0b013e31816d65bb
- Laitano O, Leon LR, Roberts WO, Sawka MN. Controversies in exertional heat stroke diagnosis, prevention, and treatment. *J Appl Physiol*. 2019;127: 1338-1348. doi:10.1152/jappphysiol.00452.2019.

Authors

Fred Varone, MD, PGY-3, Brown Emergency Medicine Residency.
William Binder, MD, Associate Professor of Emergency Medicine,
Warren Alpert Medical School of Brown University.

Correspondence

william_binder@brown.edu



Image 1. Swedish icebreaker Oden.

Frozen Obsession documentary follows URI team studying climate change in the Arctic

MARY KORR
RIMJ MANAGING EDITOR

Frozen Obsession captures a once-pristine Polar ecosystem eroding at an alarming and accelerating rate. The PBS documentary by Emmy Award-winning filmmaker **DAVID CLARK** chronicles the Northwest Passage Project (NPP) expedition studying the effects of climate change in the Canadian Arctic. A team at the University of Rhode Island (URI) Graduate School of Oceanography (GSO) and its Inner Space Center, an international research and education hub, developed the project.

Aboard the Swedish icebreaker *Oden*, (**Image 1**), the expedition departed from the U.S. Thule Air Base in Greenland in July 2019, on a 2,000-mile voyage through the Northwest Passage (**Figure 1**). During a pre-release viewing of the film, followed by a ZOOM panel discussion, planners and participants shared their experiences during the 18-day mission.


URI Associate Professor of Oceanography **BRICE LOOSE, PhD**, the project's chief scientist, described the Arctic as "a melting pot. The old Arctic is disappearing. I feel like we are watching a tragedy unfolding." He noted the dramatic shifts in the climate record. "Humans were not

on the planet the last time there was a major change in the climactic pattern like the one we are experiencing today. Previously it has been over the course of several thousand to a hundred thousand years. What we are seeing today is a profound change because of the speed with which it is unfolding. It is not even a matter of decades now, but years."

Loose described the NPP research as "basic science, data collection. It was an

opportunity to peer into a complex system and start to understand it for the first time." Clark's documentary offers fascinating glimpses into this; one scene depicts Loose and a small group helicoptered to sea ice floes, where they drilled into six-foot thick ice to collect core samples (**Image 2**). What they found was disturbing – plastics of different types, sizes and colors. "There was so much plastic that you could look at it with your naked

Figure 1. Map of the Northwest Passage Project Expedition.



Q&A with oceanographer Brice Loose, PhD, NPP project lead scientist

MARY KORR
RIMJ MANAGING EDITOR



Q. What kind of routine medical screenings were done for the pre-COVID Northwest Passage Project (NPP) and what is taking place now on research expeditions?

A. Participants have to go through a fairly rigorous medical screening with certification from their physician and dentist that the likelihood of needing emergent care is low within the next 3-6 months. People are sometimes disqualified for heart and eye conditions, for example.

Q. Was the medical team on the ship from New England?

A. Typically, the medical personnel are employed by the ship operator. In this case, the Swedish Polar Research Secretariat had a staff physician aboard.

Q. It did not sound like there were any medical emergencies, but if there were, would a patient be helicoptered out?

A. Yes, the helicopters are critical in this respect. They can rapidly transfer someone to a town with a hospital or with airline service to larger towns. In the Northwest Passage, we were often less than one day away from a community that was accessible by helicopter. These townships are serviced by two Canadian airlines, so the field setting was not as remote as it can be.

Q. Are there research findings from the Project you would like to highlight?

A. For NPP, the other results are preliminary and still in progress, but we see the influence of terrestrial glaciers affecting the distribution of plankton and microbes, and potentially even the production and degradation of greenhouse gases. Despite the extended daylight, we found very little exchange of nutrients between land and sea. That, combined with the enhanced freshwater on top of seawater, meant that very little plant nutrient was available in the surface ocean to fuel plankton growth by primary production. In these respects, extended summer melt is really transforming the region.

I'd also like to make you aware of the **MOSAIC project**: <https://mosaic-expedition.org>. In total there were four URI members who participated in MOSAIC during the course of the year-long drift. We had some serious logistical hurdles to transfer personnel during the spring crew changes, but a protocol was developed so that COVID never made it out to the vessel.

Q. Is there follow-up research to the NPP taking place in the Canadian Arctic?

A. We are in collaboration with Canadian Dept. of Fisheries and Wildlife that have long-term monitoring in this area. Their work is ongoing. One of our graduate students has been journeying to the Canadian Arctic and the NWP almost every summer for several years. ❖



eye and see all of the beads, fibers and filaments just sitting there in the bottom of the containers," Loose said.

Previous European research has shown the trajectory of plastics, determining it lofts through air masses, descends on Arctic islands, and flows with ocean currents. Some of the sea ice collected during the 2019 expedition is believed to have started in the Central Arctic, moved east through the Nares Strait by Greenland, and then into the Northwest Passage's Lancaster Sound.

Analyses into the data collected during the NPP project is ongoing; for example, identifying the state of weathering on the ice samples and trying to determine the degradation of the plastics, and potentially tracing its source to the manufacturer.

"It is important for people everywhere on Earth to see and understand how this region affects all of us," said NPP principal investigator and project director **GAIL SCOWCROFT**. "The region's meltwater, water circulation, and exchange of greenhouse gases between the ocean and the

Image 2. Dr. Brice Loose, at right, chief expedition scientist with the Northwest Passage Project and research team collecting samples from ice cores in the Arctic.

[COURTESY NORTHWEST PASSAGE PROJECT/
CAMERA: DUNCAN CLARK]



Image 3, 4. Polar bear and iceberg in the Canadian Arctic waters.
[COURTESY OF THE NORTHWEST PASSAGE PROJECT.]

atmosphere are causing wide-scale environmental and climatic changes, which increasingly affects people and wildlife diversity around the world." (Images 3, 4)

However, she pointed to encouraging signs, with the United States rejoining the Paris climate agreement this year, and also stressed the urgency for individuals to act. "We can, as citizens, reduce

our own carbon footprint and encourage our governments to move towards a zero-emissions economy."

On the voyage, GSO microbiology graduate student **JACOB STROCK** supervised the observation and collection of plankton samples, "to understand what is in the water and how susceptible these 'small but mighty' microscopic

organisms might be to climate change. Many small organisms can have a large effect. What impact will it have on the Arctic as a whole?" he pondered during the panel discussion.

Filmmaker Clark said he hoped the film's takeaway for viewers would be to "believe the science, and see that climate change is real. We look to the next generation of young scientists and decision-makers; they are the ones who are going to save us."

Frozen Obsession also explores the maritime history of the Northwest Passage, and the history and effects of climate change on indigenous populations. The latter perspective was compelling, given by student participant **MIA OTOKIAK**, Inuit scientist and community liaison.

The NPP was supported with major funding from the U.S. National Science Foundation and additional support from the Heising-Simons Foundation.

For further information on the project and how to view the documentary, visit <https://northwestpassageproject.org>.



Complete Resolution of Traumatic Indirect Carotid-Cavernous Fistula Two Years After Inciting Event

JAMES O. ROBBINS, BA, MD-ScM'22; NISARG CHHAYA, MD; JAMIE L. SCHAEFER, MD

ABSTRACT

Indirect carotid cavernous fistulas (CCFs) are most often spontaneous, but can rarely be caused by trauma. With traumatic etiology, the timeline for the development of symptoms varies significantly and can be difficult to predict. In this report, we discuss the case of a patient found to have an indirect CCF who presented for acutely worsening ocular symptoms and a history of pulsatile tinnitus that began two years prior after a suspected inciting head injury. To our knowledge, no cases have described a traumatic indirect CCF with a similarly extensive indolent course who demonstrated full symptomatic recovery following treatment.

KEYWORDS: carotid-cavernous fistula, indirect, trauma, endovascular

CASE PRESENTATION

An 82-year-old female with hypothyroidism and no significant ocular history presented to the clinic with three months of left eye pain, redness, decreased vision, and photophobia (**Figure 1A**). She also endorsed persistent left-side facial pain and an intermittent “whooshing” noise in the ipsilateral ear, which began after a fall with loss-of-consciousness two years prior.

Exam on presentation was significant for best-corrected visual acuity (BCVA) of 20/30 OD and 20/800 OS, intraocular pressure (IOP) of 16 mmHg OD and 19 mmHg OS, 2+ afferent pupillary defect (APD) OS, and diffuse visual field deficits on the left with partial sparing of the superior nasal field. External exam showed proptosis OS with exophthalmometry (15mm OD and 20mm OS). Slit lamp examination was significant for prominent episcleral corkscrew vessels OS and no clear posterior findings.


Computed tomography angiogram (CTA) of the head and neck demonstrated asymmetric enlargement of the left superior ophthalmic vein, and asymmetric enhancement of the left cavernous sinus overall suggestive of an indirect (Barrow's type B, C, or D) carotid cavernous fistula (CCF) (**Figure 2A, 2B**).

The patient was hesitant to undergo an endovascular procedure at her initial visit, but at subsequent evaluation

Figure 1. Patient photograph at initial presentation to the clinic (**A**), at one-week post-operative follow-up (**B**), and at one-month post-operative follow-up (**C**).



Figure 2. Computed Tomography Angiography (CTA) with coronal (**A**) and axial (**B**) views of the dilated left superior ophthalmic vein. Interventional angiograms with arrows demonstrating the patent fistula (**C**) and the resolved fistula following coil embolization (**D**).



decided to proceed with treatment secondary to persistent ocular and facial pain. She was admitted to the hospital and, after an initial failed endovascular approach via peripheral venous access that identified thrombosis of the distal ophthalmic veins, underwent successful coil embolization of the fistulous pouch by direct transorbital puncture of the cavernous sinus via the superior ophthalmic vein (**Figure 2C, 2D**). She was discharged on the first postoperative day and asked to return to the clinic for follow-up in one week.

Exam at one-week follow-up demonstrated BCVA of 20/40 OS, exophthalmometry of 16.5mm OS, and only trace APD OS. Slit lamp examination revealed improved episcleral corkscrew vessels and a new subconjunctival hemorrhage temporally OS (**Figure 1B**). The patient reported mild frontal headache, but otherwise felt well with no ocular pain. Patient reported near-complete resolution of presenting symptoms after one month (**Figure 1C**).

At five-month follow-up, BCVA was 20/25 OS, exophthalmometry was 15.5mm OS, and no APD was present. Slit lamp exam revealed resolution of episcleral corkscrew vessels and subconjunctival hemorrhage. The patient endorsed some residual left-side facial pain, but otherwise complete resolution of ocular symptoms and the pulsatile tinnitus that has persisted for two years prior.

DISCUSSION

Indirect CCFs, also called “low-flow” fistulas, are abnormal communications between the cavernous sinus (CS) and dural branches of the internal carotid artery (Barrow type B), the external carotid artery (Barrow type C), or both (Barrow type D). In contrast, direct CCFs (Barrow type A) connect the internal carotid artery and CS without intermediary tracts, and thus function under higher arterial pressures. As such, the presentation of indirect fistulas is more indolent as accommodations in venous drainage from the CS may delay the onset of visual loss, proptosis, and redness.

The etiology of indirect CCFs is often unclear. Unlike with direct fistulas, obvious precipitating trauma is relatively rare. It has been postulated that some indirect CCFs may occur secondary to a subclinical CS thrombosis that causes local revascularization¹. Other risk factors include pregnancy, local surgical instrumentation, and sinusitis.²

As the anatomy of indirect CCFs is highly variable, so too is the natural history. Spontaneous closure is relatively common, with estimates varying between 40–90% depending on duration of observation and means of non-invasive treatment.^{3,4} Therefore, conservative management may be a consideration depending on patient preference and severity of presenting symptoms. Factors that have demonstrated

increased risk for poor prognosis include progressive vision loss, proptosis, and signs of elevated intracranial pressure.²

Our patient reported a two-year history of ipsilateral facial pain and pulsatile tinnitus, which began after a fall with loss-of-consciousness, progressing to include ipsilateral ocular pain and decreased vision starting three months prior to presentation. Her history, combined with the cessation of tinnitus following treatment, strongly suggests that the fall two years prior was responsible for the indirect CCF. While a traumatic indirect CCF is a highly uncommon presentation, perhaps more remarkable is this patient’s timeline of symptoms.

To our knowledge, no cases of traumatic CCFs have been described with such a lengthy indolent course, particularly given the patient’s full recovery following treatment. It is unclear why the current patient was symptomatic, but stable, for two years with a precipitous worsening of symptoms. Possible causes for the worsening of her symptoms may include an acute anatomical change to the fistulous connection resulting in increased inflow to the CS, similar to the acute process responsible for traumatic, direct (Barrow type A) CCFs. Also possible is an acute congestion of the venous outflow tract, such as with a venous thrombosis, leading to decreased drainage of the sinus.

The current case demonstrates that an unusually long symptomatic time to presentation should not steer the clinician away from adding indirect CCF to the differential diagnosis for symptoms including, but not limited to, red eye, chemosis, or ocular pain that is refractory to conservative management. The misdiagnosis of indirect CCF as more common benign conditions can cause delays in treatment and possible permanent ocular damage⁵. If signs of smoldering indirect CCF are correctly identified early by caregivers, episodes of severe exacerbation with threatened vision loss and potential permanent ocular damage may be avoided.

The current case also demonstrates that even after a two-year symptomatic period with an acute clinical decompensation, endovascular therapy can be highly efficacious. The outcomes of various endovascular interventions for patients with traumatic indirect CCFs has been described, but more research is needed to delineate the relationship between symptomatic time-to-presentation and degree of recovery post-intervention.^{6,7,8} We demonstrate here full symptomatic resolution of an indirect, traumatic CCF following a time course not previously documented. In future encounters, it may be warranted to pursue consultation for endovascular therapy as soon as an indirect CCF is suspected, irrespective of etiology, degree of vision loss, or extent of symptomatic time course.

References

1. Taniguchi RM, Goree JA, Odom GL. Spontaneous carotid-cavernous shunts presenting diagnostic problems. *J Neurosurg.* 1971; 35(4):384-91.
2. VV Halbach, GB Hieshima, RT Higashida, M Reicher. Carotid cavernous fistulae: indications for urgent treatment. *Am. J. Roentgenol.* 1987; 149:3, 587-593.
3. Nukui H, Shibasaki T, Kaneko M, Sasaki H, Mitsuka S. Long-term observations in cases with spontaneous carotid-cavernous fistulas. *Surg Neurol.* 1984; 21:543-552.
4. Newton TH, Hoyt WF. Dural arterial venous shunts in the region of the cavernous sinus. *Neuroradiology.* 1970; 1:71-81.
5. Canellas, M, Cheema, N. Misdiagnosed spontaneous carotid cavernous sinus fistula. *Clin Pract Cases Emerg Med.* 2019; 3(3), 256–258.
6. Luo CBTM, Chang FC, Chang CY. Traumatic indirect carotid cavernous fistulas: angioarchitectures and results of transarterial embolization by liquid adhesives in 11 patients. *Surg Neurol.* 2009;71(2):216–22.
7. Lin, HL, Hu, TT. Isolated third nerve palsy with pupillary involvement resulting from carotid-cavernous sinus fistula: a case report. *Medicine (Baltimore).* 2019;98(6), e14472.
8. Jacobson, BE, Nesbit, GM, Ahuja, A, Barnwell SL. Traumatic indirect carotid-cavernous fistula: report of two cases. *Neurosurgery.* 1996;39(6):1235–8.

Authors

- James O. Robbins, BA, MD-ScM'22, The Warren Alpert Medical School of Brown University, Providence, RI.
- Nisarg Chhaya, MD, Division of Ophthalmology, Warren Alpert Medical School of Brown University, Providence, RI.
- Jamie L. Schaefer, MD, Division of Ophthalmology, Warren Alpert Medical School of Brown University, Providence, RI.

Correspondence

James O. Robbins
 222 Richmond Street
 Providence, RI 02903
james_robbins@brown.edu

An Unusual Presentation of Kikuchi-Fujimoto Disease with Recurrent Fever and Syncope

QUYNH-LAN DAO, MD; BRIDGET HOTTENSTEIN, MD; VIJAI RAM SELVARAJ, MD, MPH

ABSTRACT

Kikuchi-Fujimoto (KF) disease, also known as necrotizing histiocytic lymphadenopathy, is a rare, benign disorder of the lymph nodes of young adults, predominantly young females. The exact cause of KF disease is unknown. Here, we report a young female with recurrent fever and syncopal episodes diagnosed with KF disease on lymph node biopsy.

KEYWORDS: Kikuchi-Fujimoto disease, Kikuchi's disease, syncope, fever of unknown origin

INTRODUCTION

Kikuchi-Fujimoto (KF) disease, also known as Kikuchi's disease, or necrotizing histiocytic lymphadenopathy, is a rare cause of fever of unknown origin (FUO). Most common causes of FUO are infections, malignancy, or autoimmune disease. We present a unique case of KF disease in a young female with Hodgkin's lymphoma who presented with recurrent fevers and syncope.

CASE PRESENTATION

A 34-year-old female with a history of Hodgkin's lymphoma status post-six cycles of Brentuximab Vedotin- doxorubicin/ vinblastine/dacarbazine (July 2018–Dec 2018) in complete remission presented to the hospital with recurrent syncopal episodes for one month. She reported sitting on the toilet to urinate and then lost consciousness upon standing up. She denied hitting her head and did not have any prodromal symptoms. She has had approximately 9 to 10 episodes previously. She also endorsed having intermittent nonproductive cough, persistent high-grade fevers of up to 106F, nausea, vomiting, and unintentional weight loss. She had a similar presentation one week ago when an extensive workup was done that only showed new cervical and axillary lymphadenopathy, for which she underwent fine-needle aspiration of one of the lymph nodes. This showed a polymorphous lymphocytic population, favoring reactive lymphadenopathy. She improved with intravenous fluids and anti-emetics and was advised to follow up with her outpatient providers.

She denied having any chills or rigors during the febrile

episodes, which occurred randomly throughout the day and without clear association with day/night cycles. She reported mild dysphagia due to thrush for which she was started on fluconazole as outpatient. Her medical history was significant for gestational diabetes mellitus. She is originally from Cape Verde and came to the United States approximately 14 years ago. She denied any traveling since then and has remained in the state of Rhode Island. She has no pets at home, nor does she have any animal exposures. She denied any tobacco use, alcohol use, or drug use. She denied any significant outdoor activities recently. She is fully vaccinated for COVID-19. She only takes fluconazole at home. Physical exam revealed two palpable nodes in the posterior neck/retrotruncal area (biggest was 2 cm, the other 1 cm). Orthostatic vital signs were negative.

A complete blood count showed leucopenia with borderline lymphopenia. A CT scan of the neck/chest/abdomen/pelvis with contrast found interval enlargement of the cervical and thoracic lymph nodes. Infectious diseases were consulted who recommended extensive workup. Liver function panel showed mild transaminitis with AST 318 IU/L (10–42 IU/L) and ALT 187 IU/L (6–45 IU/L). CRP was 16.56 mg/L (0–10mg/L) and ESR was 25mm/h (0–20mm/h). Parasite smear, hepatitis panel, treponemal antibody, anaplasma PCR, human herpesvirus-6 (HHV-6) PCR, QuantiFERON gold, Human Immunodeficiency Virus (HIV), toxoplasma, and antinuclear antibody (ANA) were negative. Cytomegalovirus (CMV) IgG was elevated, but CMV IgM was negative. Respiratory pathogen panel including SARS-CoV2 was negative. Blood cultures had no growth after five days. Epstein Barr Virus (EBV) IgG and Nuclear Ag Ab was elevated. Alpha 1 antitrypsin was mildly elevated. Echocardiogram was normal.

Surgery was consulted, who performed a left axillary lymph node biopsy. Flow cytometry revealed 64% T-cells; CD4:CD8 ratio = 2.2. The T-cell subsets showed polytypic TRBC1 expression. 32% B-cells; sIg kappa: sIg lambda light chain ratio = 1.6. Histiocytes were positive for CD4 and CD68. The cytotoxic marker Granzyme B was positive in numerous lymphocytes. The final biopsy diagnosis was KF lymphadenitis. She was discharged from the hospital with outpatient hematology follow-up. One week following discharge, she reported mild fatigue although her fevers had resolved, and lymphadenopathies improved.

DISCUSSION

Kikuchi-Fujimoto (KF) disease, also known as Kikuchi's disease, or necrotizing histiocytic lymphadenopathy, was first described in Japan by Masahiro Kikuchi and Y. Fujimoto et al. in 1972.¹ It has a worldwide distribution and affects predominantly young adults, especially young women of Asian descent. Patients most often present with cervical lymphadenopathy (most commonly in the posterior cervical triangle and the jugular carotid chain), and sometimes associated with fever and leucopenia. Affected individuals may also develop mild fever, myalgia, night sweats, and a rash. Less common symptoms include headaches, fatigue, joint pain (arthralgia), and nausea and vomiting. There is no definitive lab test for KF disease, although 25.5% to 58.3% of patients present with leucopenia².

Various infections such as toxoplasma, brucellosis, *Yersinia enterocolitica*, *Bartonella henselae*, and viruses such as herpes viruses, HIV, EBV, rubella are thought to play a role³. Some authors also hypothesize that KF disease may represent the body's autoimmune response to various stimuli, including the bacterial and viral agents listed above and dietary sources such as raw fish. There is also a possible role for interferon-gamma, interleukin-6 and apoptotic cell death in some cases⁴. KF disease has been known to have an association with the development of SLE in patients⁵. The diagnosis of SLE has been noted to occur before, during, or after the diagnosis of KF disease⁶. Electron micron studies have identified tubular reticular structures within the cytoplasm of histiocytes and lymphocytes in KF disease, that are also seen in patients with SLE and other autoimmune conditions⁷. Some investigators also believe KF disease could be an incomplete or self-limited form of an autoimmune condition due to an exuberant T cell response to a variety of non-specific stimuli.⁸ Even though the two conditions share many clinical features, they are distinct in their histopathologic criteria.⁹

Confirmatory diagnosis is through lymph node biopsy. The minimum criteria for the diagnosis of KF disease are the presence of aggregated histiocytes with occasional crescent-shaped nuclei, plasmacytoid histiocytes, and scattered karyorrhexis.¹⁰ Immunohistochemical stains typically demonstrate CD68+/CD123 plasmacytoid dendritic cells, CD68+/myeloperoxidase+ histiocytes, immunoblasts, abundance of CD8+ lymphocytes, and paucity of neutrophils.⁸ Differential diagnoses of KF disease should include malignant lymphoma, metastatic carcinoma, Systemic Lupus Erythematosus (SLE), Hemophagocytic lymphohistiocytosis (HLH), toxoplasmosis and cat-scratch disease, and HIV/AIDS. Serologic testing for SLE is advisable whenever a diagnosis of KF disease is made. Diagnosis of SLE typically requires at least 4 of the 11 criteria to be met, and lymphadenitis is not one of the criteria. Lymph node biopsies typically reveal follicular hyperplasia, plasma cells, Azzopardi phenomenon and presence of hematoxylin bodies.⁵ In our

case, ANA and anti-ds DNA were negative. HLH was considered unlikely as the patient's triglyceride and fibrinogen levels were normal. Lymphoma was considered unlikely as fine-needle aspiration of the lymph nodes did not show findings consistent with malignancy.

The prognosis is good, and in most cases, the disease resolves spontaneously within several months. Therapy is rarely indicated in KF disease, and when required, it is geared towards symptomatic management. In some cases, glucocorticoids or hydroxychloroquine have been used with some success.^{11,12} The recurrence rate can be anywhere from 4% to 15%.¹³ To our knowledge, this is the first case of KF disease presenting with recurrent fever and syncopal episodes. There was one prior case report of a patient with FUI that had KF disease associated with EBV.¹⁴ There is no known correlation between KF disease and Hodgkin's Lymphoma. In conclusion, although KF disease is benign, physicians must be aware of KF disease as early biopsy can be instrumental in reducing unnecessary investigations and in shortening the length of hospital stay. Patients with KF disease should also have continued long-term follow-up care due to their increased risk of developing SLE.

References

1. Kikuchi, M. Lymphadenitis showing focal reticulum cell hyperplasia with nuclear debris and phagocytosis. *Nippon Ketsueki Gakkai Zasshi*. 1972; 35: 379–380.
2. Kikuchi's disease (histiocytic necrotizing lymphadenitis). A clinicopathologic study of 79 cases with an analysis of histologic sub-types, immunohistology, and DNA ploidy. Kuo TT. <https://insights.ovid.com/crossref?an=00000478-199507000-00008>. *Am J Surg Pathol*. 1995;19:798–809.
3. Lerosey Y, Lecler-Scarcella V, Francois A, Guitrancourt JA. A pseudo-tumoral form of Kikuchi's disease in children: a case report and review of the literature. *Int J Pediatric Otorhinolaryngol*. 1998;45:1–6.
4. Kubota N, Tsukamoto R, Kurakawa K, Imai T, Furusho K. Elevated serum interferon gamma and interleukin-6 in patients with necrotizing lymphadenitis (Kikuchi's disease). *Br J Haematol*. 1996;95:613–615.
5. Deaver D, Horna P, Cualing H, Sokol L. Pathogenesis, diagnosis, and management of Kikuchi-Fujimoto disease. *Cancer Control*. 2014 Oct;21(4):313-21. doi: 10.1177/107327481402100407. PMID: 25310212.
6. Martínez-Vázquez C, Hughes G, Bordon J, Alonso-Alonso J, Anibarro-García A, Redondo-Martínez E, Touza-Rey F. Histiocytic necrotizing lymphadenitis, Kikuchi-Fujimoto's disease, associated with systemic lupus erythematosus. *QJM*. 1997 Aug;90(8):531-3. doi: 10.1093/qjmed/90.8.531. PMID: 9327032.
7. Dorfman RF. Histiocytic necrotizing lymphadenitis of Kikuchi and Fujimoto. *Arch Pathol Lab Med*. 1987 Nov;111(11):1026-9. PMID: 3662766.
8. Bosch X, Guilabert A, Miquel R, Campo E. Enigmatic Kikuchi-Fujimoto disease: a comprehensive review. *Am J Clin Pathol*. 2004 Jul;122(1):141-52. doi: 10.1309/YF08-1L4T-KYWV-YVPQ. PMID: 15272543.
9. Hu S, Kuo TT, Hong HS. Lupus lymphadenitis simulating Kikuchi's lymphadenitis in patients with systemic lupus erythematosus: a clinicopathological analysis of six cases and review of the literature. *Pathol Int*. 2003 Apr;53(4):221-6. doi: 10.1046/j.1320-5463.2003.01458.x. PMID: 12675765.

10. Tsang WY, Chan JK, Ng CS. Kikuchi's lymphadenitis. A morphologic analysis of 75 cases with special reference to unusual features. *Am J Surg Pathol.* 1994;18(3):219-231.
11. Yoshioka K, Miyashita T, Nakamura T, Inoue T, Yamagami K. Treatment of histiocytic necrotizing lymphadenitis (Kikuchi's disease) with prolonged fever by a single course of methylprednisolone pulse therapy without maintenance therapy: experience with 13 cases. *Intern Med.* 2010;49:2267-2270.
12. Rezai K, Kuchipudi S, Chundi V, Ariga R, Loew J, Sha BE. Kikuchi-Fujimoto disease: hydroxychloroquine as a treatment. *Clin Infect Dis.* 2004; 39:124-126.
13. Cheng CY, Sheng WH, Lo YC, et al. Clinical presentations, laboratory results and outcomes of patients with Kikuchi's disease: emphasis on the association between recurrent Kikuchi's disease and autoimmune diseases. *J Microbiol Immunol Infect.* 2010;43(5):366-371.
14. Moyer A, Hanafi MZ, Scordino T, Bronze M. Kikuchi-Fujimoto Disease: An Atypical Presentation of a Rare Disease. *Cureus.* 2019 Feb 1;11(2):e3999. doi: 10.7759/cureus.3999. PMID: 30989008; PMCID: PMC6445562.

Authors

Quynh-Lan Dao, MD, Division of Hospital Medicine, The Miriam Hospital, Providence, RI.

Bridget Hottenstein, MD, Division of Hospital Medicine, The Miriam Hospital; Warren Alpert Medical School of Brown University, Providence, RI.

Vijairam Selvaraj, MD, MPH, Division of Hospital Medicine, The Miriam Hospital; Warren Alpert Medical School of Brown University, Providence, RI.

Correspondence

Vijairam Selvaraj, MD, MPH

The Miriam Hospital

164 Summit Ave, Providence, RI, 02906

vijairam.selvaraj@lifespan.org

An Incidental Appendix Cystadenoma in a 47-Year-Old Male

WINSTON McCORMICK, BS, MD'23; YIGIT BAYKARA, MD; JESSICA YOON, MD; STEVEN SCHECHTER, MD

CASE PRESENTATION

A 47-year-old Hispanic man presented to The Miriam Hospital Emergency Department with complaint of left-sided flank pain. He had a past medical history of diabetes mellitus, hypertension, hypercholesterolemia, and syncope. Routine labs revealed an increased creatinine, and urinalysis revealed microscopic hematuria. Non-contrast CT of the abdomen and pelvis incidentally showed dilatation of the appendix to a diameter of 2.6cm with low-density fluid and mural calcifications (Figure 1A,1B). No inflammatory stranding was seen. Concern for mucinous neoplasm of the appendix was noted, and surgical consult was recommended. Upon further questioning, the patient admitted to occasional dull right lower quadrant aching that was sub-clinical in nature.

Colonoscopy was performed 11 days after initial presentation. The appendiceal orifice and ileocecal valve were normal, and the remainder of the colonoscopy was uneventful. CEA was normal. The patient underwent elective hand-assisted laparoscopic appendectomy and partial cecectomy along with the mesoappendix. There was no evidence of

metastasis or peritoneal seeding. Grossly, the appendix measured 5.8 x 3.3 x 3.2 cm and was cystically dilated with a brown smooth silvery surface (Figure 2). Pathology confirmed LAMN with margins of excision free from neoplastic cells (Figure 3). Post-surgical recovery was uncomplicated.

Figure 1B. Coronal image of the same CT demonstrating fluid-filled appendix. The average Hounsfield units of the fluid measured approximately 18-24 indicative of proteinaceous fluid.




Figure 1A. Axial image through a non-contrast CT demonstrating fluid-filled appendix (white arrows) with mural calcifications (blue arrows) in the right lower quadrant.



Figure 2. The gross resected appendix with mesoappendix demonstrating dilatation, brown discoloration and intact serosa.



Figure 3A,3B. Histopathology of low-grade appendiceal mucinous neoplasm of the appendix is shown. Cystic space filled with mucin is evident lined by mucin-producing columnar cells that have low-grade nuclear atypia, abundant apical mucin and elongated nuclei without basement membrane invasion can be seen.



DISCUSSION

Mucocele of the vermiform appendix remains a rare pathology. It underlies approximately 0.1–0.3% of appendectomies.^{2,5,8,10} It appears to have a slight predilection for biological females over the age of 50.^{2,6} Most extant literature on the appendiceal mucocele (AM) occurs as single-case studies.¹¹ In accordance with its low incidence, there is no typical presentation for AM and it is commonly diagnosed incidentally, but patients may present with symptoms mimicking acute appendicitis, such as lower right quadrant pain.^{1,7-10}

The mucocele is a non-specific cystic dilatation secondary to mucinous accumulation that may have several etiologies. There are four types of AM: mucous retention cysts, epithelial hyperplasia, cystadenoma, and cystadenocarcinoma, with the latter two diagnoses recognized as primary appendiceal mucinous neoplasms (AMNs).^{1,6,9,10} The cystadenoma is the more common AMN, and is considered benign. It is histopathologically classified as a low-grade appendiceal mucinous neoplasm (LAMN).^{6,7} The LAMN is exceedingly rare, with an estimated incidence of 1,000–2,000 cases in the United States per year.⁷

There are multiple complications that may arise from the AM, but the most feared is pseudomyxoma peritonei (PMP), which may arise from any AM subtype.⁴ PMP arises from the accumulation of mucous-producing epithelial cells in the peritoneal cavity, often from seeding of a ruptured AM or mucinous ovarian cyst, and it frequently recurs after cytoreduction surgery and has a 10-year overall survival rate of only 63%, leading it to be considered a malignant condition.^{4,5,9} Given these considerations, especially the potential for PMP and malignancy, current standard-of-care for AM is surgical resection of the appendix and mesoappendix without spillage of mucinous contents to obtain a pathologic diagnosis and grossly negative margins.⁴

Removing an intact specimen is of paramount importance, as rupture and spillage of mucin can lead to PMP and is considered a negative outcome.⁴ A thorough pre-operative work-up for underlying carcinoma must be performed due to high rates of underlying malignancies, including adnexal, urologic, and colorectal in up to 20% of cases.^{4,6,8} The long-term prognosis post-appendectomy is not clear, but for LAMNs with complete excision, appendectomy is considered curative.^{4,7} For HAMN, further treatment with right colectomy is recommended.^{4,7} Distinct biologic and clinical differences between HAMN and LAMN are not known. The difference remains a pathologic one. Given the possibility for this pathology to present identically to acute appendicitis, one of the most common surgical urgencies, yet with very different complications, the mucocele remains a rare but important clinical pathology.³

References

- Demetrashvili Z, Chkhaidze M, Khutsishvili K, et al. Mucocele of the Appendix: Case report and review of literature. *International Surgery*. 2012;97(3):266-269. doi:10.9738/cc139.1
- Dixit A. Appendiceal Mucocoeles and pseudomyxoma peritonei. *World Journal of Gastroenterology*. 2007;13(16):2381. doi:10.3748/wjg.v13.i16.2381
- Fingar K, Stocks C, Weiss A, Steiner C. Most Frequent Operating Room Procedures Performed in U.S. Hospitals, 2003-2012. Healthcare Cost and Utilization Project. <https://www.hcup-us.ahrq.gov/reports/statbriefs/sb186-Operating-Room-Procedures-United-States-2012.jsp>. Published December 2014. Accessed July 27, 2021.
- Glasgow SC, Gaertner W, Stewart D, Davids J, Alavi K, Paquette IM, Steele SR, Feingold DL. (2019). The American Society of Colon and Rectal Surgeons, clinical practice guidelines for the management of appendiceal neoplasms. *Diseases of the Colon & Rectum*, 62(12), 1425–1438. <https://doi.org/10.1097/dcr.0000000000001530>

5. Kahn M, Friedman IH. Mucocele of the appendix. *Diseases of the Colon & Rectum*. 1979;22(4):267-269. doi:10.1007/bf02586892
6. Padmanaban V, Morano WF, Gleeson E, et al. Incidentally discovered low-grade appendiceal mucinous neoplasm: A precursor to pseudomyxoma peritonei. *Clinical Case Reports*. 2016;4(12):1112-1116. doi:10.1002/ccr3.694
7. Porter KR, Ramos CE, Neychev V. Low-grade appendiceal mucinous neoplasm in the context of acute appendicitis. *Cureus*. 2019. doi:10.7759/cureus.5159
8. Roberge RJ, Park AJ. Mucocele of the appendix: An important clinical rarity. *The Journal of Emergency Medicine*. 2006;30(3):303-306. doi:10.1016/j.jemermed.2005.05.026
9. Vyas J, Badgurjar M, Saxena P, Parihar S, Thakor P. Prudent planning in management of mucocele of appendix. *International Journal of Surgery Case Reports*. 2021;81:105766. doi:10.1016/j.ijscr.2021.105766
10. Zagrodnik DF, Rose DM. Mucinous Cystadenoma of the appendix: Diagnosis, surgical management, and follow-up. *Current Surgery*. 2003;60(3):341-343. doi:10.1016/s0149-7944(02)00728-6
11. Zanati SA, Martin JA, Baker JP, Streutker CJ, Marcon NE. Colonoscopic diagnosis of mucocele of the appendix. *Gastrointestinal Endoscopy*. 2005;62(3):452-456. doi:10.1016/j.gie.2005.04.018

Authors

Winston McCormick, BS, MD'23, The Warren Alpert Medical School of Brown University, Providence, RI.
 Yigit Baykara, MD, The Warren Alpert Medical School of Brown University, Providence, RI.
 Jessica Yoon, MD, The Warren Alpert Medical School of Brown University, Providence, RI.
 Steven Schechter, MD, The Warren Alpert Medical School of Brown University, Providence, RI.

Disclosures

The authors declare no conflicts of interest.

Correspondence

Winston McCormick, BS, MD'23
 69 Arnold St. Providence RI, 02906
 Winston_mccormick@brown.edu



Adverse Childhood Experiences and Long-Term Health among Adults in Rhode Island

TRACY L. JACKSON, PhD, MPH; KARINE MONTEIRO, MPH

Adverse Childhood Experiences (ACEs) are stressful or traumatic events in childhood that can undermine a child's sense of safety and well-being.¹ Examples of ACEs include, being a victim of or witnessing abuse, living in a home with someone with a substance abuse problem, and living with someone who was incarcerated. According to national estimates from the 2010 Behavioral Risk Factor Surveillance System, more than 60% of adults have experienced at least one type of ACE.¹ Prior research has found that the toxic stress that results from ACEs not only impacts the health of children, but has long lasting effects into adulthood as well, negatively influencing factors such as employment, mental health, chronic disease, substance use, and other aspects of health.^{1,2,3} The aim of this study was to examine the prevalence and impact of ACEs among Rhode Island adults.

METHODS

Data were from the 2020 Rhode Island Behavioral Risk Factor Surveillance System (BRFSS). The BRFSS is a telephone survey of non-institutionalized adults ≥ 18 years that is administered by the RI Department of Health (RIDOH) with support from the Centers for Disease Control and Prevention (CDC) and is used to measure risk behaviors and health. Data obtained from the survey sample are weighted to obtain statewide population estimates.

The 2020 RI BRFSS contained an eleven-item module on ACEs. These items were classified into one of eight categories:

- 1) Household mental illness
 - Did you live with anyone who was depressed, mentally ill, or suicidal?
- 2) Household substance abuse
 - Did you live with anyone who was a problem drinker or alcoholic?
 - Did you live with anyone who used illegal street drugs or who abused prescription medications?
- 3) Household incarceration
 - Did you live with anyone who served time or was sentenced to serve time in a prison, jail, or other correctional facility?
- 4) Parental divorce/separation
 - Were your parents separated or divorced?

- 5) Household domestic violence
 - How often did your parents or adults in your home ever slap, hit, kick, punch or beat each other up?
- 6) Emotional abuse
 - How often did a parent or adult in your home ever swear at you, insult you, or put you down?
- 7) Physical abuse
 - Not including spanking, how often did a parent or adult in your home ever hit, beat, kick, or physically hurt you in any way?
- 8) Sexual abuse
 - How often did anyone at least 5 years older than you or an adult, ever touch you sexually?
 - How often did anyone at least 5 years older than you or an adult, try to make you touch them sexually?
 - How often did anyone at least 5 years older than you or an adult, force you to have sex?

Similar to methods of previous studies from Kaiser-Permanente² and the Colorado Department of Public Health,³ responses were summarized into an overall ACE score where respondents received one point for each type of ACE they experienced prior to the age of 18. Those missing responses to any ACE question were excluded from the total score calculations. Possible scores ranged from 0–8. Scores were then categorized into 3 groups: No ACEs, Low ACEs (1–3 ACEs), and High ACEs (4 or more ACEs).

Descriptive analyses were conducted to examine the distribution of ACE scores in the population and the prevalence of each ACE type. Next, we examined the demographic characteristics and prevalence of selected health outcomes and behaviors by the three-level ACE measure. All health outcomes were then separately assessed in multivariable logistic regression models examining the association between the three-level ACE measure and health, adjusting for sex, sexual orientation, age group, race/ethnicity, and educational attainment.

RESULTS

Overall 5,378 respondents completed the 2020 BRFSS, representing a weighted sample of 857,269 adults. The most common type of ACE was emotional abuse (34.7%), followed by having divorced or separated parents (30.7%), household substance use (26.1%), and physical abuse (24.7%) (Figure 1).

Figure 1. Prevalence of ACEs, by type





Figure 2. Distribution of ACE Scores among Rhode Island Adults



ACE scores ranged from 0-8, with a mean of 1.63. Overall 63.2% reported at least one type of ACE with 24.0% reporting only one ACE and 1% reporting all 8 ACEs (Figure 2).

When categorized into the three-level ACE measure, 36.8% had 0 ACEs, 47.1% had Low ACEs, and 16.1% had High ACEs (Table 1). Younger age groups, gay/lesbian/bisexual/others, and those of lower socioeconomic status (SES) were more likely to have higher ACE scores than comparison groups.

Descriptive analysis revealed significant differences in prevalence of health outcomes by ACE level for all health measures assessed except diabetes, kidney disease, and cancer. For example, 45.1% of those with a high ACEs score reported a history of depression, compared to 24.3% of those with low ACE scores, and 8.6% of those with no ACEs (Table 2). Nearly one-third (30.0%) of those with a high ACE score were cigarette smokers compared to 13.4% of those with a low ACE score, and 6.7% of those with no ACEs.

In multivariable models adjusting for age, race/ethnicity, sex, sexual orientation, and educational attainment individuals with High ACE scores were significantly more likely than those with no ACEs to experience all health and behavioral outcomes assessed except obesity (Table 3). Notably, individuals with a high ACE score had more than seven times the odds of having a depression diagnosis (Adjusted

Table 1. ACE scores in Rhode Island, by selected demographics

	No ACEs N=240,345; 36.8%		Low ACEs N=306,956; 47.1%		High ACEs 104,946; 16.1%	
	N	%	N	%	N	%
Sex						
Male	119,446	38.6	142,072	45.9	47,719	15.4
Female	120,898	35.2	164,884	48.1	57,227	16.7
Sexual orientation						
Lesbian/gay/bisexual/other	7,338	20.9	12,539	35.8	15,186	43.3
Straight	217,463	37.3	281,388	48.2	84,938	14.5
Age Group						
18-29 years	39,742	30.5	66,574	51.2	23,807	18.3
30-44 years	49,679	32.3	69,326	45.1	34,859	22.7
45-64 years	74,389	34.9	103,531	48.6	35,052	16.5
65+ years	70,979	48.2	65,782	4.7	10,440	7.1
Race/ethnicity						
White, non-Hispanic	182,987	38.1	221,601	46.3	74,463	15.5
Black, non-Hispanic	10,213	36.6	13,741	49.3	3,945	14.1
Other, non-Hispanic	12,134	31.8	18,452	48.0	7,827	20.4
Hispanic	29,926	32.5	45,903	49.9	16,191	17.6
Household income						
Less than \$25,000	34,919	31.4	51,925	46.7	24,263	21.8
\$25,000-49,999	37,802	32.4	57,018	48.9	21,717	18.6
\$50,000-74,999	33,394	37.0	39,477	43.7	17,439	19.3
\$75,000+	85,589	38.6	107,205	48.4	28,921	13.0
Education level						
Less than high school	21,964	32.0	32,086	46.7	14,685	21.4
High school	64,655	34.8	86,177	46.4	34,705	18.7
More than high school	152,497	38.5	188,110	47.5	55,019	13.9

Odds Ratio [aOR]=7.37, 95% Confidence Interval [CI]:5.09-10.69) and six times the odds of experiencing frequent mental distress (FMD; aOR=6.08, 95% CI: 3.88-9.52) compared to those with no ACEs. Those with a low ACE score were significantly more likely than those with no ACEs to experience a number of health outcomes including fair or poor health, FMD, frequent physical distress, arthritis, COPD, depression, disability, cigarette smoking, and heavy drinking.

DISCUSSION

A majority of Rhode Island adults (63.2%) have experienced at least one type of ACE and approximately 1 in 6 reported four or more ACEs. There was a dose response between ACEs and poor health outcomes as those who experienced greater numbers of ACEs were more likely to engage in risk

Table 2. Prevalence of health and behavioral outcomes, by ACE level

	No ACEs		Low ACE Score		High ACE Score	
	%	95% Confidence Interval	%	95% Confidence Interval	%	95% Confidence Interval
General Health						
Fair or Poor Health*	8.2	6.5–9.9	12.0	10.1–14.0	20.9	16.1–25.7
Frequent ^a mental distress*	5.8	4.3–7.4	13.5	10.8–16.3	32.9	26.9–38.9
Frequent ^a physical distress*	6.9	5.1–8.6	10.0	8.0–12.1	17.5	13.2–21.9
Frequent ^a activity limitations*	10.2	6.8–13.6	14.7	12.2–18.1	28.5	22.0–35.0
Chronic Conditions^b						
Current asthma*	9.2	7.0–11.3	11.1	8.9–13.2	24.9	19.4–30.4
Arthritis*	23.8	21.1–26.4	23.9	21.4–26.4	32.4	26.9–38.0
Cancer	13.5	11.5–15.4	12.5	10.7–14.2	11.4	7.9–15.0
Cardiovascular disease ^{c*}	7.8	6.3–9.3	7.8	6.4–9.2	8.1	5.4–10.7
COPD*	4.7	3.5–5.8	7.0	5.5–8.4	12.2	8.0–16.3
Depression*	8.6	6.7–10.5	23.7	20.7–26.8	45.1	39.1–51.1
Diabetes	11.2	9.2–13.2	9.8	8.1–11.5	12.2	8.7–15.7
Disability ^{d*}	17.3	14.9–19.7	24.3	21.5–27.1	42.2	36.2–38.2
Kidney disease	2.3	1.3–3.3	2.5	1.6–3.5	3.5	0.8–6.2
Health risk factors/behaviors						
Cigarette smoking*	6.7	5.1–8.4	13.4	11.1–15.8	30.0	24.2–35.8
Binge drinking ^{e*}	11.0	8.5–13.5	14.9	12.3–17.4	22.8	17.4–28.2
Heavy drinking ^{f*}	2.7	1.8–3.6	6.6	5.1–8.2	10.0	6.7–13.3
Obesity ^{g*}	30.1	26.3–33.8	30.6	27.5–33.8	38.3	32.3–44.2

Notes: *indicates statistical significance ($p < .05$); Frequent defined as ≥ 14 days in the last 30; ^aindicates ever diagnosed with condition unless otherwise specified; ^cCOPD=Chronic Obstructive Pulmonary Disorder; ^dproblems with hearing, vision, cognition, mobility, self-care, or independent living; ^e ≥ 4 drinks for females or ≥ 5 drinks for males, in one sitting in the last 30 days; ^f ≥ 8 more drinks for females or ≥ 15 for males, per week in the last 30 days; ^gBody Mass Index ≥ 30 based on height and weight.

behaviors and have health problems, findings similar to studies conducted among other populations.^{2,3,4}

ACEs can be prevented. Prior research has identified a number of family and neighborhood level risk factors for ACEs including low SES, isolation from friends/family, and living in communities with high rates of violence, and lack of activities for young people.¹ The CDC recommends six main evidence-based strategies to reduce the risk of ACEs. These strategies include: strengthening economic support for families (e.g., improving financial security, family-friendly work policies), promoting social norms that protect against violence (e.g., public education campaigns, legislation to reduce corporal punishment), ensuring strong

Table 3. Adjusted odd ratio of selected health outcomes, comparing low ACE scores and high ACE scores to those with no ACEs

	Low ACE Score		High ACE Score	
	Adjusted Odds ratio	95% Confidence Interval	Adjusted Odds ratio	95% Confidence Interval
General Health				
Fair or Poor Health	1.67	1.22–2.29	3.55	2.41–5.20
Frequent mental distress	2.28	1.55–3.37	6.08	3.88–9.52
Frequent physical distress	1.53	1.05–2.23	2.77	1.76–4.38
Frequent activity limitations	1.50	0.91–2.50	2.92	1.73–4.93
Chronic Conditions				
Asthma	1.17	0.82–1.68	2.89	1.86–4.51
Arthritis	1.31	1.06–1.62	3.47	2.44–4.93
Cancer	1.25	0.98–1.60	1.74	1.14–2.66
Cardiovascular disease	1.27	0.94–1.71	2.07	1.33–3.25
COPD	1.80	1.24–2.61	4.78	2.88–7.93
Depression	3.14	2.33–4.23	7.37	5.09–10.69
Diabetes	0.97	0.72–1.30	1.70	1.10–2.62
Disability	1.71	1.34–2.17	4.35	3.12–6.08
Kidney disease	1.34	0.71–2.54	2.37	1.06–5.31
Health risk factors/behaviors				
Cigarette smoking	2.21	1.55–3.15	4.97	3.28–7.53
Binge drinking	1.35	0.96–1.90	2.19	1.43–3.36
Heavy drinking	2.40	1.56–3.69	3.72	2.16–6.38
Obesity	1.02	0.79–1.31	1.38	0.98–1.94

starts for children (e.g., home visitation programs, high-quality childcare), teaching coping and relationship skills (e.g., social-emotional learning programs, parenting skills), connecting youth to caring adults and activities (e.g., mentoring and after school programs), and intervening to lessen harms (e.g., enhanced screening in primary care, access to mental health and substance use treatment).⁵

This study had several limitations. First, data were self-reported and may be prone to recall or reporting bias, particularly the underreporting of ACEs. Additionally, our definition of ACEs was limited to the 11 measures assessed in the module – there may be other adverse experiences not measured in our study – and there may be different aspects of ACEs, such as severity and age of onset – that can have different implications on health. Also, the relationship between ACEs and SES and health is nuanced and was difficult to fully assess in our study. We controlled for adult SES in our models due to the strong association between

childhood SES and ACEs and childhood SES and adult SES, but prior research has shown ACEs can directly impact adult SES – a factor which, in turn, can impact health. We did not have measures of childhood SES so we could not fully disentangle the relationship between these factors in our study.

Overall results of this study indicate a strong association between ACEs and poor health outcomes and risky health behaviors among adults in Rhode Island. While more research is needed to fully understand the intricacy of the relationship, health professionals and community advocates should incorporate evidence-based strategies to reduce ACEs and mitigate their harmful effects.

References

1. Centers for Disease Control and Prevention. Preventing Adverse Childhood Experiences. 2021. <https://www.cdc.gov/violenceprevention/aces/fastfact.html>
2. Felitti VJ, Anda RF, Nordenberg D, Williamson DF, Spitz AM, Edwards V, Koss MP, Marks JS. Relationship of childhood abuse and household dysfunction to many of the leading causes of death in adults: The Adverse Childhood Experiences (ACE) Study. *American Journal of Preventive Medicine*. 1998;14:245–258.
3. Anderson Mellies A. Impact of adverse childhood experiences on adult health in Colorado. Colorado Department of Public Health and Environment, 2016.
4. Merrick MT, Ford DC, Ports KA, et al. Vital Signs: Estimated Proportion of Adult Health Problems Attributable to Adverse Childhood Experiences and Implications for Prevention – 25 States, 2015–2017. *MMWR Morb Mortal Wkly Rep* 2019;68:999–1005. DOI: <http://dx.doi.org/10.15585/mmwr.mm6844e1>
5. Centers for Disease Control and Prevention. Preventing Adverse Childhood Experiences: Leveraging the Best Available Evidence. Atlanta, GA: National Center for Injury Prevention and Control, Centers for Disease Control and Prevention, 2019.

Authors

Tracy L. Jackson, PhD, MPH, is a Senior Public Health Epidemiologist in the Center of Health Data and Analysis (CHDA) at the Rhode Island Department of Health (RIDOH).

Karine Monteiro, MPH, is the Acting Manager of Health Surveys in CHDA at RIDOH.

Correspondence

Tracy L. Jackson, PhD, MPH
Center for Health Data and Analysis
Rhode Island Department of Health
tracy.jackson@health.ri.gov



VITAL STATISTICS

NICOLE E. ALEXANDER-SCOTT, MD, MPH
 DIRECTOR, RHODE ISLAND DEPARTMENT OF HEALTH
 COMPILED BY ROSEANN GIORGIANNI, DEPUTY STATE REGISTRAR

Rhode Island Monthly Vital Statistics Report

Provisional Occurrence Data from the Division of Vital Records

VITAL EVENTS	REPORTING PERIOD		
	MAY 2021	12 MONTHS ENDING WITH MAY 2021	
	Number	Number	Rates
Live Births	983	10,854	10.2*
Deaths	921	11,829	11.2*
Infant Deaths	5	53	4.9#
Neonatal Deaths	4	37	3.4#
Marriages	561	5,186	4.9*
Divorces	207	2,249	2.1*

* Rates per 1,000 estimated population
 # Rates per 1,000 live births

Underlying Cause of Death Category	REPORTING PERIOD			
	NOVEMBER 2020	12 MONTHS ENDING WITH NOVEMBER 2020		
	Number (a)	Number (a)	Rates (b)	YPLL (c)
Diseases of the Heart	201	2,397	226.3	3,489.5
Malignant Neoplasms	170	2,139	210.9	4,400.0
Cerebrovascular Disease	32	436	41.2	625.0
Injuries (Accident/Suicide/Homicide)	72	960	90.6	14,104.5
COPD	30	473	45.5	550.0

(a) Cause of death statistics were derived from the underlying cause of death reported by physicians on death certificates.
 (b) Rates per 100,000 estimated population of 1,059,361 for 2019 (www.census.gov)
 (c) Years of Potential Life Lost (YPLL).

NOTE: Totals represent vital events, which occurred in Rhode Island for the reporting periods listed above.
 Monthly provisional totals should be analyzed with caution because the numbers may be small and subject to seasonal variation.



Rhode Island's Medical Staffing Experts

Favorite Healthcare Staffing provides a comprehensive range of staffing services at preferred pricing to RIMS members. Call today to see why we are the favorite choice of healthcare professionals and physician practices across the US!

 401.354.7115

 MedicalStaffing@FavoriteStaffing.com



Favorite Healthcare Staffing is a Valued
Sponsor of the Rhode Island Medical Society

My Newest Favorite Hallucination

JOSEPH H. FRIEDMAN, MD

ABOUT ONE-THIRD OF PEOPLE TREATED with medications for Parkinson's disease have visual hallucinations. I learned early in my career to ask about them because patients were often reluctant to mention them for fear of being labelled as "crazy." I also realized in asking about hallucinations early on, which generally brings a smile to most patients' faces, that if they eventually do develop, the patient and family are not panicked. They think, "So that's what Dr. Friedman was asking about...They're not so bad." It's important to know about hallucinations because the patients and their families need reassurance that they are not, in fact, "going crazy," that it's a side effect of the medications they take for mobility, and their presence alters how we treat their disease.


Early on in the disease, the hallucinations, which may occur in any sensory modality, are usually visual, unlike hallucinations in schizophrenia and other primary psychiatric disorders, which are generally only auditory. More importantly, the hallucinations in PD generally have no emotional significance. An elderly woman saw her deceased husband at night. When asked how this made her feel, responded, "How should it make me feel? He's dead." Unlike schizophrenia, in which the voices criticize the patient, telling them unpleasant things, "Can't you see that everyone hates you?...You're rotting on the inside....You have cancer and

are dying." PD patients see ordinary people, or people in clown costumes, children, pet dogs or cats and even statues. Even when the people are odd appearing, as without faces, or Lilliputian in size, possibly living in a houseplant, they are usually not disturbing. Sometimes they are entertaining. Unfortunately, sometimes they are not, but that is not for this short essay.

I have a store of hallucination stories. Most are fairly mundane, a stranger reading a book, three children playing cards, two girls dressed in tutus dancing ballet in the driveway, a dog that is sometimes an old pet that died, sometimes another dog or cat, people climbing trees in the yard. I once had a patient who took photos of the hallucinations to convince me they were, in fact, real. He saw them in his photos, too, but no one else did. Until this week I had a favorite hallucination that I heard twenty years ago. "Do you see things that aren't really there?" "Yes." "What do you see?" "I see people outside my house." "If they're outside, how do you know they're not really there?" "Well, every morning when I sit down to eat breakfast, there's a group of nuns, wearing habits, who come over to my house and start building me a deck."

"Yup. Sounds like an hallucination."

Even though most PD patients aren't bothered by their hallucinations, they generally do not like the idea that their mind is playing tricks on them, and they may worry that they mistake hallucinations for reality. They undermine one's ability to rely on the perception of reality. Occasionally patients do like the hallucinations and prefer to keep them. My patient liked seeing the nuns visiting each day. It was an entertainment. One elderly man liked seeing the baby in his room that appeared every night. "He's so good. He never makes any noise," he told me with a smile.



Chromolithographic print from 1905 titled "Auld Lang Syne."

[COURTESY OF LIBRARY OF CONGRESS PRINTS AND PHOTOGRAPHS DIVISION. [HTTPS://HDL.LOC.GOV/LOC.PNP/PPMSCA.46118](https://hdl.loc.gov/loc.pnp/ppmsca.46118)]

I guess it's like having a short, reliably entertaining LSD trip, or a very realistic daydream. Since most hallucinations occur in low-stimulus environments, like watching TV alone in a dimly lit room, a patient can usually make the hallucination disappear by turning on the lights or getting up and moving around. They always disappear when the patient approaches the hallucination and tries to touch it.

My newest favorite was described by a 70-year-old man whose hallucinations had been dramatically reduced by an adjustment of his medications. These had been bothersome mostly because they were so often present, rather than that the individual scenes and hallucinated characters were unpleasant. He now had the hallucinations only in the evening and he looks forward to them. Although it really wasn't any of my business, I asked him what he saw that was so pleasant. "Sometimes, I see my daughter and my granddaughter, both about 7 years old, playing together."

Unfortunately, patients don't get to choose their hallucinations. One sees nuns building a deck and another sees a nun in a habit with blood dripping from her mouth. One patient, who loves cats, enjoys the two cats that play by her chair and rub against her feet. She feels them but she can't pick them up because she learned from experience that they'd disappear. Unfortunately, she does not care for the man who appears on the other side of her visual field from where the cats are, who has no face. The absence of a face is not the real problem for her.

That doesn't bother her. It's simply that a strange person who doesn't belong there keeps her from enjoying the cats. "And you don't have to put out any food or clean up after them," she joked, knowing that they were hallucinations.

When I chose to train in neurology, I really had no particular interest in behavior, per se. I was interested in language dysfunction, but, with further experience, and, perhaps, lack of any faculty with a similar interest, I was attracted to my current field of movement disorders, particularly drug-induced disorders like parkinsonism and tardive dyskinesia. But, as I saw an increasing number of people with Parkinson's disease, I started to discover all sorts of fascinating, important and previously overlooked behavior problems that were either intrinsic to the disorder, or iatrogenic. I am unsure if my interest in hallucinations and delusions in PD, both unfortunately common iatrogenic problems, preceded or followed my learning from a drug rep that, "You write more prescriptions for L-Dopa than anyone in New England." Since L-Dopa is only used to treat PD, it meant that I treated more PD patients than anyone in New England. That, however, meant that I caused more psychosis than any other doctor in New England.

There is a sort of happy ending to this story, though. Doctors and patients are much more aware of these problems, and we now have some effective drugs for treating them. And the hallucinations aren't all bad. ❖

Author

Joseph H. Friedman, MD, is Editor-in-Chief Emeritus of the *Rhode Island Medical Journal*, Professor and the former Chief of the Division of Movement Disorders, Department of Neurology at the Alpert Medical School of Brown University, chief of Butler Hospital's Movement Disorders Program and first recipient of the Stanley Aronson Chair in Neurodegenerative Disorders.

Correspondence

joseph_friedman@brown.edu

Q&A with Providence College Authors of 'Today's Health Care Issues'

MARY KORR
RIMJ MANAGING EDITOR

PROVIDENCE – Providence College (PC) Professors **ROBERT B. HACKEY, PhD**, and **TODD M. OLSZEWSKI, PhD**, along with a dozen student co-authors, recently published a book on health care issues and disputed policies presented through a Democratic/Republican lens. "Today's Health Care Issues" is part of a reference series, "Across the Aisles," by publisher ABC-CLIO, LLC.

Some of the issues examined are the COVID-19 pandemic, guns and public health, health insurance marketplaces, immigrant health coverage, marijuana, Medicaid expansion, Medicare drug pricing, physician-assisted death, vaccine mandates, and veterans' health care. Chapters offer readers a historical background but focus on more contemporary health policy debates. In each chapter, the authors present:

- An "At a Glance" introduction to the topic
- A bulleted list of key positions/themes for each party
- A detailed overview of the issue that includes key milestones and decision points
- Democratic and Republican perspectives on the issue featuring key leaders and spokespersons
- Sidebars that capture policymaker positions for/against the issue

RIMJ invited the authors to further elaborate on the book and the role PC students engaged in, both in the writing and in coursework.

Q. What was the genesis for the book and author collaborations?

A. PROF HACKEY: I was approached by an editor at ABC-CLIO in 2018 about contributing a volume on health care policy to a new reference series that would span nearly 20 different policy issues. I knew I couldn't tackle such an expansive project on my own. I invited my good friend and colleague Todd Olszewski, who previously collaborated with me on evaluating health care reform in New Hampshire for the ACA Implementation Research Network, to co-author the volume. Todd's background as a historian of medicine complements my own work on federal and state health care reform, public opinion, and rhetoric. Together, we bring both



August 2021, 380pp, 7x10, eBook/Hardcover
Print: 9781440869150
ABCCLIO, LLC | 147 Castilian Dr | Santa Barbara,
CA 931175515 | USA

Excerpt

Vaccine Mandates

Vaccination represents a fundamental achievement in medicine and public health. While Democrats and Republicans have generally agreed about the effectiveness of vaccines as public health measures, partisan controversy has focused on government-imposed vaccine mandates and what exemptions, if any, should be permitted. In 2020, all 50 states and Washington, DC, had vaccination requirements for schoolchildren, and they all provided for medical exemptions to these state mandates. The District of Columbia and 45 states allow religious exemptions, and 15 states allow philosophical exemptions; however, 5 states – California, Maine, Mississippi, New York, and West Virginia – do not allow nonmedical exemptions to their state vaccination mandates.

...This philosophical divide has precedents dating to the 19th century. During that time, many states enacted compulsory smallpox vaccination laws for children and adults. Penalties for refusing immunizations against smallpox included exclusion from school systems for children and substantial fines or quarantine for adults (Colgrove 2016). The 1905 U.S. Supreme Court case *Jacobson v. Massachusetts* upheld the right of state governments to enact compulsory vaccine mandates. The court's decision reflected the widely held belief that preventing outbreaks of vaccine-preventable diseases that threaten public health warranted the forfeiture of individual liberty (Gostin 2005).

...The use of coercion to ensure vaccine compliance raises important questions about individual liberty, freedom of choice, and government regulation in the name of public health. By 2019, all states mandated immunization for enrollment in public schools, and all states also allowed medical exemptions to those mandates. However, states vary with respect to which vaccines are mandated. For example, Connecticut, New Jersey, Ohio, and Rhode Island have mandated the influenza vaccine for children in child-care facilities, but Rhode Island and Virginia (and the District of Columbia) are the only states that have mandated the human papillomavirus (HPV) vaccine for secondary school students.



Robert B. Hackey, PhD



Todd M. Olszewski, PhD

historical and contemporary perspectives to each of the chapters. Since this was an ambitious project with a tight timeline, we recruited more than a dozen current Providence College students to join us as co-authors for various chapters. Our students were exceptional colleagues who became our eyes and ears as we grappled with each policy issue.

Q. This book presents, similar to a Point /Counterpoint, Democratic and Republican views on contemporary health/policy positions, but it is not a partisan presentation. Why not Progressive, Independent, right /left positions, given the attention they are receiving on so many of these issues today?

A. PROF. HACKEY: The series presents readers with past and present Democratic and Republican positions and records on a wide range of issues. One of the primary goals of the series is improve the knowledge and ‘policy literacy’ of the public about contested issues. Our volume is written on two distinct levels – we present each topic in an accessible manner for lay readers, but each chapter has a strong evidence-base that draws upon peer-reviewed sources, public opinion polls, and public statements by policymakers and key stakeholders that will be of interest to academics and policymakers.

We provide context by reviewing the policy history of each issue, but we don’t take sides or endorse particular policy

viewpoints. Instead, we allow each party’s position to speak for itself. Our approach was guided by the series’ editorial mandate, which focused on party differences, rather than ideology. We discuss ideological differences between – and in some cases within – parties, but our goal is to highlight how each party

frames debate over these issues. The ultimate goal of our book, and the series itself, is to improve the way we talk about contested issues by helping individuals – regardless of their partisan views – to better understand where each party stands and the origins of these beliefs.

Q. I found the vaccine mandate chapter (See Book Excerpt) especially relevant given its focus today with the COVID-19 vaccination mandates. Do you see this and other historical parallels fundamental to framing policy discussions?

A. PROF. OLSZEWSKI: The vaccine mandate chapter became increasingly relevant as the COVID-19 pandemic progressed. My co-author Amanda McGrath and I started working on that chapter in January 2019 – more than a year before the onset of the COVID-19 pandemic in the United States. In our research, we documented a long-term but growing ideological divide among Republicans and Democrats regarding vaccine mandates in general and nonmedical exemptions in particular. These debates were far from new, as policymakers had already debated whether vaccine mandates represented necessary public health measures or unfair constraints on individual liberty and religious freedom. To a certain degree, some of the historical precedents included in the chapter explain how and why COVID-19 vaccine mandates have become so politically polarizing.

Q. Who is the target readership for this book?

A. PROF. HACKEY: Books in this series are intended for patrons of high school, public, and college and university libraries. We target a general audience and provide context and background for readers in each chapter. A secondary audience includes undergraduate and graduate courses on health care reform, health policy, and public health.

We’re using selected chapters from the book in a class we’re teaching together this fall. Students really embraced the volume’s approach of presenting both parties’ position in an even-handed way, without pushing a particular point of view. After reading each chapter, we ask them to wrestle with their own views on the subject so that they can decide what they think and explain their views to others. For us, that’s the essence of what we hope the book will accomplish – creating more informed and thoughtful citizens who are ready to engage in spirited public debates about important issues is what our country needs at this moment.

Q. What was the most surprising policy issue that the authors uncovered in this research that they were not fully aware of before?

A. PROF. OLSZEWSKI: Bob and I teach in the Department of Health Policy and Management at Providence College. Even though our professional training differs (Bob is a political scientist and I am an historian of medicine), we have both taught and written about historical and recent health reform efforts. I appreciated the opportunity to explore policy topics that I might not have written about otherwise. Most illuminating for me, though, was observing the inconsistent ways in which both political parties apply their respective ideological values to health policymaking.

Q. An exciting project with many student collaborators. Was this a funded project and how long did it take to complete?

A. PROF. OLSZEWSKI: We started this project in June 2018 and reviewed final page proofs in July 2021. We were fortunate to receive financial support from several sources. Providence College's Center for Engaged Learning subsidized a series of team dinners to build community among our student collaborators. PC's School of Professional Studies awarded us a summer grant to hire two students as editorial assistants; these two collaborators provided crucial support in 2020 by identifying necessary updates to chapters that we first wrote in 2019. We envisioned our student collaborators as junior scholars rather than traditional

research assistants, which made the writing experience more rewarding. Students' responsibilities were comparable to ours when preparing each chapter draft; they offered feedback on our writing and we offered feedback on theirs. By promoting trust and transparency within each chapter team, we reinforced the collaborative nature of our project and assigned equal credit for each co-authored chapter.

Q. What advice do you give today to your students who wish to pursue careers in public health and policymaking?

A. PROF. OLSZEWSKI: We have had the opportunity to co-teach our department's intensive writing seminar while working on the book. We organized our edition of the course around several chapter topics;

students write op-eds, engage in in-class debates, and write traditional research papers that expand upon their op-eds. We encourage the students to "think on their feet" and carefully consider issues from more than one point of view. By encouraging students to question everything, they learn how to formulate their own position on a given policy topic and to be able to defend that position by drawing upon multiple forms of evidence and data. Ultimately, we hope that the course prepares students to become thoughtful and principled future professionals whether they pursue careers in health care, public health, or health policy. ❖



Adventures

Aetna® is proud to support the members of
the Rhode Island Medical Society.

[Aetna.com](https://www.aetna.com)

 aetna®



Working for You: RIMS advocacy activities

October 5, Tuesday

RIMS Physician Health Committee (PHC): **Herbert Rakatansky, MD**, Chair
Harm Reduction Centers (HRC) Advisory Committee: **Elizabeth Samuels, MD**, **Rahul Vanjani, MD**

October 7, Thursday

COBRE (Center of Biological Research Excellence) Lecture Series: Compensating for Systems Failure: The Transformation of the Community Health Worker/Peer Recovery Specialist Roles during COVID-19

October 8, Friday

Virtual meeting with Doctors for Cannabis Regulation
Meeting with Senator Whitehouse staff regarding Harm Reduction Centers and US Department of Justice

October 12, Tuesday

AMA Advocacy Resource Center (ARC) Executive Committee call regarding state legislation
Harm Reduction Centers (HRC) Advisory Committee: **Elizabeth Samuels, MD**, **Rahul Vanjani, MD**
American Medical Association (AMA) Federal Advocacy Update Conference call

October 13, Wednesday

RI Department of Health (RIDOH) Board of Medical Licensure and Discipline
Governor's Overdose Intervention and Prevention Task Force: **Sarah Fessler, MD**, RIMS Past President

October 14, Thursday

RIMS' hosted Zoom Meeting regarding Harm Reduction Centers with community proponents

October 15, Friday

Care Transformation Collaborative (CTC-RI) Clinical Strategy Committee Meeting: **Elizabeth Lange, MD**, President; **Thomas Bledsoe, MD**, President-elect

October 18, Monday

Meeting with Blue Cross & Blue Shield of Rhode Island (BCBSRI): **Elizabeth Lange, MD**, President; **Thomas Bledsoe, MD**, President-elect
RIMPAC (Rhode Island Medical Political Action Committee) sponsored fundraiser for Speaker of the RI House of Representatives, K. Joseph Shekarchi

October 19, Tuesday

Office of the Health Insurance Commissioner (OHIC) Health Insurance Advisory Council (HIAC): **Catherine A. Cummings, MD**, Past President
RIDOH Harm Reduction Center Advisory Committee: **Elizabeth Samuels, MD**, **Rahul Vanjani, MD**

October 20, Wednesday

RIDOH Primary Care Physicians Advisory Committee (PCPAC): **Elizabeth Lange, MD**, President

October 21, Thursday

RIMS' hosted Zoom Meeting regarding Harm Reduction Centers with community proponents

October 26, Tuesday

The Womxn Project's Campaign for equal abortion coverage
RIDOH Harm Reduction Center Advisory Committee: **Elizabeth Samuels, MD**, **Rahul Vanjani, MD**
Medicine Grand Rounds, "The Rhode Island Medical Society: Aspects of Advocacy for Professional Satisfaction and Personal Well-being," **Elizabeth Lange, MD**, President; **Herbert Rakatansky, MD**, Physician Health Program (PHP) Chair; **Kathleen Boyd**, Director of RIMS PHP; **Newell Warde, PhD**, RIMS Executive Director

October 27, Wednesday

RIMS Finance Committee: **Kwame Dapaah-Afriyie, MD, MBA**, Chair

October 28, Thursday

Governor's Overdose Task Force Racial Equity Work Group
AMA Physician Grassroot webinar: Advocacy in a Post-COVID World: **Michael Migliori, MD**, Chair, Public Laws Committee
RIMS' hosted Zoom Meeting regarding Harm Reduction Centers with community proponents

RIMS NOTES: News You Can Use

Our biweekly e-newsletter is published on alternate Fridays exclusively for RIMS members. Contact Dulce Cosme if you've missed an issue, dcosme@rimed.org.





RIMS CORPORATE AFFILIATES

The Rhode Island Medical Society continues to drive forward into the future with the implementation of various new programs. As such, RIMS is expanding its Affinity Program to allow for more of our colleagues in health-care and related business to work with our membership. RIMS thanks these participants for their support of our membership.

Contact Marc Bialek for more information: 401-331-3207 or mbialek@rimed.org



www.nhpri.org

Neighborhood Health Plan of Rhode Island is a non-profit HMO founded in 1993 in partnership with Rhode Island's Community Health Centers. Serving over 185,000 members, Neighborhood has doubled in membership, revenue and staff since November 2013. In January 2014, Neighborhood extended its service, benefits and value through the HealthSource RI health insurance exchange, serving 49% the RI exchange market. Neighborhood has been rated by National Committee for Quality Assurance (NCQA) as one of the Top 10 Medicaid health plans in America, every year since ratings began twelve years ago.



www.ripccp.com

RIPCPC is an independent practice association (IPA) of primary care physicians located throughout the state of Rhode Island. The IPA, originally formed in 1994, represent 150 physicians from Family Practice, Internal Medicine and Pediatrics. RIPCPC also has an affiliation with over 200 specialty-care member physicians. Our PCP's act as primary care providers for over 340,000 patients throughout the state of Rhode Island. The IPA was formed to provide a venue for the smaller independent practices to work together with the ultimate goal of improving quality of care for our patients.



RIMS gratefully acknowledges the practices who participate in our discounted Group Membership Program



east bay community action program
THE BRIDGE TO SELF-RELIANCE



Diabetes & Endocrinology
Associates, Inc.

Orthopaedic Associates, Inc.



Ortho Rhode Island



HopeHealth Hulitar Hospice Center completes expansion

PROVIDENCE – Rhode Island’s only free-standing hospice center has increased its capacity by 25% with the completion of a yearlong construction project to address the region’s growing demand for specialized end-of-life care.

The HopeHealth Hulitar Hospice Center is now a 30-bed inpatient unit after finishing the approximately \$3 million expansion project in July. The project added six beds to the original 24-bed unit, located on the second and third floors of HopeHealth’s headquarters at 1085 North Main St., Providence. The project maintained the building’s 50,000 square foot footprint.

“This much-needed expansion meets an important community need for additional capacity,” said HopeHealth President & CEO **DIANA FRANCHITTO**. “We were also able to refurbish existing patient rooms after a decade of high utilization.”

Most hospice patients prefer to spend their last days at home or in a home-like setting with loved ones by their side. The Hulitar Hospice Center provides a home-like environment for hospice patients who require acute inpatient care, as their symptoms progress and cannot be safely and comfortably managed at home.

“The focus of the Hulitar Hospice Center is not just the patient but the family,” said HopeHealth Chief Medical Officer **EDWARD MARTIN, MD, MPH**.



Exterior of the Center located at 1085 North Main St. in Providence.

“It’s a calm and peaceful environment. Sometimes families are concerned about moving their loved one, but universally, once they get here, they’re so glad they did. It makes such a big difference.”

The project reconfigured the layout of the inpatient unit so that there are now 15 rooms each on the second and third floors. New flooring, paint, and carpeting refreshed all rooms, staff and family gathering areas. The second floor has a renovated interfaith chapel. The third floor

now has a relocated nurses station, renovated kitchen facilities and a dining area, and larger areas for staff, as well as new rooms where grief counselors can meet with families to provide support.

The HopeHealth Hulitar Hospice Center remained operational throughout the construction, which coincided with the COVID-19 pandemic. The center also continued to allow visitors during the pandemic so they could be with their loved one in their last days. ❖



The Four Seasons room.



Third-floor nurses station.

Federal delegation announces \$6.1M for capital improvements at RI Community Health Centers

WASHINGTON, D.C. – U.S. Senators **JACK REED** and **SHELDON WHITEHOUSE** and Congressmen **JIM LANGEVIN** and **DAVID CICILLINE** recently announced that eight community health centers have received a total of more than \$6.1 million from the American Rescue Plan to make capital improvements aimed at better serving Rhode Island patients.

The funding, which was awarded by the Health Resources and Services Administration, will be used to expand, renovate, and make other capital improvements at community health centers across the state. The awards will support the improvement of primary care infrastructure in Rhode Island and advance health equity and better health outcomes in medically underserved communities, including through projects that support COVID-19 testing, treatment, and vaccination.

The health centers receiving funding are:

- Blackstone Valley Community Health Care, Pawtucket – \$734,663
- Comprehensive Community Action, Cranston – \$677,672
- East Bay Community Action, Newport – \$616,578
- Northwest Community Health Care, Pascoag – \$682,171
- Providence Community Health Centers, Providence, \$1,166,666
- Thundermist Health Center, Woonsocket – \$1,065,455
- Tri-County Community Action Agency, Johnston – \$589,111
- Wood River Health Services, Hope Valley – \$573,876

Rhode Island community health centers provide care for about 190,000 patients, according to the National Association of Community Health Centers. ❖

Miriam receives \$11.1M grant to fund COBRE center on stress, trauma in early life

PROVIDENCE – The Miriam Hospital has received an \$11.1 million federal grant to establish a Center of Biomedical Research Excellence (COBRE) in Rhode Island devoted to a growing field of inquiry – how stress and trauma early in life can have lasting impacts on our health and wellness.

The grant from the National Institute of General Medical Sciences, part of the National Institutes of Health, will fund the creation of the STAR COBRE, which will be based out of the Center for Behavioral and Preventive Medicine at The Miriam Hospital. STAR stands for Stress, Trauma, and Resilience.

The center will build research capacity at The Miriam while also establishing partnerships with collaborators at Lifespan, the Warren Alpert Medical School of Brown University, and Care New England, specifically its psychiatric facility, Butler Hospital. Study topics will include the effects that child neglect, sexual abuse, and food insecurity have on mental and physical well-being.

The COBRE grant is for five years and is the first phase of what could ultimately span three phases over 15 years. These centers are intended to bring together experienced researchers and junior investigators to cultivate multidisciplinary research and expertise in a variety of fields. In recent years, Lifespan has succeeded in securing grants for COBRE centers devoted to a variety of critical areas of research including the following: antimicrobial resistance, opioids and overdose, childhood and adolescent sleep, skeletal health and repair, cancer research development, and stems cells and aging.

“It is a critical time for research into stress and trauma, and the pathways to resilience. The COVID-19 pandemic

has increased the global burden of stress, trauma, and adversity and has exaggerated racial, ethnic, and socio-economic inequalities. The STAR COBRE will be the first research center in Rhode Island to focus on stress, trauma, and resilience,” said the center’s principal investigator, **LAURA R. STROUD, PhD**, who is director of The Miriam’s Center for Behavioral and Preventive Medicine and a professor of psychiatry and human behavior at the Warren Alpert Medical School of Brown University. “It will support transformative research to understand how stress and trauma impact mental and physical health and develop novel approaches to interventions that will promote resilience across the lifespan.”

Proposed research initiatives at the STAR center, led by early career investigators, include:

- A project investigating how childhood maltreatment impacts executive function, rumination and mental health symptoms in adolescents building on a 12-year prospective study. (**CRYSTAL VERGARA-LOPEZ, PhD**)
- A project using real time experience sampling in pregnant women to examine links between childhood sexual abuse and hypertensive disorders of pregnancy. (**MARGARET BUBLITZ, PhD**)
- A clinical trial to examine the impact of food insecurity on diet, inflammatory and metabolic markers, and summer weight gain. (**E. WHITNEY EVANS, PhD**)

The Center will also allow The Miriam Hospital to recruit a new faculty member, and to fund pilot projects for new research initiatives with a particular focus on health disparities. ❖


Ortho Rhode Island now offering robotic technology in total and partial joint replacement at Ortho RI Surgery Center in Warwick

WARWICK – Ortho Rhode Island surgeon **DR. ROBERT MARCHAND** performed a robotic-assisted total joint replacement at the Ortho RI Surgery Center in Warwick, the first procedure of its kind at a freestanding ambulatory surgery center in Rhode Island and Massachusetts. For over a decade, Dr. Marchand and Ortho Rhode Island surgeons have pioneered the use of robotic-assisted technology for hip, knee, and spine procedures in the state, in partnership with South County Health.

Dr. Marchand operated with Stryker's Mako SmartRobotics™, which combines three key components – 3D CT-based planning, Accu Stop™ haptic technology, and insightful data analytics – into one platform that has shown better outcomes for total knee, total hip, and partial knee patients.

“The Ortho RI Surgery Center offering robotics is a key demonstration of Ortho Rhode Island’s commitment to provide the community with outstanding healthcare, and we are proud to continue to be at the forefront of innovation and technology,” said **MICHAEL P. BRADLEY**, President and CEO of Ortho Rhode Island.

Patients medically eligible for outpatient total joint procedures at the Ortho RI Surgery Center return home the same



day as their surgery, while more complex inpatient robotics cases are performed by Ortho Rhode Island surgeons at local community hospitals, including South County Health.

Dr. Marchand has performed thousands of robotic-assisted joint replacement surgeries with Stryker's technology, and uses his expertise to teach surgeons around the globe how to implement this state-of-the-art innovation. ❖

Appointments

Mukesh K. Jain, MD, appointed Dean of Medicine and Biological Sciences at Brown

PROVIDENCE [BROWN UNIVERSITY] — **DR. MUKESH K. JAIN** has been appointed dean of medicine and biological sciences at Brown University. Jain serves currently as chief academic officer at University Hospitals health system in Cleveland and vice dean for medical sciences at Case Western Reserve University.

In joining Brown, he will lead the Division of Biology and Medicine, which encompasses the Warren Alpert Medical School, four biological science departments, 14 clinical departments and two hybrid departments. He will also manage the medical school's relationships with nine affiliated hospitals. Key responsibilities include supporting expansion of research activity across the Division of Biology and Medicine, with a vision to advance Brown's position in the top tier of medical schools and basic science and clinical departments. Integral to his role will be sustaining a culture of ongoing engagement with learners at every level, and promoting and enhancing diversity, equity and inclusion among all members of the community in the execution of education, research and clinical care missions.

Jain said that a range of factors piqued his interest in Brown, including the Warren Alpert Medical School's reputation for outstanding contributions to science and medicine and track record in training top physicians. He's eager to partner with faculty at Brown's School of Public Health and Carney Institute for Brain Science as well as colleagues at other schools, institutes and departments on campus. And the potential for greater integration in medical care, research and education in Rhode Island also played a significant role in his decision.

"I hope to leverage my experiences in building robust basic and translational research programs that are aligned and integrated with excellence in clinical care to help the Warren Alpert Medical School

and its clinical affiliates achieve even greater impact in advancing standard of care to improve human health," he said. "And across all biomedical programs, I will work to ensure that Brown is recruiting and retaining the best and brightest from all backgrounds and is at the tip of the national spear in cultivating the next generation of physicians and scientists."

Increasing the representation of students and faculty from groups historically underrepresented in medicine and science will be a key priority for the dean, along with advancing efforts to ensure that medical school graduates deliver care to patients with a full understanding of social determinants of health and commitment to improving upon existing inequities in health care.

In addition to his leadership roles in Cleveland, Jain is a biomedical researcher internationally recognized for studies that established a central role for a family of genetic factors in cardiovascular biology, innate immunity and metabolism. He is the founding director of Case Western Reserve's Cardiovascular Research Institute, which established a robust cardiovascular research program that spanned basic and clinical research, secured over \$60 million in research funding and is home to nearly 120 clinical trials involving 2,500 patients.

Since 2012, Jain has served as the chief scientific officer at the Harrington Discovery Institute, which supports breakthrough biomedical research. During his tenure, the institute has provided more than \$6 million annually to scholars across North America and the United Kingdom, as well as support to physician-scientists in the Cleveland community. In 2019, he was appointed a visiting professor at the University of Oxford in the U.K.

Nationally, Jain's clinical and academic contributions have been recognized through numerous honors including

election to the National Academy of Medicine, the American Society for Clinical Investigation – an

organization for which he is a past president – the Association of American Physicians, and the Association of University Cardiologists. In 2017, Jain and five colleagues (including two Nobel laureates) founded the Physician-Scientist Support Foundation, a nonprofit that strives to build a sustainable and diverse physician-scientist workforce through financial and research support for fundamental discoveries that improve human health.

Jain earned a bachelor's degree in biochemistry from the University of Buffalo and his MD from the University of Buffalo School of Medicine. After finishing his residency in internal medicine at the Beth Israel Deaconess Medical Center in Boston, he completed fellowships in research and cardiovascular disease at Harvard School of Public Health and Brigham and Women's Hospital.

Jain's appointment follows a comprehensive search process led by a committee that included faculty, staff and administrators as well as student representatives from the Warren Alpert Medical School and Brown's Graduate School. He will begin as dean effective March 1, 2022, serving as a member of the Provost's Senior Academic Deans committee and of the President's Cabinet.

Jain will succeed Dr. Jack A. Elias, Brown's dean of medicine and biological sciences for eight years, who is transitioning to the role of senior advisor for health affairs and working with University leadership to advance the development of Brown's relationships with Lifespan and Care New England as efforts toward an integrated academic health system continue. ❖



Appointments



Mark Deitch, MD, named VP, CMO at Miriam

PROVIDENCE – **MARK DEITCH, MD, MBA**, is the new senior vice president of medical affairs and chief medical officer at The Miriam Hospital.

Deitch is a veteran health care administrator who comes to Lifespan from WellSpan Health, an integrated nonprofit health system serving South-Central Pennsylvania and Northern Maryland. He served as vice president of its orthopedic service line and was responsible for strategy, standardization, and growth of orthopedic services. Previously, he held positions as an assistant professor at Johns Hopkins Medical Institute and University of Maryland Medical System, president of Greenspring Surgery Center, partner/attending orthopedic surgeon with OrthoMaryland, and owner/president of MD3 Orthopedics, all of Baltimore.

Deitch is a graduate of the Stony Brook University School of Medicine and earned his MBA in medical services management from Johns Hopkins University. He is a member of the American Academy of Orthopedic Surgeons and the American Society for Surgery of the Hand.

This is a return to Rhode Island for Dr. Deitch who completed his bachelor of arts degree in biology with honors in research at Brown University. He was a fellow in hand, microvascular and upper extremity surgery at Cincinnati Hand Surgery Specialists, Ohio; a resident in orthopedic surgery at the Hospital for Special Surgery, New York; and an intern in general surgery at Beth Israel Deaconess Medical Center, Massachusetts.

Deitch, who arrived at The Miriam in September, succeeds G. Dean Roye, MD, FACS, who held the position since 2017 and now serves as senior vice president of medical affairs and chief medical officer at Rhode Island and Hasbro Children's hospitals. ❖



Megan Ranney, MD, appointed Academic Dean at Brown School of Public Health

PROVIDENCE – **MEGAN RANNEY, MD, MPH, FACEP**, has been appointed as the Brown School of Public Health's inaugural Academic Dean, effective December 2021. Reporting directly to the Dean, she will serve as the chief academic officer of the School, and a key

member of the leadership team.

As chief academic officer of the School, Professor Ranney will work in partnership with Associate Deans Melissa Clark and Jennifer Tidey, as well as with Director of Community Engagement, Kim Gans. She will collaborate with them to help oversee the School's educational efforts, research enterprise, community engagement, and faculty recruitment, development, evaluation, retention, and promotion. She will also partner with the rest of the Deans, including Interim Associate Dean for Diversity and Inclusion Ronald Aubert, Executive Dean of Finance and Administration Sara Walsh, and Vice Dean of Advancement Benjamin Zoll.

Professor Ranney initially joined the School's leadership as Associate Dean for Strategy and Innovation earlier this year, providing expertise to help develop our expanding portfolio strategy and further connect our School within the University.

As part of this new position, Professor Ranney will also be appointed Professor of Behavioral and Social Sciences, continuing to serve as an educator, researcher, and mentor to so many at Brown. ❖

RIMJ Seeks Volunteer Physician Editor

The *Rhode Island Medical Journal* (RIMJ) is seeking a Rhode Island physician with an interest and expertise in editing and writing to assist with issue planning, theme development, peer review, and author correspondence.

This volunteer editorial position, with a flexible time commitment, is a unique opportunity for a physician familiar with the State's healthcare landscape to engage with colleagues, faculty, students and allied healthcare professionals.

RIMJ publishes 10 online issues a year (available at rimedj.org), with a hiatus in January and July. The publisher is the Rhode Island Medical Society (RIMS), and editors are approved by the RIMS Board for a three-year tenure, with an option to renew for an additional term.

Interested candidates should submit a cover letter and CV to **Editor-in-Chief William Binder, MD**, at william_binder@brown.edu.

Appointments

Kyle S. Nuland, MD, joins University Orthopedics Adult Reconstruction Division

EAST PROVIDENCE – University Orthopedics is expanding The Joint Replacement Center with the addition of **KYLE S. NULAND, MD**.

Dr. Nuland specializes in hip and knee arthroplasty, various hip and knee disorders, minimally invasive knee surgery, orthopedic trauma, knee and hip reconstruction, and total hip revision.

Dr. Nuland recently completed the Frank E. Stinchfield Adult Reconstruction fellowship at Columbia University in New York City. Prior to that, he attended medical school at the University of Connecticut and completed the residency program at Albert Einstein College of Medicine-Montefiore Medical Center in Bronx, NY. ❖



Ross Budacki, MD, joins University Orthopedics Sports Medicine division

EAST PROVIDENCE – University Orthopedics recently announced board-certified orthopedic surgeon **ROSS BUDACKI, MD**, is joining its Sports Medicine division.

Dr. Budacki is a fellowship-trained sports medicine surgeon who treats patients with a wide variety of injuries from simple sprains and strains

all the way up to complex, traumatic injuries involving the shoulder, knee, hip, hand, wrist, and elbow. He also serves as clinical assistant professor orthopaedics at the Warren Alpert Medical School of Brown University.

Dr. Budacki completed a sports medicine fellowship at the Alpert Medical School. Prior to that, he earned his medical degree at Georgetown University School of Medicine and completed the residency program at Einstein Healthcare Network in Philadelphia. During his fellowship, he served as the assistant team physician for Brown University's Division I football team and the Providence Bruins. Dr. Budacki is a member of the American Academy of Orthopaedic Surgery and the American Orthopaedic Society for Sports Medicine. ❖

Retina Consultants

Altin Pani, MD

is pleased to announce that



Scott H. Greenberg, MD

has joined the practice specializing in diseases and surgery of the retina and vitreous.

Dr. Greenberg is a graduate of Upstate Medical University in Syracuse, New York. He completed a residency in ophthalmology and a fellowship in vitreoretinal surgery at Albany Medical College in New York.

Referrals welcome

Providence

401-274-5844

North Kingstown

401-294-9429

RetinaConsultantsRI.com

Obituaries



LOUIS A. CORVESE, MD, a retired orthopedic surgeon, died peacefully on October 16, 2021 following a brief illness. He was the son of the late Dr. Anthony and Mary Leoni Corvese and the brother of the late Dr. William P. Corvese.



Dr. Corvese graduated from Classical High School, the Catholic University of America, and in 1956 he graduated from the Yale University School of Medicine. He interned at George Washington University Hospital in Washington, DC, and further trained at Newington Children's Hospital and Yale University. He was a fellow of the American Academy of Orthopedic Surgeons and was certified by the American Board of Orthopedic Surgery.

Dr. Corvese practiced orthopedic surgery for more than forty years and was a member of the RI Hospital surgical staff. Kindness and concern for his patients was always paramount to him.

Earlier in his career, Dr. Corvese served as a surgeon on active duty at sea and ashore with the US Navy. Among the Navy vessels he served on during the war in Vietnam were the USS Enterprise and the USS Sanctuary, the Navy hospital ship anchored off Vietnam providing medical and surgical care for wounded soldiers.

Dr. Corvese generously supported the arts and charities throughout his life. He was very grateful for the many acts of kindness and concern shown to him by family and friends, especially during the last years of his life. ❖



ZHOU WENG, MD, 93, died on October 17, 2021 at HopeHealth Hular Hospice Center, Providence. He was the beloved husband of the late Li-Li Lin.

Born in China, he moved to the United States in 1992, settling in Pawtucket. Dr. Weng was a general surgeon in China before retiring. He was a member of the Chinese Christian Church of Rhode Island.

He is survived by two daughters, Liya Beaujean of New York, and Kathy Zheng of Tiverton; two sons, Bin Weng of Barrington and John Weng of New York; four sisters, Xiu Zhen Weng, Ying Weng, Yuan Weng and Bing Weng; two brothers, Huan Weng and He Weng; eight grandchildren; and many nieces and nephews. ❖